Annual burning of semi-natural grasslands for conservation favours tall-grown species with high nectar production

Per Milberg, Håkan Fogelfors, Lars Westerberg and Malin Tälle

The self-archived postprint version of this journal article is available at Linköping University Institutional Repository (DiVA):

<http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-149385>

N.B.: When citing this work, cite the original publication.

Milberg, P., Fogelfors, H., Westerberg, L., Tälle, M., (2018), Annual burning of semi-natural grasslands for conservation favours tall-grown species with high nectar production, *Nordic Journal of Botany*, 36(5), UNSP e01709. <https://doi.org/10.1111/njb.01709>

Original publication available at:

<https://doi.org/10.1111/njb.01709>

Copyright: Wiley (12 months)

<http://eu.wiley.com/WileyCDA/>

Annual burning of semi-natural grasslands for conservation: winners and losers among plant species

Per Milberg, Håkan Fogelfors, Lars Westerberg, Malin Tälle

Milberg, P. (corresponding author per.milberg@liu.se), **Tälle, M.** (malin.talle@liu.se), **Westerberg, L.** (lars.westerberg@liu.se): IFM Biology, Conservation Ecology Group, Linköping University, SE-581 83 Linköping, Sweden

Fogelfors, H. (hakan.fogelfors@slu.se): Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Box 7043, SE-750 07 Uppsala, Sweden

Abstract

Species-rich semi-natural grasslands in Europe are a main target for conservation efforts, and alternative methods to the traditional management of mowing or grazing would be welcome due to the difficulties in maintaining these management practices. One such method proposed is burning of grassland vegetation during late winter or spring. To evaluate the effects of annual spring burning vs. annual mowing on semi-natural grassland vegetation, we compared the frequency of species in eleven field experiments in southern Sweden after ca 14 years. Out of the 88 species analysed, five were more frequent in burnt plots compared with mowed plots (*Vicia cracca*, *Cirsium arvense*, *Urtica dioica*, *Galium verum*, *Convallaria majalis*). In contrast, 37 species were significantly less frequent in burnt plots compared with mowed ones, those with the largest differences being *Ranunculus acris*, *Briza media*, *Veronica chamaedrys*, *Festuca ovina*, *Plantago lanceolata* and *Anthoxanthum odoratum*. Tall-grown species and those with preferences for N-rich soils increased in frequency under an annual spring-burn regime, compared with annual mowing, as did species producing larger amounts of nectar. Hence, although vegetation composition becomes trivial with annual spring burns, there might be long-term benefits for nectar-feeding insects.

Keywords: annual fire, Management, mowing, Semi-natural grassland, Sweden

Introduction

The conservation of species-rich grasslands is a demanding task in Europe and elsewhere where fodder has traditionally been extracted from native vegetation (Poschlod et al. 2009). In most agricultural systems, such fodder – both pasturage and hay – is no longer economically viable. Therefore, management of many species-rich grasslands is mainly maintained, for biodiversity and historical reasons, through subsidies to farmers, but also by nature conservation authorities and non-government organisations. The costs for mowing (Schreiber et al. 2009, Török et al. 2011) and the increasing difficulty of finding grazers for conservation (Kumm 2003) means that other ways to maintain biodiversity-rich grassland systems would be welcome. The search for alternative grassland management methods is by no means new: field experiments evaluating such methods were set up as early as the 1970s (e.g. Sweden: Hansson 1991; Germany: Schreiber et al. 2009; Switzerland: Köhler et al. 2005). Any new method is likely to exert a selection pressure that is different from that of traditional management (e.g. Alhamad et al. 2012; Jing et al. 2017) and effects of such a method must therefore be investigated before it can be recommended, e.g. as some conservation-targeted species might benefit while others might decline under new management.

Fire, a disturbance in many grass-dominated ecosystems (Brockway et al. 2002; van Langevelde et al. 2003; Archibald et al. 2005), is a potential management method that has received recent attention (e.g. Valko et al. 2014, 2016, Koyama et al. 2017). Fire is sometimes used as a restoration method, i.e. one or a couple of burns to reduce litter and eliminate woody species in semi-natural grasslands (e.g. Valko et al. 2016, Ruprecht et al. 2016, Pereira et al. 2016). However, annual burning of grass litter in late winter or early spring, before the onset of grass growth, as a substitute for mowing or grazing, serves a completely different purpose to restoration, with potentially stronger effects on biodiversity composition. Results from longer-term experiments examining effects of burning of semi-natural grasslands are negative when it comes to vegetation (Lunt & Morgan 1999, Kahmen et al. 2002, Moog et al. 2002, Köhler et al. 2005, Milberg et al. 2014, Milberg & Bergman 2014). Still, some parts of biodiversity might benefit (Deak et al. 2014, Vegvari et al. 2016), and therefore the method is likely to be used in several parts of Europe. As spring-burning affects vegetation composition (Wahlman & Milberg 2002), organisms that depend on plants is also likely to be affected. Nectar-feeding insects is a functional group with particular value due to the pollination service they provide, and these have been reported to

benefit from spring burning in a short-term Swedish experiment (Larsson & Persson 2012). In the short-term, the degree of disturbance during the growth season is likely to be negatively related to the abundance of pollinators by reducing the number of flowers (Milberg et al. 2016). Hence, spring-burning should therefore be expected to be better than mowing or grazing simply due to higher abundance of flowers present over longer time. However, in the long-term, and with a shift in vegetation composition, it is less clear what to expect regarding pollinators.

In the present study, we set out to document how plant species were affected when subjected to annual spring burns, using annual mowing as control. We used eleven Swedish field experiments, evaluated after ca 14 years of management. Results from these experiments have previously been compiled to compare burning with annual mowing and grazing, then evaluating the proportion of indicator species in vegetation and without any attention to species-wise results (Milberg et al. 2014). In the present contribution, we focus on species-wise responses, and more specifically, evaluate the following seven hypotheses regarding plant species attributes. The purpose was to identify possible general patterns in grassland vegetation:

- (i) graminoids would be promoted under annual burning, compared with forbs, as the former are often considered as fire-adapted (Bowman et al. 2014);
- (ii) lianas (winding plants), that invest relatively little in structural support, would have an advantage in tall-grown vegetation that is left undisturbed during the growth season (McIntyre & Lavorel 2001, Saatkamp et al. 2010);
- (iii) therophytes (annuals) would decrease due to problems involving seed regeneration and competition in the tall-grown vegetation developing under a spring burn regime (e.g. Jakobsson & Eriksson 2000, Lennartsson & Oostermeijer 2001).
- (iv) N-fixing species would be promoted under annual burning as nitrogen is lost from the vegetation due to spring-burns (e.g. Hobbs et al. 1991), giving them a competitive advantage;
- (v) species with low Ellenberg N indicator values would be promoted under annual burning as nitrogen is lost from the vegetation due to spring-burns (e.g. Hobbs et al. 1991);
- (vi) tall-grown species would be promoted under annual burning as they benefit from the lack of disturbance during the growth season (e.g. Borer et al. 2014, Milberg et al. 2017);

- (vii) there is no relationship between the nectar production of species and their response to annual spring burning (cf Moranz et al. 2012).

Methods

Study sites

This experiment was established in the 1970's and involved eleven experiments at nine locations in southern Sweden (Figure 1). Mean annual precipitation in this area varies between 500 mm (east) and 1000 mm (west). The mean annual temperature is around 6 °C (Alexandersson et al. 1991) while the growing season is 180-220 days (Sjörs 1999).

The experimental sites were selected to represent a diversity in soil type and land use history of marginal grasslands which – in the early 1970s – were at risk of abandonment and becoming overgrown (Steen 1976). Before the start of the experiments, most of the sites were grazed but one site (Dämkärr) had not been managed for three years, and one site (Gränö) had been mowed for a decade. Furthermore, two of the sites (Gränö and Tagel/former field) had a history as fertilized arable field (up until ca. 10 and 20 years prior to onset of the experiments, respectively) (Table 1, Hansson 1991). These former, small arable fields were embedded in old-style, small-scale agricultural landscapes, subject to some degree of alteration between mowing and arable (Ekstam & Forshed 1996).

More extensive descriptions of the experiments can be found in Steen (1976), Fogelfors (1982), Hansson (1991). It is also worth noting previous papers evaluating one of these experiments (Hansson & Fogelfors 2000, Wahlman & Milberg 2002), or all of them (Milberg et al. 2014, 2017, Tälle et al. 2015).

Experimental design

The experiment involved several management methods but in the present study we compared two of them: annual summer mowing vs. annual spring burning. We choose mowing since grazing intensity was low in some of the experiments (Hansson 1991). The management methods were applied to 5 × 20 m² treatment plots. Treatment plots were set within a randomized block design with two replicates. All plots were fenced. Mowing took place in late July or early August, using a scythe or sickle bar mower. Spring burning was executed in early April, before the onset of growth, when litter from previous year was flammable. The

exact dates varying between sites and years due to weather and snow conditions.

Vegetation sampling

Vegetation sampling was conducted in the summer before mowing, i.e. in July. We used data collected in 1986 (10 experiments) and 1987 (1 experiment), when five 1 m² fixed subplots were used for all experiments except one (Bräcke) where three 1 m² fixed subplots were used. For the surveys, the coverage of vascular plant species was recorded, but we decided to use presence/absence in subplots for the current analyses. Nomenclature follows Karlsson (1998).

Plant species attributes

Species were classified as nectar-producing or non-nectar producing based on Baude et al. (2016). In addition to this classification, Baude et al. (2016) also contain published nectar production estimates per area (kg per ha cover per year) for 39 of our 88 species; some data were empirical and some modelled (Baude et al. 2016).

Four of the species attributes were extracted from the LEDA database (Kleyer et al. 2008): canopy height (continuous); Ellenberg N (1-9 reflecting the preference of a species to productive sites rich in soil N); growth form (liana, therophyte, other). N-fixation ability was assigned according to literature (Werner et al. 2014) to species of Fabaceae. The classification of forbs vs. graminoids (Poaceae, *Carex*, *Juncus*, *Luzula*) were made according to the plant family.

Statistical analyses

Each of the 11 experiments had two blocks, and we used the difference in frequency of a species within a block for the analyses, hence N = 22 (11 experiments * 2 blocks). No adjustment was made for one site (Bräcke) having three rather than five subplots, nor for the fact that one experiment was assessed in a different year compared with the other. In total, 235 species had been recorded in these 22 blocks. As expected, many of the species were rare in data and therefore ill-suited for analyses. We decided to use only those 88 species that were deemed as sufficiently frequent for meaningful analyses (cutoff at the occurrence in at least 8 of the 22 blocks).

A weighted average of the block-wise frequency-difference data (burning-mowing) were calculated using the software Comprehensive Meta-Analysis version 2 (www.meta-analysis.com, Borenstein et al. 2009) applying the random effects model (i.e. species identity was

considered a random factor). This analysis platform was also used to group the 88 species according to life-history traits, and to apply meta-regression in case of a continuous explanatory variable.

Results

Among the 88 species, five (6%) displayed significant, positive change in response to repeated spring burns (Figure 2; positive estimate with CI not overlapping zero), hence were promoted when subjected to this treatment compared to annual mowing (*Vicia cracca*, *Cirsium arvense*, *Urtica dioica*, *Galium verum*, *Convallaria majalis*). In contrast, 37 species (42%) were disfavoured by spring burning (Figure 2), many of which were typical of traditionally managed grasslands (e.g. *Rhinanthus minor*, *Primula veris*, *Rumex acetosa*, *Luzula multiflora*, *Leucanthemum vulgare*, *Lathyrus linifolius*, *Briza media*, *Anthoxanthum odoratum*). In addition, the estimates for a large number of the species leaned towards higher occurrences in annual mowing than annual burning (Figure 2).

Hypotheses regarding species attributes

Graminoids and forbs exhibited similar responses (Figure 2b).

Nitrogen-fixing plant species did not differ from those lacking this attribute in their response to repeated spring-burns (Figure 2b).

Lianas did well under a spring-burning regime while therophytes did not, compared to the rest of the species (Figure 2b). It is worth pointing out that both groups were small ($N = 3$ and $N = 2$, respectively).

When comparing annual burning with annual mowing, there was no difference in frequencies for species that produce nectar (-0.88; CI -1.257; -0.497) and those that do not (-0.77; CI -1.159; -0.386). However, there was a clear relationship among the 39 nectar-producing species: species with large nectar production tended to be promoted by spring burns (meta-regression slope 0.236, CI: 0.134; 0.338, $P < 0.001$, Figure 3a).

There was a positive relationship between a species' height and its response to spring-burning (slope in meta-regression 0.020; CI 0.014; 0.025; $P < 0.0001$; Figure 3b).

There was a positive relationship between a species' Ellenberg N-value and its response to spring-burning (meta-regression slope 0.164; CI: 0.236; 4.514; $P < 0.001$), contrary to the expected negative relationship.

Discussion

Annual spring burning had a detrimental effect on most plant species, as was expected from previous evaluation of these data (Milberg et al. 2014) and other studies (Moog et al. 2002, Köhler et al. 2005). Many of the species disfavoured by annual spring burns are considered indicator species for traditional management (Ekstam & Forshed 1992), while most among those favoured are of low interest for conservation of plant species. Hence, from the point of view of preserving plant species associated with traditional management, annual burning does not provide much benefit. However, spring-burning does control trees and shrubs (Hansson & Fogelfors 2000, Wahlman & Milberg 2002) and reduce the amount of litter (Ryser et al. 1995, Liira et al. 2009). Furthermore, our study showed that grassland vegetation shaped by spring-burning might benefit the occurrence of nectar-feeding insects by promoting the more prolific nectar-producers (Figure 3a). Considering the importance of the pollination service nectar-feeding insects provide to society (Klein et al. 2007, Ollerton et al. 2011), one might see value in a habitat created by spring-burning. It is worth noting that this is an additional advantage, through compositional changes, to the expected abundance of flowers caused by the absence of disturbance during the growth season (Erhardt 1985, Smith & Cherry 2014).

Treatments and species' responses

A new management method introduces a new selection pressure and we tested a number of plant attributes expected to be affected by annual burning compared with annual mowing.

Graminoids (Poaceae, *Carex*, *Juncus* and *Luzula*) did not differ from forbs in their response to annual spring burns. Thus, we found no support for our hypothesis that graminoids would benefit from spring burns. It is worth pointing out that many of the non-grasses in this group (i.e. *Carex*, *Juncus* and *Luzula*) were short in height, and therefore might do poorly under an annual spring-burn regime (see further below). However, excluding these three genera did not change the results (data not shown). It is worth noting that grass biomass is very unevenly distributed among species, so other types of endpoints (e.g. biomass) might still point out grasses as winners.

Lianas did well under spring-burning and therophytes did not, i.e. they responded according to our assumptions. Although both groups were small, the findings are in line with the literature: climbing plants often do well at low levels of disturbance (Saatkamp et al. 2010). In contrast,

annuals face difficulty in tall swards characteristic of such disturbance regimes (Lennartsson & Oostermeijer 2001).

Under the expectation that N would be lost during a spring-burn (Hobbs et al. 1991), we hypothesized (i) that N-fixing species would benefit from an annual spring burn regime, and (ii) that there would be a negative relationship between the Ellenberg N-value of a species and its response to an annual spring burn regime. Both assumptions were refuted, suggesting that either the amounts of N lost when burning dry litter in the spring is too small to exert a detectable effect (e.g. due to a large pool of N in soil), or that there are other changes in vegetation that hides such an effect. For example, in the present data, there was a positive relationship between the Ellenberg N of species and their canopy height ($R = 0.499$; $N = 82$; $P < 0.001$). Hence, species of more productive sites tended to be more tall-grown, and a positive canopy height effect might therefore eliminate a negative Ellenberg N effect. A third alternative is that repeated burns may lead to increase in plant-available soil N, something reported after low-intensity fires (e.g. Neary et al. 1999; Augustine et al. 2014).

As expected, tall-grown species benefitted from repeated spring-burns, most likely due to the lack of disturbance during the growth season allowing tall-grown species to develop fully, while short-grown species benefitted from annual mowing (e.g. Liira and Zobel 2000; Milberg et al. 2017).

Overall, it seems that tall-grown, N-demanding species and those producing much nectar are winners in spring-burned grasslands, while there were weak or no support for any of the other species attributes evaluated.

Spring burning vs no treatment?

It is likely that a part of the response to burning reported can be attributed to lack of management during the growth season, which similarly affects species according to their canopy height (Milberg et al. 2017). In a recent study using similar methodology and analyzing the same field experiments, Milberg et al. (2017) reported the response to free development (no management) for about half of the species included in the present study. As expected, when comparing effects from the present and the previous study, there was a high correlation between species-wise responses to spring burning and to no treatment ($R = 0.500$, $N = 44$). By using a regression, two species (*Centaurea jacea* and *Vicia cracca*) showed residuals that were > 2 SD higher than expected by chance during free development. Other species that did much better under burning than

expected from their response to free development (> 1 SD) were *Carex pallescens*, *Galium boreale*, *Galium verum* and *Hypericum maculatum*. Two and four species had negative residuals > 2 SD (*Anemone nemorosa*, *Veronica chamaedrys*) and > 1 SD (*Anthoxanthum odoratum*, *Lathyrus linifolius*, *Plantago lanceolata* and *Ranunculus acris*), respectively. Hence, although part of the response in the present study is attributable to the lack of biomass removal during the growth season, there are species that clearly suffer from a spring burn. Whether this is due to the location of over-wintering buds (Clarke et al. 2013, Russel et al. 2013) or the phenology of growth (Henderson 1990a, b) remains to be evaluated. At the same time, there are also species that clearly benefited from repeated spring burns more than they would if left to free development. The potential mechanism is open to speculation.

Conclusions

If the aim is to maintain a traditional vegetation type, then annual spring burning cannot be recommended over annual mowing. On the other hand, if nectar-production is in focus, and a vegetation with tall-grown species with high Ellenberg N is acceptable, then repeated spring burns has potential merit. This habitat should be considered as a new, unique type that has not existed previously, hence a “neo-habitat” (sensu Blixt et al. 2015). It is worth noting that in two field trials (Köhler et al. 2005, Milberg et al. 2014), vegetation in burnt plots has diverged over time due to the invasion of undesired species. Hence, as a management method, some care would be needed to monitor spring-burnt sites for undesired development.

Acknowledgements

We thank all those involved in initiating, maintaining and monitoring the long-term field experiments analysed here. The Swedish Environmental Protection Agency funded these experiments while the Swedish Board of Agriculture funded the present analyses. We thank anonymous referees for their input.

References

- Alhamad, M.N., Alrababah, M.A. and Gharaibeh, M.A. 2012. Impact of burning and fertilization on dry Mediterranean grassland productivity and diversity. – *Acta Oecol.* **40**: 19-26.
- Alexandersson, H., Karlström, C. and Larsson-McLann, S. 1991. Temperaturen och nederbörden i Sverige 1961-1990. Referensnormaler. –

Sveriges Meteorologiska och Hydrologiska Institut, SMHI Meteorologi
No. 81. 87 pp. In Swedish.

Archibald, S., Bond, W.J., Stock, W.D. and Fairbanks, D.H.K. 2005.
Shaping the landscape: fire-grazer interactions in an African savanna. –
Ecol. Appl. **15**: 96–109.

Augustine, D.J., Brewer, P., Blumenthal, D.M., Derner, J.D. and von
Fischer, J.C. 2014. Prescribed fire, soil inorganic nitrogen dynamics, and
plant responses in a semiarid grassland. – *J. Arid Environ.* **104**: 59–66.

Baude, M., Kunin, W.E., Boatman, N.D., Conyers, S., Davies, N.,
Gillespie, M.A.K., Morton, R.D., Smart, S.M. and Memmott, J. 2016.
Historical nectar assessment reveals the fall and rise of floral resources in
Britain. – *Nature* **530**: 85–88.

Blixt, T., Bergman, K.-O., Milberg, P., Westerberg, L. and Jonason, D.
2015. Clear-cuts in production forests are not matrix, but neo-habitats for
butterflies. – *Acta Oecol.* **69**: 71–77.

Borenstein, M., Hedges, L.V., Higgins, J.P.T. and Rothstein, H. 2009.
Introduction to Meta-Analysis. John Wiley & Sons, Ltd, Chichester, UK.

Borer, E.T., Seabloom, E.W., Gruner, D.S., Harpole, W.S., Hillebrand,
H., Lind, E.M., et al. 2014. Herbivores and nutrients control grassland
plant diversity via light limitation. – *Nature* **508**: 517–520.

Bowman, D.M., French, B.J. and Prior, L.D. 2014. Have plants evolved
to self-immolate? – *Frontiers in Plant Science* **5**: 590.

Brockway, D.G., Gatewood, R.G. and Paris, R.B. 2002. Restoring fire as
an ecological process in shortgrass prairie ecosystems: initial effects of
prescribed burning during the dormant and growing seasons. – *J.*
Environ. Manage. **65**: 135–52.

Clarke PJ, Lawes MJ, Midgley JJ, Lamont BB, Ojeda F, Burrows GE,
Enright NJ, Knox KJE. 2013. Resprouting as a key functional trait: how
buds, protection and resources drive persistence after fire. – *New Phytol.*
197: 19–35.

Deák, B., Valkó, O., Török, P., Végvári, Z., Hartel, T., Schmotzer, A.,
Kapocsi, I. and Tóthmérész, B. 2014. Grassland fires in Hungary–
experiences of nature conservationists on the effects of fire on
biodiversity. – *Appl. Ecol. Environm. Res.* **12**: 267–283.

Ekstam, U. and Forshed, N. 1992. Om hävden upphör; kärlväxter som indikatorarter i ängs- och hagmarker. – Naturvårdsverket Förlag, in Swedish.

Ekstam, U. and Forshed, N. 1996. Äldre fodermarker: betydelsen av hävdregimen i det förgångna, målstyrning, mätning och uppföljning. – Naturvårdsverket Förlag.

Erhardt, A. 1985. Diurnal Lepidoptera: sensitive indicators of cultivated and abandoned grassland. – *J. Appl. Ecol.* **22**: 849-861.

Fogelfors, H. 1982. *Det marginella odlingslandskapets öppethållande. Del II. Resultat och utvärdering av långvariga försök med olika skötselmetoder.* – Swedish University of Agricultural Sciences. Department of Ecology and Environmental Research. Report No. 11. In Swedish.

Hansson, M. 1991. *Management of semi-natural grassland. Results of a fifteen-year-old field experiment in south and central Sweden.* – Swedish University of Agricultural Sciences, Uppsala. Department of Ecology and Environmental Research. Report No. 45. In Swedish with English summary.

Hansson, M. and Fogelfors, H. 2000. Management of semi-natural grassland; results from a 15-year old experiment in southern Sweden. – *J. Veg. Sci.* **11**: 31-38.

Henderson, R.A. 1990a. Effects of spring fire timing on pasque-flower (*Anemone patens*) flower-bud survival. – *Proc. 12th North Amer. Prairie Conf.*, pp. 5-9.

Henderson, R.A. 1990b. Ten-year response of a Wisconsin prairie remnant to seasonal timing of fire. – *Proc. 12th North Amer. Prairie Conf.*, pp. 121-125.

Hobbs, N.T., Schimel, D.S., Owensby, C.E., and Ojima, D.S. 1991. Fire and grazing in the tallgrass prairie: contingent effects on nitrogen budgets. – *Ecology* **72**: 1374-1382.

Jakobsson, A. and Eriksson, O. 2000. A comparative study of seed number, seed size, seedling size and recruitment in grassland plants. – *Oikos* **88**: 494-502.

Jing, G., Li, W., Yu, K., Ratajczak, Z., Kallenbach, R.L. and Cheng, J. 2017. Effects of fertilization, burning, and grazing on plant community in the long-term fenced grasslands. – *Plant Soil Environ.* **63**: 171-176.

Kahmen, S., Poschlod, P. and Schreiber, K.-F. 2002. Conservation management of calcareous grasslands. Changes in plant species composition and response of functional traits during 25 years. – *Biol. Conserv.* **104**: 319-328.

Karlsson, T. 1998. Förteckning över svenska kärlväxter. [The vascular plants of Sweden – a checklist]. – *Svensk Bot. Tidskr.* **91**: 241-560. In Swedish.

Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C. and Tscharntke, T. 2007. Importance of pollinators in changing landscapes for world crops. – *Proc. R. Soc. B* **274**: 303–313.

Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M., Poschlod, P., Van Groenendael, J.M., Klimeš, L., Klimešová, J. et al. 2008. The LEDA Traitbase: a database of life - history traits of the Northwest European flora. – *J. Ecol.* **96**: 1266-1274.

Köhler, B., Gigon, A., Edwards, P.J., Krüsi, B., Langenauer, R., Lüscher, A. and Ryser, P. 2005. Changes in the species composition and conservation value of limestone grasslands in Northern Switzerland after 22 years of contrasting managements. – *Perspectives Plant Ecol. Evol. Syst.* **7**: 51-67.

Koyama, A., Koyanagi, T.F., Akasaka, M., Takada, M. and Okabe, K. 2017. Combined burning and mowing for restoration of abandoned semi-natural grasslands. – *Appl. Veg. Sci.* **20**: 40-49.

Kumm, K.-I. 2003. Sustainable management of Swedish seminatural pastures with high species diversity. – *J. Nat. Conserv.* **11**: 117-125.

Larsson, K. and Persson, K. 2012. *Naturvårdsbränning i gräsmarker: en jämförande studie av bete, bränning och slåtter 2010–2012*. Rapport, Länsstyrelsen i Jönköpings län. In Swedish.
www.jordbruksverket.se/download/18.1fec19c313cf285924f80004269/Naturv%C3%A5rdsbr%C3%A4nning+slutrapport+2010-2012.pdf

Lennartsson, T. and Oostermeijer, J.G.B. 2001. Demographic variation and population viability in *Gentianella campestris*: effects of grassland management and environmental stochasticity. – *J. Ecol.* **89**: 451-463.

Liira, J. and Zobel, K. 2000. Vertical structure of a species-rich grassland canopy, treated with additional illumination, fertilization and mowing. – *Plant Ecol.* **146**: 183-193.

Liira, J., Issak, M., Jõgar, Ü., Mändoja, M. and Zobel, M. 2009. Restoration management of a floodplain meadow and its cost-effectiveness: the results of a 6-year experiment. – *Ann. Bot. Fennici* **46**: 397–408.

Lunt, I.D. and Morgan, J.W. 1999. Vegetation changes after 10 years of grazing exclusion and intermittent burning in a *Themeda triandra* (Poaceae) grassland reserve in south-eastern Australia. – *Aust. J. Bot.* **47**: 537-552.

McIntyre, S. and Lavorel, S. 2001. Livestock grazing in subtropical pastures: steps in the analysis of attribute response and plant functional types. – *J. Ecol.* **89**: 209-226.

Milberg, P. and Bergman, K.-O. 2014. Vårbränning är inte ett långsiktigt alternativ till bete eller slåtter av värdefulla artrika gräsmarker. – *Svensk Bot. Tidskr.* **108**: 312-322.

Milberg, P., Akoto, B., Bergman, K.-O., Fogelfors, H., Paltto, H. and Tälle, M. 2014. Is spring burning a viable management tool for semi-natural grasslands? – *Appl. Veg. Sci.* **17**: 429-441.

Milberg, P., Bergman, K.-O., Cronvall, E., Eriksson, Å.I., Glimskär, A., Islamovic, A., Jonason, D., Löfqvist, Z. and Westerberg, L. 2016. Flower abundance and vegetation height as predictors for nectar-feeding insect occurrence in Swedish semi-natural grasslands. – *Agric. Ecosystems Environ.* **230**: 47–54.

Milberg, P., Tälle, M., Fogelfors, H. and Westerberg, L. 2017. The biodiversity cost of reducing management intensity in species-rich grasslands: Mowing annually vs. every third year. – *Basic Appl. Ecol.* **22**: 61-74

Moog, D., Poschlod, P., Kahmen, S. and Schreiber, K.F. 2002. Comparison of species composition between different grassland management treatments after 25 years. – *Appl. Veg. Sci.* **5**: 99-106.

Moranz, R.A., Debinski, D.M., McGranahan, D.A., Engle, D.M. and Miller, J.R. 2012. Untangling the effects of fire, grazing, and land-use legacies on grassland butterfly communities. – *Biodiversity Conserv.* **21**: 2719-2746.

Neary, D.G., Klopatek, C.C., DeBano, L.F. and Ffolliott, P.F. 1999. Fire effects on belowground sustainability: a review and synthesis. – *For. Ecol. Manage.* **122**: 51-71.

Ollerton, J., Winfree, R. and Tarrant, S. 2011. How many flowering plants are pollinated by animals? – *Oikos* **120**: 321-326.

Pereira, P., Cerdà, A., Lopez, A.J., Zavala, L.M., Mataix-Solera, J., Arcenegui, V., Misiune, I., Keesstra, S. and Novara, A. 2016. Short-term vegetation recovery after a grassland fire in Lithuania: The effects of fire severity, slope position and aspect. – *Land Degrad. Devel.* **27**: 1523-1534.

Poschlod, P., Baumann, A. and Karlik, P. 2009. Origin and development of grasslands in central Europe. – In: Veen, P., Jeffersson, R., de Smidt, J., and van der Straaten, J. (eds.), *Grasslands in Europe of high nature value*. KNNV Publishing, pp. 12-25.

Ruprecht, E., Enyedi, M.Z., Szabó, A. and Fenesi, A. 2016. Biomass removal by clipping and raking vs burning for the restoration of abandoned *Stipa*-dominated European steppe-like grassland. – *Appl. Veg. Sci.* **19**: 78-88.

Russell, M.L., Vermeire, L.T., Ganguli, A.C. and Hendrickson, J.R. 2015. Season of fire manipulates bud bank dynamics in northern mixed-grass prairie. – *Plant Ecol.* **216**: 835-846.

Ryser, P., Langenauer, R. and Gigon, A. 1995. Species richness and vegetation structure in a limestone grassland after 15 years management with six biomass removal regimes. – *Folia Geobot. Phytotax.* **30**: 157–167.

Saatkamp, A., Römermann, C. and Dutoit, T. 2010. Plant functional traits show non-linear response to grazing. – *Folia Geobot.* **45**: 239-252.

Schreiber, K.-F., Brauckmann, H.-J., Broll, G., Krebs, S. and Poschlod, P. 2009. *Artenreiches Grünland in der Kulturlandschaft. 35 Jahre Offenhaltungsversuche Baden-Württemberg*. – Verlag Regionalkultur, in German.

Sjörs, H. 1999. The background: geology, climate and zonation. – *Acta Phytogeogr. Suecica* **84**: 5-14.

Smith, L.M. and Cherry, R. 2014. Effects of management techniques on grassland butterfly species composition and community structure. – *Amer. Midl. Nat.* **172**: 227-235.

Steen, E. 1976. *Det marginella odlingslandskapets öppethållande. Dokumentation i långvariga försök med olika skötselmetoder.* – Lantbrukshögskolan, Avdelningen för ekologisk miljövard, Uppsala. In Swedish.

Tälle, M., Fogelfors, H., Westerberg, L. and Milberg, P. 2015. The conservation benefit of mowing vs. grazing for management of species-rich grasslands: a multi-site, multi-year field experiment. – *Nordic J. Bot.* **33**: 761–768.

Tälle, M., Deák, B., Poschlod, P., Valkó, O., Westerberg, L. and Milberg, P. 2016. Grazing vs. mowing: a meta-analysis of biodiversity benefits for semi-natural grassland management. – *Agric. Ecosystems Environ.* **222**: 200–212.

Török, P., Vida, E., Deák, M., Lengyel, S. and Tóthmérész, B. 2011. Grassland restoration on former croplands in Europe: an assessment of applicability of techniques and costs. – *Biodivers. Conserv.* **20**: 2311–2332.

Valkó, O., Török, P., Deák, B. and Tóthmérész, B. 2014. Prospects and limitations of prescribed burning as a management tool in European grasslands. – *Basic Appl. Ecol.* **15**: 26-33.

Valkó, O., Deák, B., Magura, T., Török, P., Kelemen, A., Tóth, K., Horváth, R., Nagy, D.D., Debnár, Z., Zsigrai, G. and Kapocsi, I. 2016. Supporting biodiversity by prescribed burning in grasslands—A multi-taxa approach. – *Sci. Total Environ.* **572**: 1377-1384.

Van Langevelde, F., Van De Vijver, C.A.D.M., Kumar, L. Van De Koppel, J. De Ridder, N. Van Andel, J., Skidmore, A.K. Hearne, J.W., Stroosnijder, L. Bond, W.J., Prins, H.H.T. and Rietkerk, M. 2003. Effects of fire and herbivory on the stability of savanna ecosystems. – *Ecology* **84**: 337–350.

Végvári, Z., Valkó, O., Deák, B., Török, P., Konyhás, S. and Tóthmérész, B. 2016. Effects of land use and wildfires on the habitat selection of Great Bustard (*Otis tarda* L.): implications for species conservation. – *Land Degrad. Devel.* **27**: 910-918.

Wahlman, H. and Milberg, P. 2002. Management of semi-natural grassland vegetation; evaluation of long-term experiment in southern Sweden. – *Ann. Bot. Fennici* **39**: 159-166.

Werner, G.D., Cornwell, W.K., Sprent, J.I., Kattge, J. and Kiers, E.T. 2014. A single evolutionary innovation drives the deep evolution of symbiotic N₂-fixation in angiosperms. – *Nature Comm.* **5**: p.4087.

Figure 1. Study locations in southern Sweden. Ekenäs and Tagel had two experimental sites each.

Figure 2. The difference in frequency in annually burnt and annually mowed plots, where positive numbers indicate that a species is promoted by spring burning. a) species-wise differences. The number to the right is frequency of a species (max 22); b) Weighted average of species grouped according to lifeform. Numbers to the right is number of species in the group.

Figure 3. Relationship (meta-regression) between response to spring burns (where a positive response indicate that a species is promoted by spring burning) and a) species' nectar production (slope 0.236, CI: 0.134; 0.338, $P < 0.001$), b) plant height (0.020; CI 0.014; 0.025; $P < 0.0001$), and c) Ellenberg N-value (0.164; CI: 0.236; 4.514; $P < 0.001$).

Table 1. Description of the experimental sites established in southern Sweden for the comparison of the management methods in semi-natural grassland vegetation.

Site	Start	Inventory	Vegetation type	Management at start of trial	Soil type
Österplana	1973	1986	Dry	Grazing	Gravelly clay loam
Ekenäs mesic	1975	1986	Dry-mesic	Grazing	Humus-rich loamy
Bräcke	1973	1986	Mesic	Grazing	Silt
Sättra	1973	1986	Mesic	Grazing	Slightly clayey sand
Bråbo	1973	1986	Mesic	Grazing	Rock moraine
Tagel mesic	1973	1986	Mesic	Grazing	Rocky sand
Tagel former field	1973	1987	Mesic	Grazing, fertilized	-
Dämkärr	1973	1986	Mesic	No management for 3 yrs	Humus-rich silt
Gränö	1973	1986	Moist	Mowing, fertilized	Slightly humus-rich silt
Ekenäs moist	1975	1986	Moist	Grazing	Highly humus-rich light clay
Andersby	1973	1986	Moist	Grazing	Humus-rich light clay

Fig. 1. Study locations in southern Sweden. Ekenäs and Tagel had two experimental sites each.