Neonatal imitation: Temporal characteristics in imitative response patterns

Mikael Heimann and Tomas Tjus

The self-archived postprint version of this journal article is available at Linköping University Institutional Repository (DiVA):

<http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-159706>

N.B.: When citing this work, cite the original publication.

Heimann, M., Tjus, T., (2019), Neonatal imitation: Temporal characteristics in imitative response patterns, *Infancy*, 24(5), 674-692. <https://doi.org/10.1111/inf.12304>

Original publication available at:

<https://doi.org/10.1111/inf.12304>

Copyright: Wiley (24 months)

<http://eu.wiley.com/WileyCDA/>

Neonatal imitation – Temporal characteristics in imitative response patterns

Mikael Heimann¹ and Tomas Tjus^{2,3}

¹The Infant and Child Lab, Division of Psychology, Department of Behavioural Sciences and Learning, Linköping University, Sweden

²Department of Psychology, University of Gothenburg, Sweden

³ deceased on Dec 7, 2018

Accepted *Infancy* May 18, 2019

Corresponding author: Mikael Heimann, Department of Behavioural Sciences and Learning, Linköping University, SE 581 83 Linköping, Sweden. Email: mikael.heimann@liu.se

Abstract

Neonatal imitation has been an area that has attracted intense attention within developmental psychology. Reported here are data from 33 newborn infants (16 girls; mean age: 47 hrs.) assessed for imitation of tongue protrusion and mouth opening. The stimuli were presented dynamically, in three 20 second long gesture modeling intervals, interwoven with three 20 s long intervals in which the presenter kept a passive face following the procedure used by Meltzoff and Moore (1983a). Imitation of tongue protrusion emerged among a majority of the infants during the first 60 seconds of the experiment. In contrast, mouth opening showed a protracted response and a majority exhibited imitation after 60 seconds. The individual response pattern of the participating infants varied substantially over the course of the experiment. The study provides renewed support for neonatal imitation of mouth opening and tongue protrusion, and in addition, suggests that the temporal organization of the responses observed is an important factor to consider, which in turn has methodological and theoretical implications.

Key words

Neonatal Imitation, Tongue protrusion, Mouth opening, Motor organization

How we understand a newborn infant's behavioral capacities influences our theories of how the mind is formed and develops. Thus, there is no wonder that the issue of neonatal imitation has drawn intense interest within developmental psychology. Meltzoff and Moore (1977) published two studies in *Science* that discussed methods for assessing early imitation, effective eliciting conditions, and the three mechanisms that might underpin this early social competence. They reported imitation of three facial gestures (tongue protrusion, mouth opening, and lip protrusion) as well as manual imitation. This work was related to prior, less formal reports of early imitation (Gardner & Gardner, 1970; Maratos, 1973; Zazzo, 1957), but went beyond them in terms of methodological rigor and in describing the implications for the infants' representation of action and social cognition.

Meltzoff and Moore's findings motivated further empirical work testing neonates' imitative capacity, with results both confirming and extending their findings (e.g., Field, Woodson, Greenberg, & Cohen, 1982; Heimann & Schaller, 1985; Kugiumutzakis, 1998; Meltzoff & Moore, 1983a, 1989; Nagy, Pal, & Orvis, 2014; Nagy, Pilling, Orvis, & Molnar 2013; Vinter, 1986) and in several unsuccessful studies (e.g., Hayes & Watson, 1981; Koepke, Hamm, Legerstee, & Russell, 1983), with the latter rebutted as having methodological problems (e.g. Meltzoff & Moore, 1983b).

Lately, interest in the existence and mechanisms of early imitation has intensified, as exemplified by the suggestion that early imitative responses are acquired through associative learning (Heyes, 2016) and a study which attempted a new design and reported null effects (Oostenbroek et al., 2016). However, Meltzoff, Murray et al. (2018) reported a detailed reanalysis of data of Oostenbroek et al.'s raw data, which revealed significant support for imitation of tongue protrusion at 1-, 3-, 6-, and 9-weeks, with an accompanying explanation for how the author's may have missed the pattern. Meltzoff, Murray et al. (2018) also presented five recommendations for future studies to consider in order to facilitate

comparisons between findings: Limit the number of models used in within-subjects studies, allow sufficient time for neonates to visually process the display and organize their motor responses, have a good control of the physical environment in which the experiment is carried out, a better control of the social environment (limit the infant's prior social interactions with the experimenter), and finally, use of pilot testing to ensure that the methods used are suitable for the age tested (in their new design Oostenbroek et al. used too many models in a within-subject procedure, 11 different displays, and some of the displays were motorically impossible for neonates to generate). These recommendations are relevant for the current study. Although the current study was carried out before these recommendations were codified by Meltzoff et al. (2018), many of these issues had been discussed in previously published papers on early imitation, and thus we were able to capitalize on recommended "effective eliciting conditions" in designing the current study.

Theoretically, several mechanisms or explanations have been proposed of an early imitative capacity. Two dominant accounts are the Active Intermodal Matching (AIM) mechanism (Meltzoff, 1990; Meltzoff & Moore, 1997) and various attempts to describe the neural underpinnings of early perception-action coupling (Ferrari, P. F. et al., 2012; Marshall & Meltzoff, 2014). In addition, it has recently been suggested that the characteristics of newborn's imitative responses might fit an imprinting process (Bard & Nagy, 2017; Nagy & Molnar, 2004). According to the AIM account, newborn imitation does not entail a passive association. Instead it is described as a "matching to target" process (Meltzoff & Moore, 1997, p. 180) and an "active attempt to adapt and gradually refine their own movements with respect to others" (Dumas, Kelso & Nadel, 2014; p 1). The imprinting hypotheses, in contrast, builds on Nagy's (2011) arguments for viewing the neonatal period as a sensitive period for early imitation. According to this view, neonatal imitation serves a communicative purpose and is a response most easily elicited during the neonatal phase.

Bjorklund (1987) pointed out that an observed behavioral response in neonates can reflect several underlying processes, both hard-wired and voluntarily controlled. It might very well be the fact that early facial gestures in a neonate could reflect several processes such as a response pattern important for feeding, an increase due to arousal, as well as increases due to an active effort of the infant to match an adult's behavior. The last part, the voluntary response, is probably the most demanding process for the neonate but we know that it is possible for newborns to guide their motor responses already during the first week of life (Rönnqvist & von Hofsten, 1994).

It has also been reported that it takes time for the neonate to organize a coherent response, and thus if the displays and response measurement intervals are too brief, this may suppress the imitative responses (Holmlund, 1995; Meltzoff & Moore, 1997, 1983a). Heimann, Nelson and Schaller (1989) found support for neonatal imitation during the first 60 seconds of the 150 second response period (in this study, the response period started after a 60 second baseline and a 40 second stimulus demonstration period). Moreover, Holmlund (1995), describes how 1-week-old infants orient and respond to an adult in a face-to-face situation. The infants “stay oriented for at least 20 seconds without activity in any part of the body. It seems that their attention has to remain focused for quite a long time for storing and identification processing” (Holmlund, 1995, p 56). In line with this observation, Heimann (2002) stated that “a newborn child achieving reasonable matching does so in spite of the fact that many systems are immature” (Heimann, 2002, p 81). Thus, it is possible that how different researchers design their experiments, e.g. how the stimuli are presented, how long one waits for a response to occur, or how the data are collapsed might affect the results of any specific study (and may explain some of the divergent findings in the literature). Some of these issues are addressed in the current study which focuses on two gestures, tongue protrusion and mouth opening, and how the newborn's responses emerge over time both when

the dynamic stimuli are demonstrated and when the adult's face is passive during the "burst-pause" test paradigm. The methodology builds on our earlier study (Heimann et al., 1989) and the procedure proposed by Meltzoff and Moore (1983a) which was designed for newborn infants.

Neonatal imitation, as used in this paper, is defined on purely behavioral grounds. This implies that a neonate is judged to imitate a specific behavior, e.g. mouth opening, if the observed frequency of that behavior increases while watching an adult model that behavior in comparison to when an alternate behavior is modeled (e.g. tongue protrusion). This "cross-model comparison" was introduced by Meltzoff and Moore (1977; 1983b; 1994) and has since then been used to define imitation in newborn infants in both recent (e.g., Nagy et al., 2013; Simpson et al., 2014) and earlier publications (e.g. Heimann, 1989; Meltzoff & Moore, 1983a; 1989; also see Meltzoff, Murray et al., 2018, for a recent update on how to measure neonatal imitation). The cross-model comparison paradigm also implies, as pointed out by Meltzoff and Moore (1983b), that no assumption of the underlying mechanism is needed when judging if an infant imitates or not. Furthermore, "differential imitation neutralizes the arousal explanation" (Vincini, Jhang, Buder & Gallagher, 2017, p 14).

Opponents have suggested that what looks like an imitative response might be better explained through arousal processes or a coincidental match (Jones, 1996, 2017), an innate releasing mechanism (Anisfeld, 1996), as based on learning since a neonate does not have the cognitive capacity to "solve the correspondence problem that link self with other for imitation" (Heyes, 2016, p 6), or as a response dependent on the aerodigestive system (Kevin & Akins, 2017; but see Meltzoff, 2017 and Simpson et al., 2017 for critiques of this view). More specifically, Heyes proposes that an associative learning model explains how and why our capacity to imitate develops, whereas Vincini et al. (2017) specifies an association by similarity process as most likely to explain early imitation. These opposing views, it should be

noted, are not all compatible with one another; e.g., the innate releasing view contradicts the idea of gradual associative learning and the claims that the effect is wholly reduceable to arousal or artifacts.

In contrast, those interpreting the evidence in favor of an early imitative ability usually propose that neonatal imitation is an early socio-communicative skill (e.g., Meltzoff, 2007; Trevarthen, 2011). Neonatal imitation has been described as a starting state for the process leading up to theory of mind – Meltzoff’s (2007) “Like-Me hypothesis,” as an active communicative act (Nagy & Molnar, 2004), or as a first step in the development of intersubjectivity (Trevarthen, 2011; Trevarthen & Aitken, 2003). Furthermore, neonatal imitation has also been documented among non-human primates (Bard, 2007; Ferrari, P. F. et al., 2006; Paukner, Simpson, Ferrari, Mrozek, & Suomi, 2014; Simpson et al., 2014) and studies on infant rhesus macaques suggest that neonatal imitation is important for later social development (Dettmer et. al., 2016; Simpson, Miller, Ferrari, Suomi, & Paukner, 2016).

The main aim of this paper is not to discuss whether or not neonatal imitation is present at the population level, although this issue will also be addressed to some extent. Instead, the paper primarily aims at presenting new observations from a Swedish sample of newborns using a study designed to take into account the motor limitations of neonates and also one that was sensitive to possible individual differences among neonates. Several years ago we reported replications of the findings of neonatal imitation and also introduced observations supporting the notion that individual differences in imitation are observable already during the neonatal period (Heimann, 1989; Heimann & Schaller, 1985; Heimann et al., 1989). These findings also suggested that imitation of tongue protrusion was easier to elicit than imitation of mouth opening and moreover that imitation is particularly salient after the infant observes the modeling and when the presenter is passive (Heimann, 1991, 2001, 2002; Heimann et al., 1989).

Thus, two hypotheses were formulated. First, we hypothesized that we would observe neonatal imitation of both tongue protrusion and mouth opening as evident by an increase in observed frequency when presenting the gesture (e.g., mouth openings when mouth openings are presented) as compared with observed frequencies when presenting an alternate gesture (e.g., mouth openings when tongue protrusions are presented). Formulating our expectations more precisely, we predicted that the clearest evidence of neonatal imitation would occur not as an instant reaction but when the infant had had some time to organize a coherent response. Thus, we decided to split the analysis into different time windows, one early focusing on responses during the first minute and one late window covering responses emitted later. Second, building on Heimann et al.'s earlier studies, we hypothesized that imitative responses would occur primarily during the intervals in which the presenter kept a passive face rather than during the intervals when the stimuli were demonstrated dynamically. Third, for exploratory reasons but also building on earlier reports (e.g. Heimann, 1992; Heimann et al., 1989; Meltzoff & Moore, 1983a), we aimed at presenting observations on the infants' individual imitation pattern.

Methods

Participants

Thirty-three newborn infants (16 females) were observed at two days of age ($M = 47$ hours; $SD = 22.9$), all recruited from the maternity ward at Sahlgrenska University Hospital, Gothenburg, Sweden. All newborns were healthy (Apgar score at 5 minutes: $M = 8.65$; $SD = 1.5$), full-term (mean gestational age = 40.3 weeks; $SD = 1.3$), and in a good clinical condition.

Procedure

The test took place in a separate room at the maternity ward. All sessions began when the infant was judged to be awake and alert and all observations included demonstration of

two gestures: tongue protrusion (TP) and mouth opening (MO). The order was counterbalanced: for half of the participants, MO was presented first and for half TP came first. The procedure for each gesture was 120 seconds long (see Table 1). The observation began by experimenter A (female) presenting the first gesture to the infant. Experimenter B controlled the video camera and the exact timing of the procedure; experimenter A did not know until the start of the procedure if she should present mouth opening or tongue protrusion (this verbal signal from experimenter B was absent from the edited videotapes for coding).

Insert Table 1 About Here

The aim was always to start the demonstration of a gesture at a time when the infant did not display any spontaneous mouth movements (no yawning, no obvious mouth opening or tongue protrusion). For each gesture the experimental procedure was broken up into six intervals of 20 s each ($M = 20.1$ s; $SD = 3.1$ when presenting TP and $M = 20.3$; $SD = 3.1$ for the MO part of the experiment) following the Meltzoff and Moore (1983a) “burst-pause” methodology, which alternated a demonstration period with a passive-face period: Three dynamic demonstrations of the stimuli (Demo interval 1, 3 & 5) and three passive face periods (interval 2, 4 & 6) allowed the infant to build up a response. The mean number of times tongue protrusion were presented during each demonstration period varied from 9.68 to 10.84 ($SD = 3.39 - 3.97$). For mouth opening the presentation frequency varied between 9.21 and 9.48 ($SD = 2.01 - 2.59$).

During the passive face intervals the experimenter kept eye contact with the infant but did not present any facial movement. There was a brief pause of about 20 to 30 seconds after 120 seconds, when the first gesture had been presented, in order to make sure that the infant was alert and ready for the next gesture.

During the test the majority of the infants were securely supported on experimenter A’s lap, while a few were supported in an infant chair in order to promote face-to-face interaction

between the experimenter and the infant. The distance between the experimenter's face and the infant was approximately 25 cm during the session. For all subjects except one, the mother was present during the experiment. She was seated behind the infant (approximately at an angle of 45 degrees) so that she could view the procedure but precluding any gaze contact between her and her infant while the gestures were presented. The mother was passive and silent throughout the experiment.

Attrition. Of 73 families approached, 47 indicated that they were willing to participate. However, 11 mothers left the hospital earlier than planned, one mother changed her mind, and two infants were excluded due to drowsiness. Thus, acceptable observations exist for 33 newborns although some of them became tired or fussy during the experiment. A few of these 33 infants were excluded from part of the analysis due to fussiness or not spending enough time in an alert state during the whole observation period. Thus, the actual n varies from 33 during intervals 1 and 2 (the first 40 s of the experiment) to 26 for mouth opening and 24 for tongue protrusion during intervals 5 and 6 (the last 40 s of the experiment).

Coding criteria and definition of imitation

Tongue protrusion (TP). The minimum criteria was that the tongue had to pass the posterior part of the lip. The minimum criterion was that a clear forward movement of the tongue was noted although the tongue was not protruded beyond the outer part of the lips. Responses associated with hiccups or strong bowel movements were not included in the analysis. This was true for both gestures.

Mouth opening (MO). Mouth opening was defined as a clearly visible separation of the lips, although the end state needed only to be a minor change (not exceeding the width of the lips). In order to be coded as a mouth opening, no simultaneous protrusion of the tongue was accepted. Yawning was not included as an adequate mouth opening response.

Definition of imitation. Imitation was operationalized as a difference between gestures emitted when the target gesture was modeled as compared with responses observed when the non-target gesture was demonstrated. More specifically this means that imitation of tongue protrusion was measured by comparing the frequency of relevant tongue movements when TP was modeled (imitation condition) as compared with the frequency of when MO was modeled (control condition; see Meltzoff & Moore, 1983a, 1989 for a similar analysis). Similarly, imitation of mouth opening was measured by comparing the frequency of relevant mouth movements as defined above when MO was modeled (imitation condition) with the frequency when TP was modeled (control condition).

State: In addition to the above facial categories independent coders also judged the infant's state from the obtained video recordings based on the following criteria (adopted from Theorell, Prechtl, Blair & Lind, 1973 and Wolff, 1987): State 1: Quiet and/or active sleep or drowsiness; State 2: Quiet awake, eyes open, no movements; State 3: Active awake, eyes open, gross movements; State 4: Crying. State was monitored continuously and only intervals when the infants had been judged to be in state 2 or 3 were included in the final analysis.

Reliability. All tapes were coded blind by four graduate students who were trained until they performed at a minimum level of 85 percent agreement. The coding entailed two behaviors (mouth opening and tongue protrusion) and state. The final overall agreement was calculated using Cohen's Kappa (Cohen, 1992). The obtained reliability coefficients were $\kappa > .75$ for all measures.

Ethics

The present study was conducted according to guidelines laid down in the Declaration of Helsinki, with written informed consent obtained from a parent or guardian for each child

before any assessment or data collection. All procedures involving human subjects in this study were approved by the ethical committee at the University of Gothenburg, Sweden.

Statistical analysis

The repeated measures ANOVAs conducted for the main analyses was based on the observed frequencies of gestures produced divided by seconds for each time window thus controlling for the unequal length of the time windows used. For the pre-analysis of the infants' looking pattern Student t-test was used. An $\alpha \leq .05$ was used as a cut-off for statistical differences and effect sizes are reported as η_p^2 for ANOVAs and as Cohen's d (Cohen, 1992) for Student t-test. SPSS (Statistical Package for Social Sciences 23.0) was used for all statistical analyses.

Results

The result section is organized into four parts. The first part deals with pre-analyses presented for clarity reasons, while the three following parts presents results pertinent to our hypotheses and aim: Findings regarding neonatal imitation within different time windows are presented first followed by findings relevant to our second hypothesis stating that imitation would be more likely during the intervals in which the presenter keeps a passive face. Finally, imitative response patterns are presented.

Pre-analysis

Attention. The participating infants looked significantly more towards the model's face than looking away when any of the two gestures were modelled (see Table 2). The mean percentage of looking at the model was 61.1% when mouth opening was dynamically presented and 57.7% when the model presented tongue protrusion. An additional correlational analysis between looking time and observed imitative responses revealed no significant associations.

Number of time windows. Although hypothesis-1 specified two time windows (within the first minute or during the second minute) this was changed before carrying out the final analysis based on the observation that few infants (< 40 %) imitated during the final 20 seconds of the experiment. Thus the experiment was divided into three time windows: The first 60 seconds (time window 1), 60 to 100 seconds into the experiment (time window 2), and finally the last 20 seconds of the experiment (time window 3).

Insert Table 2 About Here

Responses within the three time windows (Hypothesis 1)

Time window 1, the first 60 seconds. A 2 (condition: experimental/control) x 3 (intervals 1, 2 and 3) repeated measures ANOVA showed a significant main effect for imitation of tongue protrusion, $F(1,29) = 4.857, p = .036, \eta_p^2 = .143$; see Table 3. The main effect for time was also significant ($F[2,28] = 3.605, p = .04; \eta_p^2 = .205$) while the condition x time interaction was not ($p = .277$). A comparable analysis of mouth opening (see Table 4) during the first minute revealed no significant effects for imitation or for time or the interaction.

Insert Tables 3 and 4 About Here

Time window 2, 61 – 100 seconds. A 2 (condition: experimental/control) x 2 (intervals 4 and 5) repeated measures ANOVA revealed no main effect of imitation of tongue protrusion. Time was also non-significant while the interaction condition x time was significant ($F[1,25] = 4.859, p = .037, \eta_p^2 = .163$). A closer inspection of the results suggest that this significant interaction reflects a tendency towards imitation of tongue protrusion noted during the fourth interval ($p = .088$).

A similar analysis for mouth opening revealed a different pattern. The main effect for imitation of mouth opening was significant, $F(1,26) = 5.269, p = .030; \eta_p^2 = .169$ (see Table 4) but neither the effect of time nor the interaction time x condition.

Time window 3 (TW3), 101 – 120 seconds. As evident from Tables 3 and 4 there were no indication of imitation of any of the two gestures during the final 20 seconds of the experiment. A visual illustration of how the infants changed their tendency to respond to the target gestures across the three different time windows are depicted in Figures 1 and 2.

Insert Figures 1 and 2 About Here

Comparing demonstration and passive face periods (Hypothesis 2)

Comparing the mean frequencies across the three demonstration intervals with the three passive face intervals (see Table 1 for the design) revealed a non-significant difference as a function of whether the display was currently in view or had just been demonstrated, for both mouth opening and tongue protrusion (see Table 5). The infants were as likely (or not likely) to open their mouth or protrude their tongue when the experimenter kept a passive face as during the dynamic demonstration of the gesture.

Insert Table 5 About Here

Individual response patterns (exploratory aim)

Imitation was also analyzed based on each infants' individual tendency to imitate or not imitate during the three time windows used in the analysis, As suggested by Meltzoff and Moore (1983a), a participating infant was judged to imitate a gesture if the frequency of that gesture in the imitation condition exceeded the frequency observed in the control condition. As evident from Table 6 more than half (53.3 %) of the infants imitated tongue protrusion during the first time window but less so during time windows two (42.3 %) or three (29.2 %). For mouth opening a strong majority imitated during time window two (66.7 %) while only about one-third of the infants displayed an imitative response during time windows one or three.

Insert Table 6 About Here

Discussion

Two hypotheses and one exploratory aim were formulated at the outset of the study: (a) That the infants were expected to display their strongest imitative response when they had had time to organize a coherent response, (b) that the strongest evidence for imitation would be found during the passive face intervals, and finally (c) we also expected to describe individual variation patterns in imitation observed among the infants. Globally, we found support for neonatal imitation of tongue protrusion during the first minute and for mouth opening one minute into the experiment, thus confirming our first hypothesis. In contrast, no support was found for hypothesis number two, the infants imitative responses were not more common during the passive face intervals overall, compared to the demonstration periods. Finally, the individual imitation patterns were found to vary strongly between the three defined time windows used in the analysis.

Support for an imitative response pattern was found for tongue protrusion during the first time window (the first 60 seconds of the experiment), a finding in line with a large number of previous papers that have reported evidence for an ability to imitate tongue protrusion (see summaries in Meltzoff & Moore, 1997; Meltzoff, Murray et al., 2018; Nagy et al., 2014; Simpson et al., 2014). It has also been proposed that evidence of an early imitative capacity exists for tongue protrusion only (Anisfeld, 1996), but this is countered by our findings of neonatal imitation of mouth opening as well.

The finding that imitation of tongue protrusion was evidenced during the first time window was not according to our hypothesis. We had expected that the infants' imitative responses would build up over a longer time frame, that the strongest indication of imitation would be observed about one minute into the experiment, an expectation not confirmed for tongue protrusion (but was confirmed for mouth opening). Regardless, our finding add

support to the many previous studies having found evidence of an imitative capacity in newborns (e.g., Heimann et al., 1989; Meltzoff & Moore, 1977; 1983a,b; Nagy et al., 2013).

A different temporal response pattern was observed when the infants were faced with a model presenting mouth opening. Here, no indication of imitation was detected during the first minute of the experiment. This seemingly supports the argument in the literature that imitation of mouth opening is absent or more difficult to elicit (Anisfeld, 1996; Heyes, 2016; Oostenbroek et al., 2016). However, the picture changed after the first minute, and an imitative response pattern was observed during the middle time window, 61 - 100 seconds into the procedure, which is in accordance with our first hypothesis: namely, that it might take some time for the infants to display imitation, as also suggested by a few previous papers (Heimann et al., 1989; Holmlund, 1995; Meltzoff & Moore, 1994). It has also been reported that infants often adjust their responses over time and that their responses become more precise over trials (Kugiumutzakis, 1998; Meltzoff & Moore, 1977, 1983a, 1994, 1997). In the current study, it took a full minute until the infants displayed a differential response indicating imitation of mouth opening. This supports our expectation that infants this young are able to imitate mouth opening and, in addition, the hypothesis that they need time to organize a coherent imitative response. This result, if upheld, also suggests that mouth opening, as a specific response, takes longer time to organize for the newborn than tongue protrusion. Although speculative, mouth opening might be controlled differently than tongue protrusion (e.g., proprioceptive information from lip and jaw movements may be different from simple tongue movements) and also be more prone to arousal fluctuations¹. After the newborn period, mouth opening might also be a more socially reinforced behavior (Murray, DePascalis, Bozicevic, Hawkins, Scafani & Ferrari, 2016; Rayson, Bonaiuto, Ferrari &

¹ Mouth opening is, at least in our study, a more common behavior as evident by the overall higher mean frequencies compared to tongue protrusion (but clearly this depends on the response definitions adopted).

Murray, 2017) and be influenced by the speech and smile movements infants see in social interactions (Kuhl & Meltzoff, 1996; Meltzoff & Moore, 1994). Rayson et al. (2017) found that the mothers' mirroring of smiles and mouth opening when the infant was 2 months predicted "infant motor system activity during observation of the same expressions at nine months" (Rayson et al., p 1). Murray and colleagues also followed mother-infant dyads over the first two months finding that the degree of the mothers' mirroring increased over time and especially so for social behaviors such as mouth movements. These observations suggest that early infant behaviors and mirroring (and imitation on the infant's part) can be influenced by infant-parent face-to-face interactions as suggested for early imitative behaviors by Heimann (1989) and Meltzoff, Murray et al. (2018). These observations do not, however, explain why imitation of mouth opening took a minute to manifest itself in the current newborn study, except inasmuch as the proprioceptive information may take some time to be used by young infants.

Our second hypothesis specified that imitation would be most strongly evidenced not when the experimenter demonstrated the gesture but instead during the passive face periods. This prediction was not confirmed by the results. Comparing the total number of responses emitted when a gesture was dynamically presented with the responses during the passive face intervals did not reveal any significant differences. This was true for both mouth opening and tongue protrusion. This suggests that our initial idea that infants imitate differently in these two situations – seeing the action or seeing a passive face – might be false. In retrospect, previous microanalyses of newborns' response profiles also reported imitation during both the stimulus-demonstration and the passive-face intervals (e.g., Meltzoff & Moore, 1989). Thus, using the burst-pause procedure with 20-s intervals, the obtained effect seems to be a robust phenomenon. However, it might also be that longer passive face period as for instance used in

some other studies (e.g., Heimann et al., 1989; Nagy et al., 2013) would have been better suited to achieve support for our second hypothesis.

In order to understand more about the variability in imitation we also explored the individual response patterns of the infants. This analysis revealed that the number of infants actually showing an imitative response varied between the different time windows used. The strongest indication of imitation was observed for mouth opening during time window 2 when nearly 70% of the infants imitated. A similarly strong figure was not noted for tongue protrusion. Even during the first time window, when the overall result was significant, only slightly more than 50% of the infants actually imitated tongue protrusion. This kind of variation in newborn infant's proneness to imitate should not come as a surprise; several previous studies have reported on individual variations in neonatal imitation (Ferrari, P.F. et al., 2009; Heimann & Schaller, 1985; Heimann et al., 1989; Meltzoff & Moore, 1983a, 1989; Nagy, Kompagne, Orvos, & Pal, 2007), and the causes and sequelae of individual differences are a topic of importance to pursue in future studies.

Imitation is a response that emerges from the interplay between what the newborn perceives (e.g. the adult demonstration), the infant state (motivational state, alertness and attention) and what the infant selects to do (motor output). Thus, to imitate is a complex process that relies on several systems, many of which are immature at birth, e.g. systems that controls vision, motivation, attention and motor responses. Sometimes the outcome is a clear match between perception and action, and an imitative response is observed quickly. At other times it takes some time for the infant to imitate, and sometimes there is no match at all and imitation is not observed. As a note for further studies we found that the infants did not focus 100 percent on the experimenter's face when the gestures were demonstrated. In fact, they spent approximately 60 percent of their time looking at the experimenter and 40 percent of the time gazing away. It is not possible to evaluate what this pattern means but we suspect that

gaze away might in part reflect periods when the infant is processing the information available, allowing time to prepare for a possible motor response. Such a pattern has been evidenced in older children (e.g. Previc, Declerck & de Brabander, 2005).

Meltzoff and Moore (1997) and Heimann (2001, 2002) offered proposals about the role of experience in early imitation. According to Meltzoff and Moore (1997, p 184) “infants move their limbs and facial organs in repetitive body play analogous to vocal babbling”. The authors labeled this as “body babbling,” and their AIM theory proposes that infant motor experience, specifically including prenatal movements, may play a major role in early imitation (Meltzoff & Moore, 1997). In fact, fetuses have been found to respond to some maternal vocalizations with related movements already at gestation week 25 (Ferrari, G. F et al., 2016). We suspect that newborns differ in how these patterned movement representations actually are constructed over time. Moreover, if “comparative processes between own movements and other movements are involved in neonatal imitation, and if prenatal differences in patterned movements exist, then this comparison must lead to different outcomes between children” (Heimann, 2002, p 81). This development might then, in part, explain why individual newborn infants differ in their proneness to imitate at birth. If neonatal imitation can be elicited for both tongue protrusion and mouth opening, as the current findings suggest, it raises the issue about the neuropsychological mechanisms behind this ability.

Nagy (2011) has argued for a unique place in development for the neonatal period, being a stage defined by the transition from intrauterine to extrauterine environment representing a sensitive period and an “experience-expectant stage of development”. More specifically, she highlights the first week of life, the perinatal period, as a moment in time when the infant might be especially prone to produce imitative responses (Nagy et al., 2013; 2014). In line with this, Nagy and Molnar (2004) have suggested that, since temporal cortex is

immature at birth, it is likely that also subcortical structures are involved in imitation during the neonatal period. One such candidate is the superior colliculus (Bjorklund, 1987; Heimann et al., 1989; Vinter, 1986), a sub-cortical structure relatively mature at birth suggested to process multisensory information (Stein & Meredith, 1993; Stein & Stanford, 2013). A recent computational modeling experiment successfully demonstrated how visual and somatotopic input could be combined through a multimodal visuo-tactile layer (Pitti, Kuniyoshi, Quoy, & Gaussier, 2013), but this computer modeling finding is suggestive at the most. A more common and probably also widespread theoretical account proposes that infant body representations may also play a role, as described by Meltzoff and Moore's (1997) AIM theory. Interestingly, there are emerging electrophysiological studies (EEG and MEG) documenting cortical body maps in infants as young as 60 days old (Meltzoff, Saby, & Marshall, 2018; Saby, Meltzoff, & Marshall, 2015), with arguments offered about how these neural body maps may link self and other in imitation (Meltzoff, Ramírez, Saby, Larson, Taulu, & Marshall, 2018). Taken together, it seems possible that several different processes, e.g. prenatal experience, biological preparedness (e.g. a spatial representation of the body), and perceptual-motor skills all might contribute to the imitative capacity we observe in a newborn infant.

In conclusion, we interpret our findings in support of the active intermodal matching (AIM) model proposed by Meltzoff (1990; Meltzoff & Moore, 1997) although other theories exist that might accommodate our results. An alternative theory of special interest is the idea of rapid learning based on a filial imprinting process as proposed by Nagy and colleagues (Nagy et al., 2014). Although our finding that for mouth opening the infants needed a minute to build up an imitative response could be viewed as an instance of rapid learning the AIM model is, in our view, still the most comprehensive theoretical framework for explaining neonatal imitation. Furthermore, we hypothesize that this system is mainly governed

subcortically early in development. It also seems possible that early imitation processes rely more on distributed networks while later imitation will rely on more specialized circuitry as recently found for the development of working memory (Bathelt, Gathercole, Johnson, & Astle, 2017). Associative learning seems less plausible as an explanation of imitation in newborns, although it probably comes into play as a child's imitative ability or motivation to imitate develops over time. We view it as highly unlikely that a mechanism such as associative learning can explain the imitative response observed for newborn tongue protrusion. The age of infants – only around 2 days – leaves little room for establishing a capacity for facial imitation through parent-infant interactions and parental training. This is especially true for the act of tongue protrusion, a behavior not commonly observed within spontaneous social interactions between parents and their baby; tongue movements are rather seen as important for sucking in feeding situations during the neonatal period (Kevin & Akins, 2017).

Limitations

The sample size used in this study was not large and the low N was further hampered by loss of infants due to changes in their state. That is, they moved from an alert state to drowsiness or sleep. Thus we cannot exclude that our study, at least in part, suffer from low statistical power. Another limitation is the decision not to include a separate passive face period prior to the demonstrations in which to collect baseline data. This decision was based on our aim to focus on the differential response between the two gestures and the fact that our participants were newborns with a limited window of awake time. Although useful, such baseline periods are probably more suitable for older infants. The within-group design used might also have introduced unwanted noise in the data since as pointed out by Ullman (2016), there is a risk that representations that resemble each other create interference. A use of a wider range of gestures would also be useful. Nonetheless, by counterbalancing the

demonstrations of the two facial gesture in the experiment and having second experimenter strictly controlling when to start and when stop presenting the gestures we were able to conduct a study that focused on our key questions and was short enough to be tolerated by these young infants.

Conclusion

Building on the methodology used in a seminal paper on newborn imitation (Meltzoff & Moore, 1983a) this study provides further support for the idea that human newborns have the capacity to match mouth opening and tongue protrusion gestures in an imitative like fashion. However, our findings also provide useful details, texture, and refinements to the overall picture. First, whether imitation is observed depends partly on the time window used. This was true for both tongue protrusion and mouth opening; imitation being observed within different time windows for the two gestures (which may explain why some investigators, using brief test periods, have found it difficult to elicit mouth opening using other designs). Second, we did not find support for our initial idea that neonates are more prone to imitate during the periods when the presenter kept a passive face, and finally, we note as several others have done before us (e.g. Heimann et al., 1989; Heimann, 2002; Meltzoff, 1983a; Simpson et al., 2014) a large variability in imitation among the infants. The causes and consequences of these individual differences are of great interest.

Acknowledgement

The authors wish to thank all of the parents and children who participated. We are also grateful to Agneta Swerlander and Eva Ullstadius for collecting part of the data and to Andrew N Meltzoff and Vasudevi Reddy for providing insightful comments on an earlier draft. This study was supported by grants from the Bank of Sweden Tercentenary Foundation (89/313), the Swedish Council for Working Life and Social Research (2008-0875) and the Swedish Research Council (2008-2454) to Mikael Heimann. The authors declare no conflicts of interest with regard to the funding source for this study.

Finally, this paper honors the memory of Tomas Tjus, second author of the paper, who passed away a few days before the final revision was submitted. His scientific insight, especially his analytic skills, was fundamental to the completion of the study.

References

- Anisfeld, M. (1996). Only tongue protrusion modeling is matched by neonates. *Developmental Review, 16*(2), 149-161.
- Bard, K. (2007). Neonatal imitation in chimpanzees (*Pan troglodytes*) tested with two paradigms. *Animal Cognition, 10*, 233–242. <https://doi.org/10.1007/s10071-006-0062-3>
- Bard, K., & Nagy, E. (2017). Status of the human newborn. In B. Hopkins, E. Geangu, & S. Linkenauger (Eds.), *The Cambridge Encyclopedia of Child Development* (pp. 249-256). Cambridge: Cambridge University Press. doi:10.1017/9781316216491.043
- Bathelt, J., Gathercole, S. E., Johnson, A., & Astle, D. E. (2017). Differences in brain morphology and working memory capacity across childhood. *Developmental Science*. e12579.
- Bjorklund, D. F. (1987). A note on neonatal imitation. *Developmental Review, 7*(1), 86-92.
- Cohen, J. (1992). A power primer. *Psychological Bulletin, 112*, 155-159.
- Dettmer, A. M., Kaburu, S. S., Simpson, E. A., Paukner, A., Sclafani, V., Byers, K. L., . . . Miller, G. M. (2016). Neonatal face-to-face interactions promote later social behaviour in infant rhesus monkeys. *Nature communications, 7*: 11940.
- Dumas, G., Kelso, J. S., & Nadel, J. (2014). Tackling the social cognition paradox through multi-scale approaches. *Frontiers in Psychology, 5*: 882. Doi: 10.3389/fpsyg.2014.00882
- Ferrari, G. A., Nicolini, Y., Demuru, E., Tosato, C., Hussain, M., Scesa, E.,...& Ferrari, P. F. (2016). Ultrasonographic investigation of human fetus responses to maternal communicative and non-communicative stimuli. *Frontiers in Psychology, 7*:354. 10.3389/fpsyg.2016.00354

- Ferrari, P. F., Paukner, A., Ruggiero, A., Darcey, L., Unbehagen, S., & Suomi, S. J. (2009). Interindividual differences in neonatal imitation and the development of action chains in rhesus macaques. *Child Development, 80*, 1057-1068.
- Ferrari, P. F., Vanderwert, R. E., Paukner, A., Bower, S., Suomi, S. J., & Fox, N. A. (2012). Distinct EEG amplitude suppression to facial gestures as evidence for a mirror mechanism in newborn monkeys. *Journal of Cognitive Neuroscience, 24*, 1165-1172.
- Ferrari, P. F., Visalberghi, E., Paukner, A., Fogassi, L., Ruggiero, A., & Suomi, S. J. (2006). Neonatal imitation in rhesus macaques. *PLoS Biology, 4*(9), e302.
- Field, T. M., Woodson, R., Greenberg, R., & Cohen, D. (1982). Discrimination and imitation of facial expression by neonates. *Science, 218*(4568), 179-181.
- Gardner, J., & Gardner, H. (1970). A note on selective imitation by a six-week-old infant. *Child Development, 41*, 1209-1213.
- Hayes, L. A., & Watson, J. S. (1981). Neonatal imitation: fact or artifact? *Developmental Psychology, 17*(5), 655.
- Heimann, M. (1989). Neonatal imitation, gaze aversion, and mother-infant interaction. *Infant Behavior and Development, 12*(4), 495-505.
- Heimann, M. (1991). Neonatal imitation: A social and biological phenomenon. In T. Archer & S. Hansen (eds.), *Behavioral Biology: Neuroendocrine axis* (pp. 173-186). Hillsdale, N.J.: Lawrence Erlbaum.
- Heimann, M. (2001). Neonatal imitation-a "fuzzy" phenomenon. In C. v. Hofsten, F. Lacerda, & M. Heimann (Ed.), *Emerging cognitive abilities in early infancy* (pp. 231-246). Mahwah, NJ: Erlbaum.
- Heimann, M. (2002). Notes on individual differences and the assumed elusiveness of neonatal imitation. In A. N. Meltzoff & W. Prinz (eds.), *The imitative mind: Development, evolution, and brain bases* (pp. 74-84). Cambridge, UK: Cambridge University Press.

- Heimann, M., Nelson, K. E., & Schaller, J. (1989). Neonatal imitation of tongue protrusion and mouth opening: methodological aspects and evidence of early individual differences. *Scandinavian Journal of Psychology*, 30, 90-101.
<https://doi.org/10.1111/j.1467-9450.1989.tb01072.x>
- Heimann, M., & Schaller, J. (1985). Imitative reactions among 14-21 days old infants. *Infant Mental Health Journal*, 6, 31-39.
- Heyes, C. (2016). Homo imitans? Seven reasons why imitation couldn't possibly be associative. *Philosophical Transactions of the Royal Society B*, 371(1686), 20150069.
- Holmlund, C. (1995). Development of turntakings as a sensorimotor process in the first 3 months: A sequential analysis. In K.E. Nelson & Z. Réger (eds.) *Children's Language volume 8* (pp. 41-64). Hillsdale, NJ: Lawrence Erlbaum.
- Jones, S. (1996). Imitation or exploration? Young infants' matching of adults' oral gestures. *Child Development*, 67, 1952 – 1969. <https://doi.org/10.1111/j.1467-8624.1996.tb01837.x>
- Jones, S. (2017). Can newborn infants imitate? *Wiley Interdisciplinary Reviews: Cognitive Science*. doi:10.1002/wcs.1410
- Keven, N., & Akins, K. A. (2017). Neonatal imitation in context: Sensorimotor development in the perinatal period. *Behavioral and Brain Sciences*, 40, e381.
- Koepke, J. E., Hamm, M., Legerstee, M., & Russell, M. (1983). Neonatal imitation: Two failures to replicate. *Infant Behavior and Development*, 6, 97-102.
- Kugiumutzakis, G. (1998). Neonatal imitation in the intersubjective companion space. In S. Bråten (Ed.), *Intersubjective communication and emotion in early ontogeny* (pp. 63-88). Cambridge, UK: Cambridge University Press.

- Kuhl, P. K., & Meltzoff, A. N. (1996). Infant vocalizations in response to speech: Vocal imitation and developmental change. *Journal of the Acoustical Society of America*, 100, 2425 – 2438.
- Maratos, O. (1973). *The origin and development of imitation in the first six months of life*. Paper presented at the Annual Meeting of the British Psychological Society, Liverpool, UK.
- Marshall, P. J., & Meltzoff, A. N. (2014). Neural mirroring mechanisms and imitation in human infants. *Philosophical Transactions of the Royal Society: Biological Sciences*, 369: 20130620.
- Marshall, P. J., & Meltzoff, A. N. (2015). Body maps in the infant brain. *Trends in Cognitive Sciences*, 19(9), 499-505.
- Meltzoff, A. N. (1990). Towards a developmental cognitive science. The implications of cross-modal matching and imitation for the development of representation and memory in infancy. *Annual Report of the New York Academy of Science*, 608, 1-31.
<https://doi.org/10.1111/j.1749-6632.1990.tb48889.x>
- Meltzoff, A. N. (2007). 'Like me': a foundation for social cognition. *Developmental Science*, 10, 126-134.
- Meltzoff, A. N. (2017). Elements of a comprehensive theory of infant imitation. *Behavioral and Brain Sciences*, 40, e396.
- Meltzoff, A. N., & Moore, M. K. (1977). Imitation of facial and manual gestures by human neonates. *Science*, 198(4312), 75-78.
- Meltzoff, A. N., & Moore, M. K. (1983a). Newborn infants imitate adult facial gestures. *Child Development*, 54, 702-709.

- Meltzoff, A. N., & Moore, M. K. (1983b). The origins of imitation in infancy: Paradigm, phenomena, and theories. In L. P. Lipsitt (Ed.), *Advances in Infancy Research Volume 2* (pp. 265–301). Norwood, NJ: Ablex Publishing Corporation.
- Meltzoff, A. N., & Moore, M. K. (1989). Imitation in newborn infants: Exploring the range of gestures imitated and the underlying mechanisms. *Developmental Psychology*, 25, 954-962.
- Meltzoff, A. N., & Moore, M. K. (1994). Imitation, memory, and the representation of persons. *Infant Behavior and Development*, 17, 83-99.
- Meltzoff, A. N., & Moore, M. K. (1997). Explaining facial imitation: A theoretical model. *Early Development and Parenting*, 6, 179-192.
- Meltzoff, A.N. & Marshall, P.T. (2018). Human infant imitation as a social survival circuit. *Current Opinion in Behavioral Sciences*, 24,130–136.
<https://doi.org/10.1016/j.cobeha.2018.09.006>
- Meltzoff, A.N., Murray, L., Simpson, E., Heimann, M., Nagy, E., Nadel, J.... & Ferrari, P. F. (2018). Re-examination of Oostenbroek et al. (2016) – Evidence for neonatal imitation of tongue protrusion. *Developmental Science*, 21:e12609.
- Meltzoff, A. N., Ramírez, R. R., Saby, J. N., Larson, E., Taulu, S., & Marshall, P. J. (2018). Infant brain responses to felt and observed touch of hands and feet: an MEG study. *Developmental Science*, e12651.
- Meltzoff, A. N., Saby, J. N., & Marshall. P. J, (2018). Neural representations of the body in 60-day-old human infants. *Developmental Science*, e12698.
- Murray, L., Pascalis, L. De, Bozicevic, L., Hawkins, L., Scafani, V., & Ferrari, P. F. (2016). The functional architecture of mother-infant communication, and the development of infant social expressiveness in the first two months. *Scientific Reports*, 6:39019.

- Nagy, E. (2011). The newborn infant: A missing stage in developmental psychology. *Infant and Child Development*, 20, 3– 19.
- Nagy, E., Kompagne, H., Orvos, H., & Pal, A. (2007). Gender-related differences in neonatal imitation. *Infant and Child Development*, 16, 267-276.
- Nagy, E., & Molnar, P. (2004). Homo imitans or homo provocans? Human imprinting model of neonatal imitation. *Infant Behavior and Development*, 27(1), 54-63.
- Nagy, E., Pal, A., & Orvos, H. (2014). Learning to imitate finger movements by the human neonate, *Developmental Science*, 17, 841 – 857.
- Nagy, E., Pilling, K., Orvos, H., & Molnar, P. (2013). Imitation of tongue protrusion in human neonates: Specificity of the response in a large sample. *Developmental Psychology*, 49, 1628-1638.
- Oostenbroek, J., Suddendorf, T., Nielsen, M., Redshaw, J., Kennedy-Costantini, S., Davis, J., . . . Slaughter, V. (2016). Comprehensive longitudinal study challenges the existence of neonatal imitation in humans. *Current Biology*, 26, 1334-1338.
- Paukner, A., Simpson, E. A., Ferrari, P. F., Mrozek, T., & Suomi, S. J. (2014). Neonatal imitation predicts how infants engage with faces. *Developmental Science*, 17, 833-840.
- Pitti, A., Kuniyoshi, Y., Quoy, M., & Gaussier, P. (2013). Modeling the minimal newborn's intersubjective mind: The visuotopic-somatotopic alignment hypothesis in the superior colliculus. *PLoS One*, 8(7), e69474.
- Previc, F.H., Declerck, C., & de Brabander, B. (2005). Why your “head is in the clouds” during thinking: The relationship between cognition and upper space. *Acta Psychologica*, 118, 7–24.
- Rayson, H., Bonaiuto, J. J., Ferrari, P. F., & Murray, L. (2017). Early maternal mirroring predicts infant motor system activation during facial expression observation. *Scientific Reports*, 7: 11738

- Rönnqvist, L., & von Hofsten, C. (1994). Neonatal finger and arm movements as determined by a social and an object context. *Early Development and Parenting*, 3(2), 81-94.
- Saby, J. N., Meltzoff, A. N., & Marshall, P. J. (2015). Neural body maps in human infants: Somatotopic responses to tactile stimulation in 7-month-olds. *Neuroimage*, 118, 74-78.
- Simpson, E. A., Maylott, S.A., Heimann, M., Subiaul, F., Paulkner, A., Soumi, S.J., & Ferrari, P.F. (2017). Animal studies help clarify misunderstandings about neonatal imitation. *Behavioral and Brain Sciences*, 40, e400.
- Simpson, E. A., Miller, G. M., Ferrari, P. F., Suomi, S. J., & Paukner, A. (2016). Neonatal imitation and early social experience predict gaze following abilities in infant monkeys. *Scientific Reports*, 6, 20233.
- Simpson, E. A., Murray, L., Paukner, A., & Ferrari, P. F. (2014). The mirror neuron system as revealed through neonatal imitation: presence from birth, predictive power and evidence of plasticity. *Philosophical Transactions of the Royal Society B*, 369(1644), 20130289.
- Stein, B. E., & Meredith, M. A. (1993). *The merging of the senses*: Cambridge, MA: MIT Press.
- Stein, B. E., & Stanford, T. R. (2013). Development of the superior colliculus/optic tectum. In J. L. R Rubesteing & P. Rakic (eds.), *Comprehensive developmental neuroscience: Neural circuit development and the function in the healthy and diseased brain* (pp. 41 – 59). London, UK: Academic Press.
- Theorell, K., Prechtl, H. F. R., Blair, A. W., & Lind, J. (1973). Behavioural state cycles of normal newborn infants: A comparison of the effect of early and late cord clamping. *Developmental Medicine and Child Neurology*, 15, 597 – 605.
- <https://doi.org/10.1111/j.1469-8749.1973.tb05170.x>

- Trevarthen, C. (2011). What is it like to be a person who knows nothing? Defining the active intersubjective mind of a newborn human being. *Infant and Child Development*, 20, 119-135.
- Trevarthen, C., & Aitken, K. (2003). Regulation of brain development and age-related changes in infants' motives: The developmental function of regressive periods. In M. Heimann (ed.) *Regression Periods in Human Infancy* (pp. 107-184). Mahwah, NJ: Erlbaum.
- Ullman, M. T. (2016). The Declarative/Procedural Model: A neurobiological model of language learning, knowledge, and use. In G. Hickok & S.L. Small (eds.), *Neurobiology of Language* (pp. 953 – 968). Amsterdam: Academic Press.
- Vincini, S., Jhang, Y., Buder, E. H., & Gallagher, S. (2017), Neonatal imitation: Theory, experimental design, and significance for the field of social cognition. *Frontiers of Psychology*, 8:1323.
- Vinter, A. (1986). The role of movement in eliciting early imitations. *Child Development*, 57, 66-71.
- Wolff, P.H. (1987). *Behavioral States and Expression of Emotions in Early Infancy*. Chicago: University of Chicago Press.
- Zazzo, R. (1957). Le problème de l'imitation chez le nouveau-né. *Enfance*, 10(2), 135-142.

Table 1.

Overview of the Procedure: Imitation of Tongue Protrusion and Mouth Opening were each Observed Separately During Six Intervals: Three Dynamic Demonstrations when the Target Gesture was Modeled (TP or MO; intervals 1, 3, and 5) Intertwined with Three Passive Face Periods (intervals 2, 4, and 6).

Interval	Gesture A ¹						Pause ²	Gesture B ¹					
	1	2	3	4	5	6		1	2	3	4	5	6
Stimuli	Demo	Passive	Demo	Passive	Demo	Passive		Demo	Passive	Demo	Passive	Demo	Passive
Length (s)	20	20	20	20	20	20		20	20	20	20	20	20
Cumulative time (s)	20	40	60	80	100	120		20	40	60	80	100	120

^aThe order was counterbalanced: for half of the infants TP was presented as the first gesture during the experiment, else MO was presented first. ²The length of the pause was 20 to 30 seconds or longer if the infant was not in an alert state.

Table 2.

The Mean length in seconds (s) of the three Demonstration Periods and the Amount of Time the Infants were Judged to be Attentive to the Model (looking toward the face) when the Gesture was Presented.

	Demo intervals		Looking ¹ (s)					
Gesture modeled	Length in s		At model		Away		<i>P</i> ² <	<i>d</i>
	M	SD	M	SD	M	SD		
Tongue protrusion	20.1	3.1	11.6	6.2	7.7	5.9	.01	0.64
Mouth opening	20.3	3.1	12.4	5.5	6.5	4.4	.01	1.18

¹ Not codable frames (e.g. eyes closed): On average 0.6 s for tongue protrusion and 1.4 s for mouth opening; ² Comparison between looking at model or away (t-test).

Table 3.

Imitation of *Tongue Protrusion* (TP): Rate of TP per second after Modeling of TP (TP Demo)
 Compared with when Mouth Opening was Modeled (MO Demo) During Three Different
 Time Windows.

Time Window ^a	N	Interval ^b	Rate of TP/s				ANOVA (<i>p</i> -value)		
			TP Demo		MO Demo		C ^c	T ^d	CxT ^e
			M	SD	M	SD			
1. Early, 0-60s	30	1, 2 and 3	.091	.079	.059	.063	.036	.040	ns
2. Middle, 61-100s	26	4 and 5	.097	.086	.090	.012	.ns	.ns	.037
3. Late, 101-120s	24	6	.056	.074	.083	.094	-	ns	-

Notes. ^a Only children judged to be alert within the complete time window are included in the analysis; thus the variation in N. ^b Each interval is 20-s long; ^c C = condition; ^d T = time; ^e CxT interaction.

Table 4.

Imitation of *Mouth Opening* (MO): Rate of MO per second after Modeling of MO (MO Demo) Compared with when Tongue Protrusion was Modeled (TP Demo) During Three Different Time Windows.

Time Window ^a	N	Interval ^b	Rate of MO/s				ANOVA (<i>p</i> -value)		
			MO Demo		TP Demo		C ^c	T ^d	CxT ^e
			M	SD	M	SD			
1. Early, 0-60s	31	1, 2 and 3	.109	.085	.130	.083	ns	ns	ns
2. Middle, 61-100s	27	4 and 5	.162	.109	.119	.076	.030	ns	ns
3. Late, 101-120s	26	6	.136	.126	.139	.125	-	ns	-

Notes. ^a Only children judged to be alert within the complete time window are included in the analysis; thus the variation in N. ^b Each interval is 20-s long; ^cC = condition; ^dT = time; ^eCxT interaction.

Table 5.

Mean Frequencies Across the Three Demo and the Three Passive Face Intervals for *Tongue Protrusion* During the Tongue Protrusion Part of the Experiment and for *Mouth Opening* During the Mouth Opening Part of the Experiment.

Behavior	N ¹	Intervals				<i>t</i>	<i>p</i> =
		Demo ²		Passive Face ³			
		M	SD	M	SD		
Tongue Protrusion	24	1.97	1.63	1.79	1.27	-.644	ns
Mouth Opening	26	2.47	1.78	2.81	1.65	1.23	ns

Notes. ¹ Only children with valid data within in all intervals are included in the analysis; ² Demo = intervals when the gesture is modeled; ³ Passive = the experimenter keeps a passive face.

Table 6.

Number of Infants Displaying an Imitative (+), Non-Imitative (-) or an Equal Response (0)

Pattern During the Three Time Windows^a in the Experiment

Time windows (TW) ^a	Individual response pattern						
	Imitation			Unclear		No imitation	
	(+) ^b			(0) ^b		(-) ^b	
	N	n	%	n	%	n	%
<i>Tongue protrusion</i>							
TW1	30	16	53.3	5	16.7	9	30.9
TW2	26	11	42.3	4	15.4	11	42.3
TW3	24	7	29.2	8	33.3	9	37.5
<i>Mouth opening</i>							
TW1	31	11	35.5	2	6.5	18	58.1
TW2	27	18	66.7	1	3.7	8	29.6
TW3	26	10	38.5	4	15.4	12	46.2

Notes. ^aTW1 = 0 - 60s, TW2 = 61-100s, and TW3 = 101-120s, see text for detailed information; ^b A plus (+) sign indicates imitation; a minus (-) sign indicates that the frequency of the target behavior is higher in the control condition, and a zero (0) that the frequency is equal in both the imitation and control condition.

Figure Captions

Figure 1.

Mean Frequencies (Y-axis) of *Tongue Protrusion* (TP) Plotted Over the Six Intervals Constituting the Three Time Windows (TW) used in the Main Analysis. Observed Frequency of TP when TP is Demonstrated (TP Demo) is Represented by a Blue Line and the Mean Frequencies of TP when Mouth Opening is Demonstrated (MO Demo) is Represented by the Red Line. Intervals 1, 3 and 5 Represents Active Presentation of the Gesture while Intervals 2, 4, and 6 Depicts Periods when the Experimenter Keeps a Passive Face. Error bars = SE.

Figure 2.

Mean Frequencies (Y-axis) of *Mouth Opening* (MO) Plotted Over the Six Intervals Constituting the Three Time Windows (TW) used in the Main Analysis. Observed Frequency of MO when MO is Demonstrated (MO Demo) is Represented by a Blue Line and the Mean Frequency of MO when Tongue Protrusion is Demonstrated (TP Demo) is Represented by the Red Line. Intervals 1, 3 and 5 Represents Active Presentation of the Gesture while Intervals 2, 4, and 6 Depicts Periods when the Experimenter Keeps a Passive Face. Error bars = SE.

Figure 1

Figure 2