Immune maturation and modulation in childhood allergies

Aspects of epigenetic, mucosal
and systemic immune mediators in
allergy development and
prevention

Linköping University Medical Dissertations

No. 1755

Immune maturation and modulation in childhood allergies

Aspects of epigenetic, mucosal and
systemic immune mediators in allergy
development and prevention

Johanna Huoman

Division of Inflammation and Infection
Department of Biomedical and Clinical Sciences
Faculty of Medicine and Health Sciences, Linköping University,
SE-581 83 Linköping, Sweden

Linköping 2020

© Johanna Huoman 2020

All rights reserved

Artwork on cover by Jessica Andersson. Illustrations in thesis by Johanna Huoman.

Papers II and IV were published by the Journal of Pediatric Allergy and Immunology, and have been re-printed with permission from John Wiley & Sons Ltd.

ISBN 978-91-7929-787-9

ISSN 0345-0082

Printed by LiU-tryck, Linköping, Sweden 2020

“If we knew what it was we were doing,
it would not be called research, would it?”

Albert Einstein

”Utan tvivel är man inte klok.”

Tage Danielsson

“You fail to recognise that it matters
not what someone is born,
but what they grow to be.”

Albus Dumbledore

Harry Potter and the Goblet of Fire

Main supervisor

Professor Maria Jenmalm, Linköping University, Sweden

Co-supervisor

Professor Lennart Nilsson, Allergy Center and Linköping University, Sweden

Faculty opponent

Professor Petra Ina Pfefferle, Phillips University of Marburg, Germany

Funding

The work presented in this thesis was supported by the Swedish Research Council, the Swedish Heart and Lung Foundation, the Cancer and Allergy Foundation, the United Kingdom Medical Research Council, the Manchester Biomedical Research Centre, the Foundation Samariten, the Ellen, Walter & Lennart Hesselman foundation, the Ekhaga Foundation, the Åke Viberg Foundation, the Olle Engkvist Foundation, the Medical Research Council for South-East Sweden, the Swedish Asthma and Allergy Association, the Cancer and Allergy Association, the Konsul Th C Berghs Foundation for Scientific Research, the Research and Scholarship Foundation in Östergötland, and the University Hospital of Linköping, Sweden.

Original publications and manuscripts

- Paper I Circulating chemokine levels and the development of allergic phenotypes from infancy to adolescence: a population-based birth cohort study

Johanna Huoman*, Sadia Haider*, Angela Simpson, Clare S Murray, Adnan Custovic & Maria C Jenmalm

*shared first authorship

Manuscript – submitted to Journal of Allergy and Clinical Immunology

- Paper II Pre- and postnatal *Lactobacillus reuteri* treatment alters DNA methylation of infant T helper cells

Anna Forsberg*, **Johanna Huoman***, Simon Söderholm, Ratnesh Bhai Mehta, Lennart J Nilsson, Thomas R Abrahamsson, Jan Emerudh, Mika Gustafsson & Maria C Jenmalm

*shared first authorship

Pediatric Allergy and Immunology 2020;31:544–553

- Paper III Epigenetic effects of pre- and postnatal *Lactobacillus reuteri* and ω -3 supplementation in neonatal T helper cells

Johanna Huoman, David Martínez-Enguita, Elin Olsson, Jan Emerudh, Lennart J Nilsson, Karel Duchén, Mika Gustafsson & Maria C Jenmalm

Manuscript

- Paper IV Sublingual immunotherapy alters salivary IgA and systemic immune mediators in timothy allergic children

Johanna Huoman, Georgia Papapavlou, Anna Pap, Johan Alm, Lennart J Nilsson, & Maria C Jenmalm

Pediatric Allergy and Immunology 2019;30:522–530

Other relevant publications

Does allergy begin *in utero*?

Gabrielle A Lockett, ***Johanna Huoman*** & John W Holloway

Pediatric Allergy and Immunology 2015;26:394–402

Abstract

The prevalence of allergic diseases has in the past century increased among children in affluent societies. Underlying causes are incompletely disentangled, but decreased diversity in environmental and microbial exposures could drive allergy development. Allergic individuals possess imbalanced immune responses, skewed in favour of Th2 cells along with lesser Th1- and Treg-responses. As allergy development early in life increases the risk of developing further allergic manifestations later, early prevention is key. Thus, interventions in pregnancy, early life and childhood may modulate immunity towards tolerance, although underpinnings of immune maturation and modulation in allergy prevention throughout childhood are not entirely understood. In this thesis, these questions are addressed in children with a high propensity of developing allergic disease or who already have manifested allergies.

Chemokines are crucial for immune cell recruitment to the allergic reaction site, and associate with allergy development in childhood. In Paper I, circulating levels of the allergy-related chemokines CCL17, CCL18, CCL22, CXCL10 and CXCL11 were studied in the natural course of allergic disease. Elevated levels of the Th2/Treg-regulated chemokine CCL18 in infancy and childhood associated with development of asthma and/or sensitisation. Moreover, this finding conferred higher odds of developing asthma and sensitisation from early school age until adolescence. Additionally, increased levels of the Th1-associated chemokines CXCL10 after birth, and decreased levels of CXCL11 at birth, preceded asthma development later in life. Hence, Paper I showed that circulating chemokine levels in different ways precede allergy development.

Epigenetic modifications, such as DNA methylation, comprise a link between the genetic setup and environmental exposures, and regulate processes such as Th cell differentiation. Perinatal treatment with *Lactobacillus reuteri* and ω -3 fatty acids prevent development of some IgE-mediated manifestations. However, the drivers of the immunostimulatory and pro-resolving effects of these treatments are sparsely examined. In Papers II and III, epigenome-wide DNA methylation patterns in CD4+ cells upon pre- and postnatal *L. reuteri* supplementation alone or in combination with ω -3 fatty acids were studied. In Paper II, the greatest epigenome-wide differential methylation was evident at birth, mainly directed towards hypomethylation, indicating transcriptional availability of affected genes. Network analyses revealed several immune-related pathways, and a relationship of differentially methylated genes to allergy development. Thus, prenatal *L. reuteri* treatment seemingly poises Th cells towards immune activation at birth, possibly influencing immune maturation as well as allergy development in the child.

In Paper III, epigenome-wide DNA methylation patterns were surveyed at birth. In this ongoing trial, mothers are treated during the latter half of pregnancy with a combination of *L. reuteri* and ω -3 fatty acids. Four different treatment groups were studied, and the largest differential methylation was seen in the double active treatment group. In contrast to Paper II, most CpGs and genes were hypermethylated, indicating repressed gene transcription. In line with Paper II, network analyses showed that T cell- and immune-mediated pathways were affected by treatment, and synergistic effects of the double treatment were indicated. Taken together, prenatal treatment with *L. reuteri* and/or ω -3 fatty acids altered the epigenome to

different extents at birth, mainly towards hypermethylation, and often affected immune-related pathways.

Immunomodulatory effects of sublingual immunotherapy in children and adolescents are scarcely investigated. In Paper IV, circulating and salivary immune mediators were investigated in timothy grass pollen-allergic children treated with sublingual immunotherapy. Actively treated children had elevated levels of timothy grass pollen-specific IgA-antibodies in saliva, along with increased circulating levels of the Th1-associated chemokines CXCL10 and CXCL11, both after treatment ending and two years later. Taken together, sublingual immunotherapy modulates local and peripheral immune responses in children with timothy grass pollen-induced allergy, by augmenting Th1-responses, lessening Th2-responses and inducing immunomodulatory responses, suggesting induction of tolerance, also partly in the long-term.

Altogether, the studies in this thesis have shown altered immune regulation in children developing allergies. Moreover, immunomodulatory effects of prenatal treatment with probiotics and ω -3 fatty acids, and sublingual immunotherapy in children with grass pollen-induced allergic disease, were revealed. DNA methylation patterns and immunologic mediators in blood and saliva could potentially serve as appropriate biomarkers for allergic disease. Long-term health benefits can be reached by intervening early in life, and further knowledge about the mechanisms behind this could promote the prevention of allergic diseases and hence improve the quality of life for children and adolescents.

Populärvetenskaplig sammanfattning

Förekomsten av allergiska sjukdomar, som böjveckseksem, hösnuva och astma, har under det senaste århundradet ökat markant bland barn i industrialiserade samhällen. De bakomliggande orsakerna är inte helt klarlagda, men samhällseliga förändringar har minskat vår mångfaldiga exponering för bakterier, virus och parasiter. Detta skulle kunna ligga till grund för immunförsvarets felaktiga reaktion mot egentligen ofarliga ämnen som ses vid allergier. Hos allergiska individer är immunförsvaret obalanserat, med en relativ övervikt av det så kallade T-hjälpar-2 (Th2)-svaret gentemot Th1- och det regulatoriska T-cells (Treg)-svaret. Allergiska sjukdomar utvecklas ofta tidigt i livet, vilket ökar risken för att utveckla vidare allergier senare i livet. Därför är det viktigt att motverka den allergiska marschens framfart tidigt genom förebyggande behandlingar. Ett tillvägagångssätt är att påbörja behandling under graviditeten och tidiga barndomen hos barn med hög risk för att bli allergiska, då grunden för immunsystemet läggs redan under fosterlivet. För redan utvecklade allergier är det tänkbart att omforma dessa immunsvår med immunterapi, som kan minska symptom av befintliga allergier samtidigt som det är möjligt att motverka utvecklingen av senare allergier. Det är dock inte helt klarlagt hur immunutmognaden under barndomen är reglerad, eller hur dessa typer av behandlingar skulle kunna påverka allergiutveckling under den perioden. I denna avhandling undersöks immunutmognad vid allergiutveckling hos barn, och möjliga immunmodulerande förebyggande behandlingar hos barn med genetisk benägenhet att bli allergiska eller som redan utvecklat allergisk sjukdom.

För att celler ska rekryteras till platsen för en allergisk reaktion krävs bland annat s.k. kemokiner. I det första arbetet undersöktes dessa lockelseämnen, då våra tidigare studier visat att nivåerna av vissa kemokiner vid födseln förutspår utvecklingen av allergi hos barn. De allergirelaterade kemokinerna CCL17, CCL18, CCL22, CXCL10 och CXCL11 analyserades i blodprover vid födseln, 1 och 8 års ålder hos barn från en populationsbaserad observationsstudie. Förhöjda nivåer av CCL18, ett kemokin under reglering av både Th2- och Treg-svar, uppmättes vid 1 och/eller 8 års ålder hos barn som hade astma (särskilt svår astma) och/eller var sensibiliserade. De ökade nivåerna associerade också till högre odds för utveckling av astma från tidig skolålder upp till övre tonåren, med liknande resultat för sensibilisering. Även ökade nivåer av de Th1-associerade kemokinerna CXCL10 efter födseln och minskade nivåer av CXCL11 vid födseln föregick utvecklingen av astma senare i livet. Det första arbetet visade alltså på att cirkulerande kemokiner på olika vis föregår utvecklingen av allergier hos barn och ungdomar.

Som länk mellan arv och miljö står s.k. epigenetiska modifieringar, vilka reglerar genaktiviteten utan att förändra den genetiska koden i arvsmassan. Till dessa modifieringar räknas DNA-metylering, en process som bl.a. styr utmognad av de allergirelaterade T-hjälparcellerna. Vi har i tidigare separata studier med den probiotiska stammen *Lactobacillus reuteri* och omega-3-behandling visat förebyggande av vissa IgE-medierade allergier. Vad som föranleder de immunstimulerande och immunmodulerande effekterna av behandlingarna är dock otillräckligt undersökt. I det andra och tredje arbetet undersöktes hur *L. reuteri* separat eller i kombination med omega-3-fettsyror påverkar DNA-metyleringsmönster i CD4⁺ Th-celler hos barn som behandlats före och efter födseln. I det andra arbetet undersöktes DNA-metyleringsmönster både lokalt och i hela genomet vid födseln, ett och två års ålder. Behandling med *L. reuteri* förändrade DNA-metyleringsmönster i allergirelaterade T-hjälparceller mest vid födseln mot s.k.

hypometylering, vilket pekar på ökad tillgänglighet av generna för proteinuttryck. Vidare nätverksanalyser visade att flera immunrelaterade processer påverkades av behandlingen. Därtill var generna från nätverket till stor del associerade med allergiutveckling. Maternell behandling med *L. reuteri* under den sista graviditetsmånaden tycks alltså förändra DNA-metyleringsmönster i T-hjälparceller hos fostret mot ökad immunaktivering vid födseln, vilket i sin tur skulle kunna påverka både immunutmognad och allergiutveckling hos barnet.

I likhet med det andra arbetet undersöktes i det tredje arbetet DNA-metyleringsmönster i hela epigenomet, fast endast vid födseln. I denna pågående studie behandlas mödrarna under den andra halvan av graviditeten med en kombination av *L. reuteri* och omega-3-fettsyror. Fyra olika behandlingsgrupper undersöktes och den största förändringen i DNA-metylering återfanns i den kombinerade aktiva behandlingsgruppen. I motsats till det andra arbetet var dock de flesta CpG positionerna och generna hypermetylerade, vilket tyder på att genernas tillgänglighet för proteinuttryck hämmas. I linje med det andra arbetet framkom T-cells- och immunrelaterade signalvägar i nätverksanalyser på dessa gener, och det fanns indikationer på synergistiska effekter mellan behandlingarna. Det tredje arbetet visade att behandling med *L. reuteri* och/eller omega-3-fettsyror under senare delen av graviditeten förändrar T-hjälparcellernas epigenom i olika grad främst mot hypermetylering, och ofta påverkar immunrelaterade signalvägar. Relevansen av dessa fynd kommer i framtida studier att undersökas på proteinnivå och i relation till allergiutveckling.

Med allergenspecifik immunterapi är det möjligt att modulera immunsvaret hos allergiska individer mot ett tolerant immunsvaret, men effekter av sublingual immunterapi på immunförsvaret hos barn och ungdomar är knapphändigt undersökta. I det fjärde arbetet undersöktes olika immunologiska mediatorer i blod och saliv hos barn med gräspollenallergi, som deltagit i en studie med sublingual immunterapi. Nivåerna av allergirelaterade cytokiner och kemokiner undersöktes i blodprover från inklusionstillfället, efter tre år med behandling samt två år efter avslutad behandling i plasmaprover och allergenstimulerade blodceller. Dessutom mättes total-IgA, sekretoriskt IgA och gräspollensspecifikt IgA i saliv vid samma tillfällen. Barn som fått aktiv behandling hade högre nivåer av gräspollensspecifika IgA-antikroppar i saliv både när behandlingen avslutades och två år efter. Därtill ökade nivåerna av de Th1-associerade kemokinerna CXCL10 och CXCL11 i blodet vid samma tidpunkter. Sammantaget visade resultaten från det fjärde arbetet att behandlingen med sublingual immunterapi hos barn med gräspollenallergi modulerar immunsvaret både lokalt och i cirkulationen genom att öka Th1-svar, minska Th2-svar och inducera immunreglerande svar, vilket indikerar att tolerans har utvecklats hos dessa barn, delvis även på lång sikt.

Sammanfattningsvis har studierna i denna avhandling visat på förändrad immunreglering hos barn som utvecklar allergi. Dessutom påvisades immunmodulerande effekter av prenatal behandling med probiotika och omega-3-fettsyror samt av sublingual immunterapi hos barn med gräspollenallergi. DNA-metyleringsmönster och immunologiska mediatorer i blod och saliv skulle kunna fungera som lämpliga biomarkörer för allergisk sjukdom, vilket är ett viktigt led i att kunna förutsäga allergiutveckling och förbättra den kliniska behandlingen av allergier bland barn och ungdomar. Långsiktiga hälsofördelar kan uppnås genom att ingripa tidigt i livet, och vidare kunskap om mekanismerna bakom detta skulle kunna främja förebyggandet av allergiska sjukdomar och således kunna förbättra livskvaliteten för barn och ungdomar.

Table of contents

Abbreviations	1
Prologue · Early beginnings - the case of an allergic child	3
Background	4
The allergy epidemic – a consequence of microbial dysbiosis?	4
Immunity and inflammation – endogenous protection in a world full of dangers	6
T helper cells – friends and foes in health and disease	8
Allergy development – dysregulated inflammatory responses to innocuous antigens	12
Epigenetic regulation – transcriptional modifications modulating interplay between genome and environment	15
Chemokines – key orchestrators in immune cell recruitment	18
Perinatal immune maturation – the importance of maternal exposures during pregnancy and early life	22
Mucosal immunity – immunological barriers balancing host defence and tolerance induction	24
Commensal microbiota – symbiotic bacteria important for appropriate immune maturation	27
IgA-antibodies – mucosal drivers of host defence and tolerance induction	29
Combating the allergy epidemic – approaches to immunomodulating treatment	31
Probiotics – amongst all microbes, who's a good bug?	32
ω -3 fatty acids – modulators of inflammatory responses, as a matter of fat	34
Immunotherapy – re-creating tolerance by desensitisation	36
Aims and hypotheses of the thesis	39
Study design and methodological considerations	40
Paper I – Chemokine studies in the Manchester Asthma and Allergy Study (MAAS)	40
Paper II – DNA methylation studies in the <i>Lactobacillus reuteri</i> (<i>L. reuteri</i>) cohort	41
Paper III – DNA methylation studies in the PRObiotics and OMega-3 (PROOM-3) study	43
Paper IV – Immune response studies in the GRAZAX® Asthma Prevention (GAP) trial	46
Ethical considerations	48
Methodological foundations	52
Enzyme-linked immunosorbent assay (ELISA)	52
Multiplex bead assay (Luminex)	52
Flow cytometry	53
DNA methylation analyses - Th subset locus-specific and epigenome-wide approaches	53
Allergen stimulation of peripheral blood mononuclear cells	54
Methodological considerations	55
Statistical and bioinformatic analyses	56

Results and discussion	60
Paper I · Circulating chemokine levels in infancy and childhood predict outcomes of allergic disease later in life.....	60
Childhood CXCL10 and CCL18 levels influence present and future development of sensitisation	60
Neonatal CXCL11 and childhood CCL18 levels predict different outcomes in terms of asthma development	61
Infant and childhood levels of CCL18 predict long-term development of sensitisation and asthma.....	63
Paper II · Prenatal intervention with <i>L. reuteri</i> modulates epigenetic landscape in cord blood CD4+ cells.....	65
Probiotic intervention leads to greatest epigenome-wide changes at the time of birth and affects immune-related pathways.....	65
Network analyses reveal relationships between DMGs, immunological pathways and allergy development upon probiotic intervention	67
Paper III · Differential epigenomic effects of combined prenatal intervention with <i>L. reuteri</i> and ω -3 fatty acids in cord blood CD4+ cells	70
Combined <i>L. reuteri</i> and ω -3 fatty acid treatment leads to greatest differential methylation effects.....	70
Single treatment with <i>L. reuteri</i> or ω -3 fatty acids show fewer but interesting findings.....	71
Shared genes between networks reveal common pathways, while unique genes provide insights into treatment-specific effects.....	72
Paper IV · Three years of SLIT alters immune responses towards tolerance in grass pollen-allergic children.....	77
<i>P. pratense</i> SLIT modulates immune responses in the oral mucosa by inducing timothy grass pollen-specific IgA-antibodies in saliva	77
Alterations in IgE-mediated mast cell degranulation and presence of circulating IgG ₄ after SLIT treatment	78
SLIT re-balanced Th1- and Th2-responses, with maintained shift after treatment ending...79	
Concluding remarks and future perspectives	82
Epilogue · Never too late – a break in the atopic march?	87
Acknowledgments	88
On the origins of doctor Huoman – or the people thanks to whom she got there at all.....	88
References.....	93

Abbreviations

AA – arachidonic acid
AD – atopic dermatitis
AIT – allergen immunotherapy
APC – antigen presenting cell
ARC – allergic rhinoconjunctivitis
ARS – allergic rhinitis score
Betula verrucosa – birch allergen
BMIQ – beta mixture quantile
Breg – regulatory B cell
CBMC – cord blood mononuclear cell
CCL – CC chemokine ligand
CCR – CC chemokine receptor
CFU – colony forming units
CpG – Cytosine-phosphate-Guanine
CXCL – CXC chemokine ligand
CXCR – CXC chemokine receptor
DC – dendritic cell
DHA – docosahexaenoic acid
DMG – differentially methylated gene
DMP – differentially methylated probe
ELISA – enzyme-linked immunosorbent assay
EPA – eicosapentaenoic acid
FDR – false discovery rate
GAP - GRAZAX[®] Asthma Prevention trial
GEE – generalised estimating equation
GW – gestational week
IFN- γ – interferon gamma
Ig – immunoglobulin
IL – interleukin

ILC – innate lymphoid cell

L. reuteri – *Lactobacillus reuteri*

LCPUFA – long chain polyunsaturated fatty acid

M cell – microfold cell

MAAS – Manchester Asthma and Allergy Study

MDS – multidimensional scaling

MHC – major histocompatibility complex

MMD – mean methylation difference

MØ – macrophage

MSD – mesoscale discovery

P. pratense – timothy grass pollen (*Phleum pratense*) allergen

PBMC – peripheral blood mononuclear cell

PEA – proximity extension assay

Phl p 1 – component 1 of *Phleum pratense* allergen

PROOM-3 – PRObiotics and OMega-3 study

PUFA – polyunsaturated fatty acid

SCIT – subcutaneous immunotherapy

SIgA – secretory IgA

SLIT – sublingual immunotherapy

SPM – specialised pro-resolving mediators

SPT – skin prick test

STAT – signal transducer and activator of transcription

SVD – singular value decomposition

TCR – T cell receptor

TGF- β – transforming growth factor beta

Th – T helper

TLR – Toll-like receptor

TNF – tumour necrosis factor

Treg – regulatory T cell

TSLP – thymic stromal lymphopoietin

VAS – visual analogue scale

Prologue

Early beginnings - the case of an allergic child

The calm of the pleasant early summer air, filled with scents of blooming flowers and the fumes of barbecues, is only interrupted with the cheerful screeches and laughs of three children playing in the backyard garden of a house. Chasing the ball they are playing with, the lively and animal loving 9-year-old Linus is a boy like many other boys his age. Living with his family in a medium-large city in Sweden, he enjoys drawing, playing outside with his two siblings and building large imaginary worlds in computer games. While on the outside a seemingly healthy child, he experiences symptoms of allergic disease that limit him in his everyday life from time to time. His mother has been allergic herself since childhood, whereas neither his father nor his siblings are allergic.

For Linus, it all began when he was a baby, when he got itching eczema on his face, torso and on his arms and legs. After examination at the paediatric allergy clinic, he was diagnosed with atopic dermatitis, and skin prick tests revealed that he was sensitised towards milk, eggs as well as cats. This meant that the family cat had to be given away to close friends of the family, and dairy products as well as egg-containing foods were removed from his diet. Along with the regular use of emollient and cortisone cream, the allergic eczema eventually disappeared. The allergic march had, however, just begun. When he at his first birthday enjoyed his grandmother's home-made birthday cake, he shortly thereafter experienced trouble breathing, got hives all over his body, and started vomiting. The cake had contained peanuts, and he had developed an anaphylactic reaction towards it. Yet another allergen had to be removed from his diet.

A couple of years later, Linus had outgrown his allergies towards milk and eggs, but instead he, to his great dismay, started to have difficulties breathing when playing for even short moments with furred pets. The presence of recurrent wheezing and coughing with obstructive bronchitis now suggested he had developed allergic asthma. This had been worsened by his recently acquired timothy grass pollen allergy, which made playing outside in the summer troublesome. As the allergic symptoms caused by the grass pollen allergy were very severe and at times exhausting to cope with, he started with immunotherapy treatment in an attempt to reduce his symptoms. Watching Linus lying in the grass, laughing until he hoops at some joke one of his siblings made, it is hard to imagine the impact of the allergic symptoms on this young boy's life. Only recently, this would not have been possible for him.

In many ways, Linus is a stereotypic case of a multi-allergic child: a boy, growing up in a Westernised society, living in an urban area, with a family history of allergies and development of a number of consecutive allergic diseases. The above, in reality imaginary but clinically highly relevant, examples of the atopic march raise some questions. What are the underlying immunological causes of allergic disease? How did Linus become this allergic, whereas for instance his siblings did not? Could the development of allergies have been prevented by intervening early in life? And is it possible to modulate immune responses of on-going allergic disease by intervention? In this thesis, attempts will be made to answer these questions, and the implications of the findings will be related to his possible benefit.

Background

The allergy epidemic – a consequence of microbial dysbiosis?

Allergic diseases, such as allergic rhinoconjunctivitis (ARC), asthma and atopic dermatitis (AD) among children and adolescents, pose a considerable economic and societal burden worldwide.¹ Previously, the prevalence of allergic disorders mainly increased among children in industrialised societies.² While the occurrence of some allergic diseases, such as asthma, may have plateaued in affluent countries, the general prevalence of allergic disorders is, however, presently still increasing among children in countries that have not adapted a Westernised lifestyle.^{3,4}

The term allergy was coined in the early 1900's, composing the combination of the two Greek words *allos*, meaning other or different, and *ergia*, energy or action, and hence defining allergy as a change in the [immune system's] capacity to react.⁵ This definition still holds true today, as allergies may be seen as misdirected reactions of the immune system to innocuous antigens, or as they are more commonly known, allergens. There are several types of hypersensitivity reactions, and immunoglobulin E (IgE)-mediated allergic symptoms constitute type I hypersensitivity reactions⁶, which will be the focus of this thesis. Henceforth, the term allergy will be used to designate these types of immune responses. The clinical definition of allergic disease includes the presence of allergic symptoms upon encounter with an allergen, such as rhinitis, conjunctivitis, itching or gastrointestinal symptoms, along with atopy, which comprises the heritable propensity to produce IgE-antibodies towards innocuous antigens.³ This represents the process of sensitisation, and may be examined by measuring allergen-specific IgE-antibodies in the circulation, or indirectly by surveying mast cell degranulation upon allergen challenge using skin prick tests (SPT). While a family history of atopy predisposes for development of allergic disease, and a number of genetic variants in genes such as human leukocyte antigen molecules and the T helper (Th) 2 cytokine locus have been associated to food allergies, asthma, AD and eczema⁷⁻¹⁴, the identified risk variants still only explain a small part of heritability in allergies. Inspecting the underlying causes of allergic diseases, environmental factors may contribute to the increased prevalence of allergies throughout the past century.¹⁵ It is unlikely that genetic alterations in the human genome as a result of evolutionary adaption would have taken place in such a short time span. Therefore, explanations are more likely to be found in structural changes that have taken place throughout the last 100 years in our societies, including behavioural, environmental and socioeconomic factors. This is further complicated by the more recent realisation that allergic disorders are highly heterogeneous.¹⁶ This diversity may be described by so-called endotypes, which represent groups of diseases with similar symptomatic characteristics, but with different aetiological causes. The characterisation of specific features of endotypes in both allergic symptoms and sensitisation are, however, just beginning to be uncovered.¹⁷

The development of allergic disorders often begins already in childhood, with AD and food allergies, followed by symptoms of wheezing, hay fever and ultimately asthma development during adolescence and early adulthood.¹⁸ This phenomenon has become known as the allergic march. The presence of one of the allergic manifestations does, however, not inevitably lead to the development of other allergic symptoms, neither do all children develop all of these diseases in succession. More recently, alternative models have been elaborated, suggesting that rather than being a progressive march, the development of AD may predispose for multimorbidity with later development of one or several other allergen-specific immune responses, with different

trajectories depending on the endotypes of the allergic manifestations.^{16,19} Both models provide an explanatory framework for different patterns of allergy development throughout childhood, which either way supports the rationale of intervening early in life, even *in utero*, to prevent development of allergies.²⁰

As both genetic and environmental factors interact in the development of this group of highly heterogeneous multifactorial diseases, multiple hypotheses describing the environmental impact on the recent increase in prevalence of allergic diseases have emerged. Already in 1976, the first observations of coinciding lesser exposure to bacteria, viruses and helminths in urban areas with higher prevalence of allergic diseases were reported.²¹ In the late 1980's, Strachan proposed what has become known as the hygiene hypothesis, suggesting that atopic diseases such as eczema and hay fever were possibly caused by smaller family sizes, as having older siblings seemed protective, along with cleaner living environments.²² Around the same time, Barker suggested that pre- and perinatal exposures during pregnancy and early life have an impact on future development of non-communicable diseases in adulthood.²³ This idea evolved to the developmental origins of health and disease hypothesis, explaining the phenomenon by mismatches between the programming of the immune system *in utero* and the actual outside living environment postpartum.²⁴ Later theories, such as the old friends hypothesis, puts forward how the progressive loss of previous exposures to microbial and parasitical communities throughout evolution may cause downregulated immunomodulatory responses, a feature common for immune and inflammatory diseases which are more prevalent in urban than rural settings.²⁵ Similar ideas of diminished immunoregulation due to loss of symbiotic relationships between our natural living environments and our commensal microbiota were suggested in the biodiversity hypothesis.²⁶ Altogether, these hypotheses discuss several plausible causes of the allergy epidemic, essentially revolving around one main outcome of modern life – microbial dysbiosis.²⁷

Over the past 70–80 years, major changes to our lifestyles have taken place that have altered the exposures we are evolutionarily equipped for.¹⁵ Events such as the introduction of refrigeration of foods, general vaccination programmes and antibiotic use were accompanied by the possibility of family planning and caesarean section. The centralisation of living in urban compared to rural areas has led to less contact with farm animals and natural environments, while simultaneously increasing our exposures to pollution. We spend more time indoors, use multitudes of cleaning agents for household and personal hygiene, and spend less time exercising. The intake of processed foods, saturated fats and carbohydrates has increased, while the intake of fruits, vegetables, fibres and unsaturated fats has decreased remarkably. Although the single contribution of each of these alterations is seemingly small, they have collectively reduced the diversity of microbial and environmental exposures – a deprivation with immense effects on our commensal microbial communities.²⁷ As decreased microbial diversity in the commensal microbiome may impair innate immune responses and in turn causes adaptive immune responses to become reactive instead of promoting tolerance, it has fundamental implications for development of appropriate immune responses.²⁸ This becomes particularly significant in the context of perinatal development, when maternal exposures of environmental and dietary origin direct and influence the evolvement of immunity in the foetus and new-born child.^{29,30}

The impact of the allergy epidemic on society and individual beings is immense, and early intervention is key to prevent the allergic march from progressing. Great efforts have been made to disentangle the contribution of genetic susceptibility and environmental factors in this group of highly heterogenous multifactorial diseases, but a lot still remains to be investigated.

Immunity and inflammation – endogenous protection in a world full of dangers

As in other jawed vertebrates, the immune system in humans consists of two tightly interrelated parts known as the innate and the adaptive immune systems.^{31,32} Each of these parts consist of functionally distinct cells that provide immune responses of varying longevity and specificity, whose main aim is to protect us from both intrinsic and extrinsic dangers, while at the same time providing support for beneficial microbes.³³ Hence, the immune system is challenged with the task of distinguishing between pathogenic microbes, cancer cells and toxins, which should be eradicated by inflammation, and symbiotic commensal microbes and other harmless exposures towards which tolerance should be induced. Upon recognising infection, infestation or danger, the immune system will respond by creating a local inflammatory reaction at the site of recognition, as a means to dispose of the danger at hand and eventually restore homeostasis in the tissue.³⁴

The innate immune system consists of cells such as mast cells, granulocytes (eosinophils, basophils and neutrophils) as well as dendritic cells (DCs) and macrophages (MØs), commonly referred to as antigen presenting cells (APCs).³¹ Furthermore, epithelia at our mucosal surfaces compose both physical and chemical barriers against intrusion. Innate immune responses are induced upon recognition of intruders through binding of *e.g.* microbe-associated molecular patterns and endogenous danger-associated molecular patterns to pattern recognition receptors such as Toll-like receptors (TLRs), NOD-like receptors and C-type lectin receptors, expressed both intracellularly and on the cell surface of different immune cells.³⁵ This ultimately leads to the production of inflammatory cytokines such as interleukin (IL)-12p70 by DCs and MØs³⁶, IL-1 β and IL-18 by the inflammasome, and stimulation of IL-6, interferon (IFN)- γ , and tumour necrosis factor (TNF)-production by cells such as T cells and MØs. These cytokines are instrumental in maintenance of inflammation, by promoting *e.g.* cell-mediated killing, phagocytosis and T cell activation. However, immunomodulating cytokines such as IL-10 and TGF- β are also produced, contributing to resolution of inflammation by inhibiting pro-inflammatory T cell-responses and inducing apoptosis.

Innate lymphoid cells (ILCs) are also important innate immune cells, which despite their lack of rearranged antigen-specific receptors provide a first line defence in response to dangers and stress in mucosal linings.³⁷ Their secretion of cytokines upon activation by epithelial-derived signals influences the nature of adaptive immune responses. The adaptive immune system consists of T and B lymphocytes, providing specific cellular as well as humoral immune responses against a wide variety of possible intruders.³⁴ Upon antigen uptake, the APCs will process the antigen, and migrate to draining lymph nodes, where they present the antigen on major histocompatibility complex (MHC) molecules on their surface to naïve T cells possessing a T cell receptor (TCR) with the same specificity for the antigen. This process represents a type of handover from the innate immune system to the adaptive immune system, allowing for development of specific immune responses with possibility to create immune memory. The recognition of the MHC-presented antigen on the APC by the TCR on the T cell is commonly referred to as signal 1.³¹ However, a second co-stimulatory signal, consisting of molecules such as the APC-T cell interacting B7-CD28 complex is indispensable for activation of the naïve T cell. Notwithstanding, a third signal provided by cytokines produced in the microenvironment by *e.g.* the APCs, epithelial and endothelial cells, is also important for appropriate differentiation

of the naïve Th cells into distinct Th cell subsets with distinct specificities and functions.³⁸ While a majority of these effector cells become apoptotic upon clearance of the antigen, a small proportion may develop into memory T cells, with the ability to mount faster and stronger specific immune responses upon repeated antigen encounter.³⁹

One of several roles for the Th cells is, as the name implies, to provide help to naïve B cells and promote the production of antigen-specific antibodies by these cells.⁴⁰ The B cells may in turn develop into memory B cells, which are present at mucosal sites all over the body as well as in the blood, from where they may re-circulate to secondary lymphoid tissues, such as lymph nodes or the bone marrow. They may also become long-lived plasma cells, which mainly reside in the bone marrow and at mucosal sites.⁴¹ This means that, upon re-exposure to the same antigen, memory Th cells recognise the antigen more readily and redirect the signal to several cell types to induce a faster and stronger immune response. The forwarded signal can drive development of memory B cells into plasma cells, which may secrete antigen-specific antibodies promoting the neutralisation of the antigen by *e.g.* phagocytosis (IgG-mediated opsonisation)⁴², immune exclusion (IgA at mucosal surfaces)⁴³ or release of pro-inflammatory mediators (IgE-mediated mast cell or basophil degranulation).⁶ Resolution is established by decreased recruitment of inflammatory immune cells, induced apoptosis and clearance of dead cells, along with presence of regulatory T cells (Tregs) and immunomodulatory mediators.⁴⁴ In this way the innate immune response, while exhibiting limited specificity of antigen recognition and lacking long-term memory, initiates and paves the way for both specific and long-term cellular and humoral immune responses, ultimately leading to restoration of homeostasis in the inflamed tissue.

Inflammatory responses induced by cells and tissues of the innate and adaptive immune systems are necessary for the protection of the organism from both intrinsic and extrinsic dangers. The process not only encompasses the induction of inflammation, but also includes the restoring of homeostasis in the affected tissue. Overriding pro-inflammatory responses and/or poor pro-resolving responses may pave the way for diseases, such as allergic disease.

T helper cells – friends and foes in health and disease

The differentiation of Th cells is a highly regulated process, which may stably induce polarisation of naïve Th cells into one of several effector Th cell subsets.³⁸ This process takes place in secondary lymphoid tissues such as lymph nodes, and upon differentiation and proliferation the effector Th cells enter the circulation and migrate into tissues in the quest for their cognate antigen.³¹ However, the differentiation into any of the subsets is largely dynamic, as even differentiated Th cells may adapt to changing environmental stimuli and virtually become other Th cell subsets.⁴⁵ In the adaptive immune response, Th cells confer specific protection towards a variety of external threats, with distinct subsets being adapted to handle different pathogens (Figure 1).

In the healthy state, Th1 cells provide protection against intracellular pathogens, *i.e.* viruses and intracellular bacteria. However, dysregulated immune responses by Th1 cells and their signature cytokines have been implicated in the development of several autoimmune diseases, such as multiple sclerosis and type 1 diabetes.⁴⁶ Similarly, the Th17 subset, which normally handles intrusion of extracellular bacteria and fungi⁴⁵, is also involved in the development of autoimmune disorders as well as psoriasis.⁴⁶ Th17 cells are key players in upholding tolerance in the gut, and provide a low-grade inflammation in the mucosae and epithelia of the body in the steady state, in the protection against possible intruders.⁴⁷ Th2 cells on the other hand, normally combat considerably larger multicellular pathogens, namely parasitic helminths.⁴⁵ What distinguishes Th2 cells from the other Th cell subsets is the immunologic responses induced by the recognition of a possible parasite.⁴⁸ Whereas Th1-responses target intracellular pathogens for degradation by the phagocytic abilities of MØs, and Th17 cell-responses induce inflammation by recruiting neutrophils to eradicate extracellular pathogens, Th2 cells recruit other immune cells with abilities to take on the sheer size of the parasite.⁴⁵ Hence, upon recognition of a parasite, Th2 cells secrete IL-4, IL-5 and IL-13, cytokines which will enhance barrier function through *e.g.* elevated mucus-production, and promote killing and expulsion of the parasite by eosinophil activation and IgE-dependent mast cell degranulation.⁴⁸ However, in the present-day context, many human beings do not encounter large amounts of microbial or helminthic exposures, and even less suffer from parasitic infestations. This has redirected these immune responses towards innocuous antigens, causing the symptoms that we have come to recognise as allergic reactions.

As a counterbalancing subset among the different inflammation-promoting Th cells stand the Tregs. These cells are indispensable for immune regulation of inflammation and the return to tissue homeostasis, and are commonly divided into the natural and induced Tregs.⁴⁹ While natural Tregs are induced in the thymus by self-antigens, and thereby mainly sustain self-tolerance, induced Tregs are induced upon exposure to antigens and transforming growth factor (TGF)- β in peripheral lymphoid tissues, providing immune dampening responses to non-self antigens of *e.g.* microbial or dietary origin.

Naïve Th cells have gone through maturation from lymphoid progenitors in the foetal liver and later the bone marrow, followed by several steps of thymic maturation prior to their migration to lymph nodes as single positive CD4 cells.⁵⁰ Professional APCs such as DCs encounter antigens in tissues, and migrate to draining lymph nodes to present the processed antigen on MHC II molecules to naïve Th cells (Figure 1). The directional fate of the polarisation of Th cells is dependent on both the type of antigen presented by the APCs, but also how strong the provided signal is, in terms of affinity and temporal length of interaction.^{51,52} Particularly interesting for

the development of allergic disease is that high-dose antigenic exposure favours Th1 cell differentiation, while low-dose antigenic stimulation favours the differentiation of Th2 cells. Additionally, the presence of different types secreted immune mediators, mainly cytokines, by the cells in the microenvironment surrounding the naïve Th cell also play an important role in determining Th cell fate.⁵⁰

The differentiation into different Th cell subsets requires presence of particular cytokines, along with molecules known as signal transducer and activation of transcription (STAT), for their further polarisation into their respective effector Th cells.⁵³ Furthermore, lineage-specific master transcription factors define the outcome of the maturation into each Th cell subset (Figure 1). The differentiation into Th1 cells is driven mainly by the cytokine IL-12p70, which through the transduction of the signal via STAT4 and autocrine IFN- γ mediated STAT1 translocation, activates T-bet – the master regulator of Th1 cells.³⁸ The effector functions of Th1 cells, mediated mainly by IFN- γ , include increasing phagocytic capacities of MOs and increasing the expression

Figure 1. T helper cell differentiation at a glance in health (middle circle) and disease (outer circle).

of MCH I, which is expressed on most nucleated cell types.⁴⁵ Taken together, these are the measures the Th1 cells initiate to combat intracellular viruses and bacteria.

Epithelial-derived cytokines such as IL-25, thymic stromal lymphopoietin (TSLP) and IL-33 may promote the differentiation of Th2 cells⁵⁴, by creating pro-inflammatory DCs that preferentially drive Th2 differentiation⁵⁵, or by the induction of group 2 ILCs (ILC2), which promote the differentiation of Th2 cells through the actions of IL-4.³⁷ Th2 cells become destined to their lineage by the effects of IL-4 on STAT6 expression, leading to the induction of the Th2-distinguishing GATA-3 transcription factor. The main functions of the Th2 cells include the recruitment of eosinophils and Th2 cells to the sites of infestation, along with increased mucus-production, thereby aiding the killing and expulsion of the parasite from the body.⁴⁸ This is achieved by the actions of the Th2-signature cytokines IL-4, IL-5 and IL-13.⁶

The differentiation into Th17 cells depends on the expression of IL-6 and TGF- β , which in turn activate STAT3 and ultimately the master regulator ROR- γ t.⁵⁶ These Th17 cells are considered non-pathogenic with a more Treg-like phenotype, while concomitant IL-23 exposure to differentiated Th17 cells promotes Th1-like pathogenic cells⁴⁷ and enhanced IL-17-production.⁵⁷ IL-17A, IL17F and IL-22 are the main cytokines that exercise the function of Th17 cells, both in maintenance of barrier integrity at mucosal tissues as well as in inflammatory responses towards external insults.⁵⁸

Tregs are mainly poised towards their identity by the cytokines IL-2 and TGF- β , which induce STAT5 expression and subsequent activation of the master transcription factor FoxP3.⁴⁵ The effector cytokines of Tregs mainly include IL-10 and TGF- β , which act in an immunomodulatory fashion on inflammation.⁵⁹ Upon differentiation and proliferation in the lymph nodes, the polarised and expanded Th cell populations leave the lymph nodes, and begin to patrol in the peripheral circulation in the search for their matching antigen. Thereafter, they may be attracted to the site of an infection, where the Th cells exercise their effector functions. A proportion of the effector cells will develop into different populations of memory Th cells. These cells may either be tissue-resident in *e.g.* the mucosae, or re-circulating between the blood and secondary lymphoid organs or non-lymphoid tissues, thereby providing early ignition for induction of inflammation upon re-encounter with an antigen.³⁹

During Th cell differentiation, amplification of cytokine-production by the differentiating cell creates positive feedback loops to promote development of the particular Th cell subset.⁵³ For instance, the autocrine production of IFN- γ and IL-4 by differentiating Th1 and Th2 cells, respectively, support and stabilise the maturation into their specific lineages. This is further reinforced by the antagonistic effects of the transcription of one master transcription factor inhibiting the transcription of another master regulator, exemplified by the inhibitory effects of T-bet on GATA-3 expression, preventing Th1 cells from becoming Th2 cells. These phenomena are largely regulated by epigenetic mechanisms, including both histone modifications, but also DNA methylation.⁶⁰ These modifications affect the transcriptional activity of the genome and are partly responsible for the dynamic adaptability of Th cells to changes in environmental cues, creating a in reality considerably more heterogenous population of Th cells consisting of different phenotypical hybrids.⁴⁵ Likewise, memory Th cell populations also show some degree of plasticity and wide heterogeneity.³⁹

Th cells make up a largely heterogenous group of antigen-specific adaptive immune cells, which are instrumental in the combating of viruses, bacteria and parasites. Exposures in the microenvironment provide cues for developmental polarisation of naïve Th cells in the lymph nodes, initiating the development of lineage-specific traits, which is accomplished by the encounter of cognate antigen in peripheral sites. Dysregulated differentiation of Th cells, due to alterations in antigen presentation and environmental stimuli, is believed to bring about the imbalances seen in the development of several diseases, including autoimmune and allergic diseases.

Allergy development – dysregulated inflammatory responses to innocuous antigens

In an allergic individual, immune responses that in evolutionary history were aimed at the protection against venoms, environmental toxins and the eradication of parasitic infections, are initiated towards innocuous antigens causing the development of allergic symptoms common to many allergic diseases.⁴⁸ A common feature of allergic immune responses is the imbalance in the adaptive Th cell compartment. There is an apparent skewing of immunity in atopic individuals towards Th2-responses, while Th1 and Treg cells are altered both in frequencies and function.^{61,62} The initiation and maintenance of allergic inflammation depends on the recruitment of both adaptive and innate immune cells to the site of the allergic reaction.⁵⁹

For allergic immune responses to develop, the immune system requires repeated exposure to an allergen, at least twice. The local microenvironment at the site of allergen uptake dictates whether the immune response towards the allergen will be tolerant or inflammatory in nature.⁶ In allergic individuals, the barrier function of epithelia in the skin and the mucosal linings of the airways and the gut are often compromised, leading to higher exposures to microbes, pollutants and allergens.⁶³ This may induce the production of pro-inflammatory mediators such as IL-25, TSLP and IL-33 by the epithelium.⁶⁴ These mediators may in turn promote the activation of ILC2s, which respond to stress and danger signals of the epithelium³⁷, thereby shaping initial immune responses at the mucosal barrier by the expression of cytokines such as, in the case of allergic immune responses, IL-4, IL-5, IL-13.^{65,66}

Upon the primary encounter with an allergen at any of the human body's epithelial linings, tissue resident DCs take up the allergen, process it and transport it via the afferent lymphatic vessels to the closest draining lymph node (Figure 2).⁶ In an allergy-promoting setting, the allergen is presented on MHC II molecules to naïve Th cells, initiating TCR signalling and preferential differentiation into Th2 cells, which when activated produce several cytokines including IL-4, IL-5, IL-9 and IL-13.⁶⁷ The isotype-switch of B cells to produce allergen-specific IgE-antibodies is induced by the actions of IL-4 upon Th cell help and antigen recognition. Subsequently, the secreted allergen-specific IgE-antibodies may bind to the high-affinity receptor FcεRI on mast cells, eosinophils and basophils, thereby completing the process of sensitisation. Upon repeated encounter with the same antigen, the sensitised cells may become activated, as the allergen cross-links several FcεRIs on the cell surface, ultimately leading to the degranulation and secretion of pre-formed and *de novo* produced biogenic amines and lipid mediators into the surrounding tissues.⁶ As mast cells are often located close to both nerve endings and blood vessels⁶⁸, this enables rapid induction of immediate hypersensitivity responses by releasing histamine, prostaglandins and leukotrienes into the circulation as well as locally in mucosal tissues or the skin. While the prostaglandins and leukotrienes elicit vasodilation, increase membrane permeability and recruit eosinophils to the allergic reaction site, histamine is responsible for symptoms such as gastrointestinal hyperactivity, itch, bronchoconstriction and the typical wheal and flare responses of the skin, which are used in SPTs to diagnose sensitisation.^{48,68} These types of reactions develop within minutes to hours after exposure, and are termed as early phase hypersensitivity reactions.⁶ The worst case scenario of systemic early phase reactivity is known as anaphylaxis. Late phase hypersensitivity reactions develop within the first 24 hours after repeated allergen exposure and originate from newly produced growth factors, cytokines and chemokines of the mast cells. These mediators contribute to the local accumulation of eosinophils in the affected tissues, and

Figure 2. A schematic overview of the process of sensitisation, and early and late phase effects upon allergen re-exposure.

attract allergen-specific Th2 cells to the site of the allergic reaction. Consequently, this leads to perpetuation of allergic inflammation, as eosinophil recruitment and activation at the site of the allergic reaction is upheld by IL-5⁶⁹, while IL-13 stimulates the production of mucus at mucosal sites, and IL-4 may continue to promote the production of IgE-antibodies.⁶ As immunoregulatory T cell-responses are often dysregulated in allergic individuals⁷⁰⁻⁷³, there is a substantial risk that the allergic inflammation becomes a chronic state, especially if the exposure is continuous. As with any type of inflammatory response, resolution of allergic inflammation usually takes place by means of the actions of Tregs (IL-10, TGF- β)⁷⁴, M ϕ s (phagocytosis and promotion of tissue regeneration)⁷⁵ and fatty acid mediators known as specialised pro-resolving mediators (SPM).⁷⁶

Allergic reactions develop upon repeated exposures to allergen in environments favouring inflammatory responses. Symptoms of allergic disease are caused by IgE-dependent activation of mast cells and the actions of Th2-derived cytokines on epithelia, B cells and eosinophils. In the characteristic imbalance of the Th cell immune responses towards Th2-immunity in allergic individuals, which is further amplified by diminished Th1- and Treg-responses, the development of chronic inflammation is impending. The reversal of sensitisation is therefore crucial in the quest for restoration of tolerant immune responses in improving and preventing allergic disease.

Epigenetic regulation – transcriptional modifications modulating interplay between genome and environment

The coining of the term epigenotype by Waddington in the early 1940's marked the beginnings of the work in the field of epigenetics.⁷⁷ Over the decades it has evolved to describe epigenetics as the bridging mechanisms regulating how the genome may adapt to changing environmental conditions, while at the same time serving as explanatory models for the underlying causes of non-communicable multifactorial diseases. Epigenetic modifications constitute several layers of transcriptional regulation, including DNA methylation, histone modifications and microRNAs, which do not affect the genomic nucleotide sequence itself (Figure 3).⁶⁰ DNA methylation is involved in processes such as silencing of repetitive elements in the genome, X-inactivation and allele-specific gene imprinting throughout development.⁷⁸

DNA methylation is considered the most stable epigenetic modification, as the patterns are heritable throughout the cell division process, thereby giving rise to epigenetic memory.⁷⁹ However, at the same time, DNA methylation is highly dynamic in its interaction with environmental exposures, as corroborated by studies showing that patterns change at different rates⁸⁰ throughout life⁸¹, and that individual differences in DNA methylation patterns are common.⁸²

The process of adding a methyl group ($-CH_3$) to the fifth carbon in cytosine residues is covered by a family of proteins called DNA methyltransferases, which are able to preserve and create new DNA methylation patterns.⁸³ On the other hand, the active process of demethylation is handled by other enzymes, known as ten eleven translocation enzymes, which form an intermediate between unmethylated cytosines and methylcytosine, namely hydroxymethylcytosine. By binding to methylated cytosines in the genome, methyl Cytosine-phosphate-Guanine (CpG)-binding domain proteins recruit further protein complexes, which in turn regulate the transcriptional fate of the site of interest. The sites at which this process takes place are mainly CpG sites, hence sites in the genome where a cytosine is followed by a guanine joined by a phosphate group.⁸⁴ These sites tend to be clustered in CG rich regions known as CpG islands, often located in close proximity to promoter regions, and exist mainly in a demethylated state.⁸⁴ About 10% of all CpGs in mammalian genomes are localised to CpG islands, and between 60% to 80% of all CpG sites are methylated at any given time point.⁸⁵ In promoter regions, DNA methylation mainly leads to transcriptional silencing, whereas in other genomic regions such as intergenic regions the opposite is true⁸⁶, emphasising the context dependent feature of epigenetic regulation.⁸⁷ DNA methylation may act directly to inhibit transcription, by the physical blocking of transcription factor binding to their binding sites.⁸⁸ Additionally, the recruitment of other proteins to the methylated CpG sites, either by transcription factors recognising methylated cytosines or in an indirect fashion by methyl CpG-binding domain proteins, has also been described as possible mechanisms for altering transcriptional accessibility at DNA methylation sites.⁷⁹

During the course of human development, from prenatal life throughout childhood, adolescence, adulthood and aging, DNA methylation patterns continuously adapt to transforming environmental exposures⁸⁰⁻⁸², and play an important role in different stages of maturation including that of the immune system. *In utero*, the growing foetus is programmed in preparation for the exposures of the *ex utero* environment, which the pregnant mother is exposed to.⁸⁹ The transplacental transfer of gases, nutrients and metabolites promotes the growth

of the child, while at the same time epigenetically imprinting the foetal genome to phenotypically match the challenges to be expected *post partum*. However, if there is a discrepancy between the expected outside environment and the imprinted epigenome, the resulting mismatch may lead to dysregulation of multiple organ systems, including that of the immune system²⁰, leading to the development of *e.g.* allergic disease.⁸³ Apart from exposures in

Figure 3. A schematic overview of epigenetic regulation, from chromosomes via nucleosomes and histones to DNA methylation.

the microenvironment of the lymph nodes, such as cytokine secretion, and antigen-dependent co-stimulated activation of TCR signalling, DNA methylation patterns are crucial players in the differentiation from naïve Th cells into specific Th cell subsets.⁷⁹ In the polarisation process separating distinct Th cell fates, cytokines of the microenvironment induce the binding of lineage-specific transcription factors to induce the expression of cytokines which amplify the differentiation and promote the effector functions of the Th subset at hand.⁴⁵ Furthermore, the promotion of the designated lineage is favoured by concurrent repression of other Th cell lineages, which is regulated at the epigenetic level. As an example, the expression of the Th2 cell characteristic transcription factor GATA-3 is induced by the Th2-associated cytokine IL-4, and while it positively supports the transcription of the IL-4, IL-5 and IL-13 locus, it also inhibits the expression of IFN- γ locus.⁷⁹ This is achieved by alterations in the accessibility of chromatin in enhancer and promoter regions of the cytokine gene loci, with increased permissive histone marks at Th2-associated loci and repressive counterparts at Th1-associated loci. Moreover, active demethylation takes place at cytokine gene loci responsible for the respective effector functions of the cells, emphasising the role of DNA methylation processes in development and differentiation of Th cells.^{90,91} Effector cells may also develop into memory cells, which is regulated by epigenetic marks on both the DNA methylation and histone modification levels, thereby providing the epigenetic basis of rapidly responding antigen-specific immune responses.⁹⁰

As epigenetic modifications regulate gene expression based on influence by environmental exposures, and DNA methylation plays a crucial role in the differentiation of Th cells, the potential involvement of dysregulated epigenetic regulation in the development of allergic disease is conceivable. Indeed, differential DNA methylation patterns have been revealed in allergic diseases in several studies performed in children and adolescents, most notably in the development of wheezing,⁹² asthma⁹³⁻⁹⁶, food allergies⁹⁷⁻⁹⁹, eczema¹⁰⁰ and sensitisation.¹⁰¹⁻¹⁰³ Most of these studies have surveyed whole blood or mononuclear cells from peripheral or cord blood samples, whereas few have investigated CD4+ T cells more specifically. The exception is presented by the works of Martino *et al.* on food allergy in children. Comparisons of the DNA methylomes at birth and age 1 year in food-allergic children demonstrated 92 differentially methylated probes (DMPs) that were enriched within the MAPK-signalling pathway.⁹⁸ As CD4+ cells in food-allergic children were poised towards hypomethylation in differentially methylated sites of activated T cells⁹⁷, this altogether could explain the suboptimal maturation and function of these cells in food allergies. Evidence also suggests that exposures of previous generations may affect phenotypic outcomes of allergy in the developing foetus or child, even if the exposure is no longer present. An example of this is the effect of parental/grandparental smoking on the development of asthma on the children/grandchildren.¹⁰⁴ The heritable feature of epigenetic modifications becomes particularly interesting in the interaction between mother and child in modulating allergy development. For instance, hypermethylation in the *SMAD3* gene, which is an important mediator of TGF- β signal transduction, was observed in cord blood mononuclear cells (CBMC) of asthmatic children, but only if their mothers were also asthmatic.⁹³

Epigenetic modifications provide transcriptional control of genomic expression in adapting to environmental changes. Dysregulation in processes of foetal and neonatal immune maturation, including differentiation of Th cells, contributes to the development of allergic disease, implying that epigenetic mechanisms may be involved. Evidence of altered DNA methylation patterns in allergic disease corroborate this notion, but how these modifications may be regulated in allergic children upon immunomodulating intervention is not well-established. In Papers II and III, the epigenetic effects of perinatal intervention with probiotics with or without combined ω -3 fatty acid treatment will be studied.

Chemokines – key orchestrators in immune cell recruitment

The migration of immune cells throughout the human body is a highly temporally and spatially regulated process, and the main mediators coordinating this process are known as chemokines.¹⁰⁵ Chemokines are a family of low molecular weight chemotactic cytokines which, depending on the position of their N-terminal cysteines, are further subdivided into the CC, CXC, CX₃C and XC families of chemokines.³⁶ The chemokines bind to receptors that are termed either conventional or atypical, distinguished by the downstream activation of several common immune-related signal transduction pathways in the former case.¹⁰⁵ The type of chemokine ligand they bind is mirrored in the receptor name. Most conventional chemokine receptors are G-protein coupled receptors. Chemokines are produced by different cell types, such as MØs and DCs as well as endothelial and epithelial cells upon induction by cytokines. Thereafter, the chemokines are secreted into the interstitial space and bind to glycosaminoglycans in the extracellular matrix.¹⁰⁵ This process is indispensable for creating gradients of the chemokines, which not only affects the half-life and function of the chemokine itself, but more importantly provides a direction in physical space in which the cells binding the chemokines may migrate.

The specific binding of the extracellular matrix-bound or soluble chemokines to G-protein coupled receptors on the cell surface of immune cells constitutes the starting point of chemotaxis.¹⁰⁶ As chemokines exhibit instrumental functions in processes such as leukocyte extravasation, they indirectly orchestrate the induction and maintenance of inflammatory responses, as well as the restoration of homeostasis in several tissues. The case of allergic inflammation is no exception in this regard. The recruitment of immune cells such as mast cells, Th2 cells and eosinophils to the site of an allergic reaction constitutes a prerequisite for the induction and maintenance of allergic inflammation.⁵⁹ The presence of certain chemokines represents the immune responses which initiated the production in the first place. Hence, there are chemokines which are highly associated to different immune responses.

For instance, the CXC chemokines CXCL10 and CXCL11 are tightly regulated by the Th1-response, due to their induction by the Th1-associated cytokine IFN- γ (Figure 4).¹⁰⁷ These chemokines bind to the CXC receptor 3 (CXCR3), which is expressed on cells of both the innate and adaptive immune responses, namely on CD4+ Th1 cells and CD8+ cytotoxic T cells, but also natural killer, natural killer T cells and plasmacytoid DCs. Thus, CXCL10 and CXCL11 contribute to the potentiation of Th1-associated inflammation in general, and in the combating of viral infections in particular. However, in a disease setting, they are also known to contribute to inflammatory Th1-responses in autoimmune disorders¹⁰⁸ as well as counterbalance Th2-responses in allergy development.⁵⁹ CXCL10, or inducible protein 10 as it is also called, is primarily produced by monocytes, fibroblasts, endothelial cells as well as thymic epithelial cells.¹⁰⁹ On the contrary, CXCL11 or IFN-inducible T cell alpha chemoattractant, is expressed by cells of the thymus, spleen and lung, as well as by peripheral blood leukocytes.¹¹⁰⁻¹¹² In several previous studies from our research group, we have shown associations of the peripheral levels of CXCL10 and CXCL11 with allergic outcomes. For instance, elevated levels of CXCL11 and a tendency for the same trajectory for CXCL10 was observed in children developing recurrent wheezing.¹¹³ This is in agreement with findings from children who wheezed at age three, and later developed allergic asthma at age six, where the plasma levels of CXCL10 were augmented.¹¹⁴ A large number of other studies corroborate these results in asthmatic individuals, showing increased concentrations of either or both of the two Th1-associated chemokines in tissues such as nasal

Cytokine production upon T cell activation

Chemokine production by various cell types

Chemotactic effects of chemokines upon binding to GPCR

Figure 4. A summarising outline of the production and effects of allergy-related cytokines and chemokines.

lavage fluid^{115,116}, serum¹¹⁷, bronchial epithelial cells^{118,119}, sputum¹¹⁷ and bronchoalveolar lavage.¹²⁰ In relation to sensitisation, however, the Th1-associated chemokines show a more complicated mechanism of action. In one of our previous studies, the levels of CXCL11 at birth and 2 years of age were actually decreased in children later becoming sensitised compared to in healthy children, and high CXCL11 levels were inversely associated with development of sensitisation.¹¹³

Likewise, a number of chemokines are highly connected to the Th2-response, amongst them CCL17 (thymus and activation-related chemokine) and CCL22 (macrophage-derived chemokine) (Figure 4).¹²¹ Both CCL17 and CCL22 bind with high affinity to the CC family chemokine receptor CCR4, which is primarily expressed on Tregs and Th2 cells as well as skin-homing T cells. While CCL17 mainly induces chemotaxis in T cells, CCL22 may additionally support migration of monocytes, DCs, and natural killer cells.^{121,122} Thymic as well as myeloid cells are predominant producers of CCL17 and CCL22¹²², and whereas CCL17 may be induced by IL-4 in T cells¹²³, CCL22 may be induced by TNF, IL-1 β and lipopolysaccharide in DCs, and IL-4 and IL-13 in cells of the myeloid lineage.¹²² The implication of these Th2-associated chemokines in allergic diseases has been widely documented in several studies, amongst them our own. In sensitised children with allergic symptoms^{124,125}, and in children developing wheeze¹¹³ and asthma¹²⁵, the levels of CCL17 and CCL22 were elevated. The levels of both Th2-associated chemokines at birth also predicted the development of allergic disease, as increased concentrations of CCL17 and CCL22 in cord blood preceded the development of allergic symptoms and sensitisation, respectively, throughout the first 6 years of life.^{124,125} This was more recently corroborated by Björkander and co-workers, displaying that elevated plasma CCL17 and CCL22 levels at 6 months and 1 year of age preceded the development of allergic disease throughout the first decade of life.¹²⁶ Furthermore, low cord blood CCL22 levels at birth were inversely associated with dust mite sensitisation as well as asthma development in childhood.¹²⁷

Another chemokine of interest in development of allergic diseases is the CC chemokine CCL18, or pulmonary and activation-related chemokine (Figure 4). CCL18 is constitutively expressed in *e.g.* the lung and circulation by tissue-resident APCs, and to a lesser extent in thymus, bone marrow and lymph nodes.¹²⁸ This suggests that it has regulatory capacities in times of homeostasis. However, during inflammatory responses it may be further upregulated, which is also the case upon allergen exposure. The expression of this chemokine is induced by both the allergy promoting Th2-response and the immune dampening Treg-responses, as it may be induced by IL-4 and IL-13, along with the immunomodulating cytokine IL-10.¹²⁹ In a similar fashion as the induction of Th cell-related cytokines, the cytokines needed for induction of Th cell-associated chemokines will inhibit the production of other chemokines. For instance, the expression of the two purely Th2-associated chemokines CCL17 and CCL22 is inhibited by the effects of IL-10^{130,131}, while expression of CCL18 is inhibited by the CXCL10 and CXCL11-inducing cytokine IFN- γ .¹²⁹ Until rather recently, the receptor for CCL18 eluded science, but in 2013 Islam *et al.* suggested that the CC receptor CCR8 was the cognate receptor for this chemokine in humans.¹³² However, CCR8 is probably not the only relevant receptor for CCL18, as naïve T cells migrate readily in response to CCL18, despite their lack of CCR8 expression.¹²⁸ In general, immune cells such as Th2, Tregs and skin-homing cutaneous lymphocyte antigen positive T cells express CCR8, displaying that CCL18 may hold roles in both driving inflammation as well as resolving it. As is the case for the previously described chemokines, CCL18 also has been implicated in allergy development, with augmented expression in allergic

diseases such as asthma^{133,134}, allergic rhinitis¹³⁵ and AD.^{136,137} We have previously shown that the circulating levels of CCL18 were elevated in children who develop recurrent wheeze and eczema in infancy.¹¹³ Interestingly, CCL18 possesses features other than purely chemotactic ones, including the possibility to polarise naïve T cells into Tregs producing both IL-10 and TGF- β .¹³⁸ In allergic disease, however, the regulatory capacities of CCL18 are altered, as its binding to immune cells is diminished^{138,139}, which is particularly intriguing as allergic individuals often show less induction of Tregs.⁶¹ Additionally, it stimulates the production of collagen in both skin and lung fibroblasts, providing a mechanism of involvement in allergic diseases such as asthma.¹⁴⁰

Chemokines recruit immune cells to the site of the allergic reaction, thereby initiating and maintaining allergic inflammation. Different immune responses induce the production of related chemokines, and the chemokines CCL17, CCL18, CCL22, CXCL10 and CXCL11 have all been implicated in the development of allergic disease. The role of these chemokines in childhood have been partly explored, but they have not been studied in a larger setting of an observational cohort studying allergy development from birth until young adulthood. These aspects are covered in the analyses performed in Paper I.

Perinatal immune maturation – the importance of maternal exposures during pregnancy and early life

The immune system of the mother, as well as her unborn child, is during pregnancy skewed towards Th2-responses¹⁴¹, along with high presence of Tregs¹⁴², in order to uphold tolerance towards the semi-allogeneic foetus. Normally, vaginal delivery promotes the enhancement of Th1-responses, through microbial exposures of the birth canal, thereby preparing the neonate for the outside world.¹⁴³ However, dysregulated immune responses caused by *e.g.* maternal allergic disease, potentially harmful environmental exposures during pregnancy or delivery by caesarean section may prevent the Th1-response from developing, hence allowing for the prenatal Th2-skewing to persist and promote development of allergic disease.

During foetal development, the foundations of immunity in the protection against internal and external threats are being laid. At the time of birth, the neonatal immune system is practically naïve and immature, and begins to mature in response to the extensive exposures of environmental stimuli.¹⁴³ As this takes place throughout the first decade or two of life, appropriate immune responses against a multitude of bacteria, viruses, fungi and parasites continuously evolve throughout the course of childhood and adolescence, and eventually create immunological memory, which may last even for a life-time.²⁹

The development of the human immune system begins in foetal life, already throughout the first trimester, with haematopoietic stem cell progenitors of both myeloid and lymphoid cells being present at four weeks after conception.¹⁴⁴ DCs are present in the foetal liver at about six weeks after conception, with foetal B cell precursors appearing shortly thereafter.¹⁴⁴ Foetal T cells develop into naïve Th cells upon migration from the foetal liver to the thymus at about eight weeks after conception, and may thereafter migrate to peripheral tissues in the end of the first trimester to develop into foetal effector T cells. The majority of the T cells are $\gamma\delta$ -T cells until the proportion shifts in favour of $\alpha\beta$ -T cells by the middle of the third trimester.³⁰ These cells mainly develop into Tregs, and as foetal DCs furthermore favour T cell-production of IL-4 and inhibit secretion of TNF¹⁴⁴, both foetal-maternal and self-tolerance in the womb may be maintained. Interestingly, memory cells have also been demonstrated in the foetal intestine, suggesting that maternal exposures have induced long-term responses *in utero*.¹⁴⁴

Although the major constituents of the adult immune system are in place early on in human development, there are major differences in terms of functionality and frequencies of these compartments. Foetal and neonatal immune responses resemble each other in both innate and adaptive immune responses. While the number of neutrophils, natural killer cells and monocytes increase after birth, neither frequencies nor function of these cells correspond to their adult counterparts.³⁰ Furthermore, the new-born complement system shows diminished function compared to mature immune responses of adults.²⁹ Postnatal adaptive immune responses are generally weak, due to the previously limited antigen exposure, but also caused by the lack of germinal centres in secondary lymphoid organs, which are crucial sites for the Th cell-induced production of antigen-specific antibodies by B cells.³⁰ Furthermore, antibodies produced in early postnatal life are limited in their affinity maturation, and the induction of memory B cells and plasma cells is restricted, thereby restraining neonatal humoral immune responses.²⁹ As mentioned above, Th cell-responses *in utero* tend to be tolerant with a prominent Th2-dominance in both the mother and her child at the end of pregnancy and early postnatal life.²⁹ Altogether, this renders new-born babies more susceptible to viral and bacterial infections, as

constituents of innate immune responses are diminished, while virtually naïve and separated T and B cell compartments hamper the induction of long-lived antigen-specific humoral immune responses.²⁹

The development of the foetal and neonatal immune systems in pregnancy and early postnatal life is mainly influenced by maternal exposures.^{15,33,145} Prenatally, the transplacental transmission of dietary nutrients and metabolites as well as antigens and antibodies supplies the foetus not only with building blocks for general growth, but also environmental cues for induction of foetal immune responses. Transplacental communication of IgG-antibodies, either as is or complexed with antigens of microbial or allogeneic origin, is enabled through transport by the neonatal Fc receptor¹⁴⁶, which by endocytotic processes transports and releases the contents into the foetal circulation in the placenta.¹⁴⁷ This begins early in the second trimester, enabling both the passage of passive immunity from the mother, while at the same time influencing immune maturation. This is further sustained postnatally throughout breastfeeding, by supply of nutrients through the maternal milk, along with immune cells, IgA-antibodies, cytokines and microbiota.^{30,148} In this way, the maternal and infant immune responses are aligned, by vertical transmission of both passive and active immunity and by sharing of components of the maternal immune system, which have major impacts on foetal immune maturation.

The presence of limited immune responses in the foetus has prompted the question of whether or not the *in utero* environment is sterile. Although some studies suggest that the placenta exhibits a distinct microbiome¹⁴⁹, and bacteria are present in meconium¹⁵⁰, solid evidence for colonisation of the foetus by placental microbes are still missing.¹⁵¹ During labour and delivery, however, the neonate is exposed to an extensive variety of microbiota, originating from the birth canal and maternal gut as well as the air and the surroundings of the environment in which the new-born takes its first breath.²⁹ The immense impact of this primary explosion of microbial exposures for immune maturation is evident from children born with caesarean section, who have a lower microbial diversity compared to children born vaginally¹⁵² with a higher risk of developing diseases such as allergies and type 1 diabetes.¹⁵³ The use of antibiotics during pregnancy has also been shown to alter microbial communities and immune maturation in the offspring.³⁰ The finetuned interplay between the maternal microbiome, her ingested diet and environmental exposures affects immune maturation in the foetus by the effects of primary and secondary metabolites.¹⁵⁴ Thus, the impact of maternal influence on foetal immune maturation is tremendous, and the composition of both the maternal commensal communities and her ingested diet is key in the process.^{15,155,156}

A large body of evidence points out the significance of perinatal exposures to a wide variety of microbes and dietary antigens for appropriate immune maturation. This begins already in the womb, and continues after birth throughout breastfeeding, giving particular importance to dietary and environmental maternal exposures on foetal and neonatal immunity. Perturbations in the diversity of antigenic exposures during this critical time-period may hamper development of innate and adaptive immune responses, and pave way for immune mediated diseases such as allergies, with possible long-term consequences on health.

Mucosal immunity – immunological barriers balancing host defence and tolerance induction

The mucosal linings of the gastrointestinal tract and airways make up the greatest surface area of exposure to the outside environment, constituting barriers which are continuously exposed to a large plethora of antigens.¹⁵⁷ An important task of the mucosal barriers consists of distinguishing between innocuous agents such as dietary antigens and commensal bacteria, towards which tolerance should be induced, and pathogenic bacteria, viruses and parasites, which should be eradicated by inflammatory responses. The uptake of nutrients from the diet has to be maintained while preventing microorganisms, whether pathogenic or commensal, from penetrating the mucosae. The process of actively suppressing the induction of immune responses to antigens derived from commensal bacteria or dietary origins which are encountered in *e.g.* the gut is known as oral tolerance.¹⁵⁸ This is achieved by means of both clonal deletion and anergy of T cells reacting to innocuous antigens, as well as by the induction of immunoregulatory responses mediated mainly by Treg cells. The distinction between the myriads of both beneficial and dangerous exposures is achieved by a refined process involving several parts of both the mucosal barrier itself, along with immune cells instructed by the luminal contents of the mucosae.

Mucosae consist of a single layer of epithelial cells covered by one or several layers of mucus on the luminal surface, making up the mucosal epithelium, which is in contact with the contents of *e.g.* intestines or lungs.¹⁵⁹ In the gut, the mucus layer(s) consist of mucin glycoproteins produced by goblet cells interspersed within the epithelium, which minimises direct contact of the epithelium with microbial intruders through stratification.¹⁶⁰ This is further reinforced by the large amounts of antimicrobial mediators, such as cathelicidins, defensins and C-type lectins secreted by *e.g.* enterocytes and Paneth cells, and substantial quantities of antigen-specific secretory IgA (SIgA) produced by plasma cells in the underlying lamina propria, rendering the mucosal epithelial lining virtually devoid of bacterial intruders.

On the apical side, the lamina propria makes up a collection of immunological structures of gut-associated lymphoid tissue such as the Peyer's patches, which consist of cells such as DCs, MØs and tissue-homing effector T and B cells. Microbe-associated molecular patterns in the form of microbial antigens from the gut lumen may be taken up by specialised epithelial cells known as microfold (M) cells in the mucosal epithelium¹⁶¹ or by DCs or MØs within the lamina propria.³⁵ Subsequently, these cells pass on the information to naïve Th cells in the draining mesenteric lymph nodes or Peyer's patches. There are both resident and migrating populations of APCs within the gut mucosa. Whereas migrating CD103+ DCs mainly induce immunomodulatory responses such as gut homing and induction of IgA class-switch in B cells^{162,163}, CX3CR1+ DCs and MØs chiefly participate in innate immune clearance of pathogens and supporting the responses of Th17 cells.¹⁶⁴⁻¹⁶⁶

The immune response is in the homeostatic state driven towards immunoregulation by the effects of tolerogenic APC and epithelial-derived TSLP, TGF- β and retinoic acid-production.¹⁵⁷ This drives the differentiation of Th cells towards IL-10 and TGF- β producing Treg cells, hence dampening inflammatory immune responses towards commensals. Furthermore, retinoic acid promotes a T cell-independent class-switch of B cells to become IgA-producing plasma cells, thereby further enhancing barrier integrity. Moreover, T and B cells of the lamina propria become imprinted for gut homing through retinoic acid dependent up-regulation of integrin $\alpha 4\beta 7$ and CCR9, rendering the cells re-circulating back to the gut upon egress from the lymph

nodes.¹⁶⁷ In case of bacteria penetrating the mucosal epithelium, the contact with the immune system is kept at a minimum by the compartmentalisation of the mucosal immune structures, and the commensal bacteria are rapidly eradicated by monitoring MØs as well as Th cells within the lamina propria.¹⁶⁰

The differentiation of Th cells within the mucosa-associated lymphoid tissues is largely dependent on microbial stimulation.¹⁶⁸ Different types of commensal species modulate immunity towards tolerance, emphasising the importance of exposure to a diverse microbiome for proper immune maturation. To control the continuous exposure to the extraordinary amounts of bacterial content in the gut, and prevent microbes from entering the circulation, the gut mucosa maintains a low-grade inflammatory response, represented by a high presence of mainly Th17 cells^{47,56}, and immune regulation by induced Tregs.⁴⁷ In animal models, segmented filamentous bacteria have been shown to drive the differentiation of Th17 cells^{169,170}, and very recent results suggest similar colonisation patterns in humans.¹⁷¹ Additionally, Tregs may also be induced by the microbiota.¹⁷² While Th1-derived IFN- γ promotes phagocytosis by MØs, thereby ameliorating microbial clearance, the Th17 cytokines IL17A, IL-17F and IL-22 drive recruitment of neutrophils and enhance barrier integrity by producing antimicrobial mediators.¹⁵⁹ Hence, the homeostasis in the gut mucosa is maintained by a balance between various types of Tregs producing TGF- β and IL-10, and the presence of ILCs, Th17 cells secreting IL17A, IL17F and IL-22 conferring low-grade inflammatory responses, all in all serving to maintain tolerance towards the commensal microbiome.¹⁶⁰

The oral mucosa differs from the gut mucosa in that the lamina propria is highly vascularised and covered by a stratified squamous epithelium, rendering it thicker than its gastrointestinal counterpart.¹⁷³ Despite this fact, it possesses a higher permeability, making it well-adapted for uptake of its rigorous exposure to a variety of antigens.⁴³ Similar to the gut mucosa, mucosa-associated lymphoid tissues such as salivary glands, tonsils and cervical lymph nodes compose inductive sites of oral immune responses.^{173,174} At these sites, antigen uptake takes place either directly by DCs or by M cell-like mediated delivery to DCs, which further orchestrate downstream effector T and B cell-responses within the oral mucosa lamina propria. The mucus of the oral mucosa – saliva – contains large quantities of IgA-antibodies, as well as antimicrobial substances and enzymes.¹⁷³ The induction of tolerance in the oral mucosa is similar to the induction of oral tolerance in *e.g.* the gut, which is characterised by the induction of protolerogenic antigen-specific T cell-responses, along with anergy, deletion and inhibition of present pro-inflammatory T cell-responses. However, the immense exposures to dietary antigens and commensal bacteria warrant a pro-tolerogenic default setting of the oral immune system, in order to not create inappropriate immune responses. For instance, DCs of the oral mucosa express the high-affinity IgE receptor Fc ϵ RI, with particularly high expression in atopic individuals¹⁷⁵, which may ligate to allergen-specific IgE.⁴³ Allergen recognition results in cross-linking of several receptors, and may lead to the production of indoleamine-2,3-dioxygenase as well as IL-10 by the DCs, which thereby promotes the development of tolerant immune responses.¹⁷⁴ Moreover, activation of TLRs on DCs and lymphoid cells upon microbial exposure leads to induction of tolerogenic immune responses, while at the same time silencing potential inflammatory responses to antigens. Additionally, APCs in the oral mucosa constitutively express B7-H, a molecule involved in the inhibition of co-stimulation needed for *e.g.* T cell activation.⁴³ Hence, T cell-responses within the oral mucosa are generally dampened. These mechanisms are similar to those present upon sublingual immunotherapy (SLIT).¹⁷⁴ As in the gut mucus, the

saliva contains considerable amounts of IgA produced by plasma cells present in salivary glands, along with antimicrobial mediators.⁴³

In the case of allergic inflammation, there is a breach in both the mucosal epithelial barriers as well as in tolerance.^{176,177} The impaired mucosal barriers enable entry of microbes and allergens, the increased exposures of which lead to the production of inflammatory cytokines such as IL-33, TSLP and IL-25 by the mucosal epithelium, and the promotion of Th2 cell inflammation in the mucosa-associated lymphoid tissues by pro-inflammatory DCs.¹⁶⁷ There is also a large body of evidence showing that allergic individuals possess lesser diversity in their microbiomes, which further may cause dysregulation of Treg-responses and promote uncontrolled inflammatory Th1- and Th17-responses.^{158,178-182}

Mucosal immune responses are indispensable for the development of tolerance towards beneficial bacteria and dietary components, while at the same time keeping pathogenic intruders at bay. This balance is maintained through the actions of Tregs and Th17 cells within the mucosa, as well as by the presence of large quantities of IgA and antimicrobial mediators within the mucus. Mucosal membrane integrity is often impaired in allergic subjects, leading to a pro-inflammatory environment with higher exposures to allergens, which predisposes for dysregulated pro-inflammatory immune responses.

Commensal microbiota – symbiotic bacteria important for appropriate immune maturation

Commensal microbes are beneficial to its host by the ability to protect the host from pathogenic invaders, for instance by competing for colonisation niches with appropriate adhesive or pH properties, as well as for binding sites and nutrients.^{183,184} Furthermore, the production of antimicrobial proteins and other metabolites may create an unwelcoming environment for pathogenic species.¹⁸⁴ The commensal microbiome is also of great importance for the sufficient uptake of nutrients from dietary components we are unable to digest. An example includes the digestion of complex carbohydrates in *e.g.* fibres, which is indispensable for endogenous vitamin-production and generation of short-chain fatty acids.¹⁵⁴ The composition of the microbiome is affected by several factors, including whether one lives in a rural or urban setting¹⁸⁵, the number of siblings or pets present in the family¹⁸⁶, and general environmental exposures.¹⁸⁷ However, considerable long-term contributions to the microbiome originate from maternal exposures throughout pregnancy, such as antibiotic use, dietary intake, smoking and exposure to pollution, as these affect the perinatal setup of the microbiome upon which postnatal exposures will build further.¹⁸⁸

The gut mucosa in humans is fully developed by the time of birth, whereas the immune compartments that have begun to develop *in utero* need diverse environmental exposures postnatally in order to develop the mature and tolerant immune responses seen in adults.¹⁸⁹ The communication between the developing microbiome and the maturing immune system early in life is indispensable for the establishment of a stable microbiome and the education of balanced immune responses.¹⁵⁶ This process equips the neonate with tools for tolerating or combating the multitudes of exposures encountered in postnatal life. The neonatal intestinal microbiome gradually evolves from certain bacterial communities to others, reaching a stable adult-like state at around 2-3 years of age.¹⁸⁸ This loosely defined time period, including foetal life, provides a window of opportunity for influencing the microbial communities and accompanying immune responses in the child.^{15,178} Extrinsic exposures throughout pregnancy and early postnatal life thereby have a huge impact on development of future health or disease.

The main process of colonising the different mucosal niches in the body begins throughout the process of birth, with urogenital maternal exposures laying the foundations for the tremendous colonisation and expansion of the microbiome in the new-born child.¹⁵⁷ The importance of this process becomes particularly clear in children born by caesarean section who have an altered microbiome, with decreased microbial diversity¹⁵², that mainly consists of species originating from the skin¹⁹⁰, while also being less similar to their mother's microbiome.^{152,183,191} This stands in contrast to the microbiome communities in children born vaginally that reveal high resemblance to their mother's microbiome¹⁸³ and show predominant survival of species originating from the maternal gut microbiome in their gut microbiota.^{192,193} Breastfeeding further reinforces the shaping of the vulnerable neonatal microbiome, not only by transferring nutrients, immune cells, cytokines and hormones, but mainly through the presence of SIgA and a unique microbiota in human milk.^{30,148} Accordingly, decreased microbial richness in the breast milk microbiome associates to the development of allergic disease.¹⁹⁴ Furthermore, human milk oligosaccharides favour the colonisation of commensals within the developing gut microbiome, by providing substrates for these species.¹⁴⁸ Indeed, the neonatal gut microbiome consists of species present in breast milk, such as *Bacteroides*, lactobacilli and bifidobacteria.¹⁹² The SIgA

provided by the mother constitutes passive immunity, through specific protection against pathogens encountered by the mother.¹⁵⁶ In the new-born child, the sensitivity to microbial stimulation is reduced, by means of lessened sensitivity of TLRs in epithelial cells.¹⁸⁹ Supposedly, this is in order to make way for inducing tolerance towards commensal species. Additionally, the presence of TGF- β and IL-10 in breast milk dampens the immune responses of the neonatal gut further, hence promoting the tolerance towards commensals.^{148,156} Altogether, commensal-induced effects on the intestinal epithelium promote tolerogenic DC development and subsequent Th cell differentiation, leading to tolerant unresponsiveness of the immune system towards beneficial communities of bacteria.

There is mounting evidence for altered commensal gut communities among allergic children. For instance, in children developing food allergy at age 1, the diversity of the gut microbiome was decreased and poised towards Enterobacteriaceae.¹⁹⁵ These findings confirm earlier findings, revealing lower gut microbial diversity among children developing IgE-associated eczema^{179,181} and allergic symptoms along with sensitisation until age 5.¹⁹⁶ Similarly, reduced gut microbial diversity in early life conferred a higher risk of being multi-sensitised¹⁹⁷ and developing asthma in early childhood.^{197,198} Conversely, the presence of lactobacilli and bifidobacteria in the first two months decreased the likelihood of allergy development at age 5 years.¹⁹⁶ Interestingly, in children with faecal presence of a triad of lactobacilli during the first two months of life, the risk of developing allergic disease was reduced until 10 years of age.¹²⁶ Furthermore, this led to decreased levels of the Th-associated chemokines CCL17 and CCL22 in the circulation of the same children, indicating that probiotics may have systemic effects as well.

Commensal microbes are necessary for optimal mucosal barrier function, and the composition of the microbiota is heavily dependent on early life exposures. Details of the timing, amount and diversity of environmental exposures needed for establishment of a healthy microbiome remain to be discovered. However, the impact of microbiota colonisation on immune development and function in the new-born child has far-reaching consequences for health and disease later in life.

IgA-antibodies – mucosal drivers of host defence and tolerance induction

IgA-antibodies make up the main isotype present in the secretions at mucosal surfaces, and it constitutes the most abundantly produced antibody in mammals.¹⁹⁹ The IgA-antibodies may exist as monomers, but more frequently they are present as homodimers joined by the joining J chain and parts of the poly-Ig receptor known as the secretory component.¹⁹⁹ The secretory component is a remnant from the poly-Ig receptor-mediated transcytosis through the epithelium, which protects the homodimeric SIgA complex from degradation.¹⁹⁹ SIgA-antibodies act through the process of immune exclusion, both in the sustenance of tolerance towards commensal bacteria, as well as in the targeted neutralisation of pathogens.⁴³ In this way, SIgA restricts the contact of microbes with the mucosal epithelium by neutralising the pathogenic intruders. The main sites of induction of SIgA in the gut are B cells within the Peyer's patches²⁰⁰, whereas plasma cells in salivary glands compose the chief producers in the oral mucosa.²⁰¹

It has been shown that SIgA increases the diversity of the microbiome, by constraining overrepresentation of single species^{202,203} and facilitating symbiosis between bacteria.²⁰⁴ Furthermore, commensal bacteria in the gut are coated with SIgA to a large degree^{205,206}, indicating the importance of SIgA in sensing the microbiome of the gut. Indeed, the proportions of IgA-bound faecal bacteria are lower in children developing allergic manifestations such as asthma throughout the first 7 years of life, compared to those who stay healthy.²⁰⁷ By transporting complexes of SIgA with bound antigen through M cells, the gut environment may be sampled and provide cues for promoting tolerance within the lamina propria, thereby preventing inappropriate immune responses from developing.²⁰⁵ Conversely, microbial colonisation is indispensable for proper induction of SIgA-responses, since intestinal DCs within the lamina propria need TLR-mediated signalling for their activation.²⁰⁸ Upon activation, intestinal DCs may sample the luminal content and through B-cell activating factor and APRIL²⁰⁹, to directly induce class-switching of B cells through the effects of TGF- β and IL-10^{209,210}, rendering them IgA-producing plasma cells.¹⁵⁹ This is a mucosa-specific process, which takes place alongside classical Th cell aided B-cell activation. These B cells have in the activation process become imprinted to home back to the gut mucosae and the lamina propria, where they become resident plasma cells secreting SIgA.

Differential regulation of IgA-responses has been implicated in relation to environmental exposures. For instance, throughout the first 5 years of life, Estonian children with higher exposures to bacterial endotoxin displayed elevated levels of SIgA in saliva compared to Swedish children.²¹¹ Interestingly, poly-Ig receptor expression is elevated upon bacterial exposures, hence increasing the transcellular transport of IgA to mucosal surfaces.²¹² The gut composition also affects the production of SIgA, as the number of present bifidobacteria species in faeces correlated with total SIgA levels²¹³. In relation to allergic disease, we have in previous studies shown that high salivary levels of SIgA at age one seemingly protects against development of late-onset wheeze at age four in sensitised children.²¹⁴ Moreover, lower levels of SIgA in saliva were observed in children with sensitisation and allergic symptoms, while this was not the case for children being only sensitised.²¹¹ Furthermore, colonisation with lactobacilli early in life associated with elevated salivary SIgA levels at six months and age 5 years²¹³, which is especially interesting as early life colonisation with lactobacilli associated with less allergy development at age five, regardless of parental allergy.²¹⁵

IgA-antibodies compose key players of mucosal immunity, contributing to promotion of tolerance while simultaneously preventing entry of pathogens through the mucosa. Dysfunction of mucosal barriers may lead to allergy development, and indeed altered gut and salivary IgA-responses have been observed in allergic individuals. However, the immunomodulatory effects of SLIT in the oral mucosa of children and adolescents have previously only been sparsely examined. In Paper IV, the impact of timothy grass pollen-specific SLIT on the salivary levels of total, secretory and timothy grass pollen-specific IgA-antibodies will be studied.

Combating the allergy epidemic – approaches to immunomodulating treatment

Today's treatment options for allergic disease comprise to a great extent symptomatic treatment, accompanied by considerable side effects.²¹⁶ While most of the available treatments on the market for allergic disease act mainly to alleviate present IgE-mediated symptoms (such as antihistamines) and dampen allergic inflammation (corticosteroids), large efforts are being made to introduce options that may modulate present and future development of allergic disease. Examples of this include the recently emerged biologics, which have improved therapeutic performance in allergic diseases such as asthma and eczema.²¹⁷ Effects of IgE-mediated mast cell and basophil degranulation may be blocked by the use of recombinant monoclonal anti-IgE-antibodies in severe IgE-associated asthma.²¹⁷ The inhibition of the effects of the Th2-associated cytokines may also be targeted with monoclonal antibodies against IL-4, IL-5 and IL-13 or their receptors, which have shown varying but promising success in clinical practice and/or clinical trials in several allergic diseases. However, none of these treatments may be used in prevention of allergic disease. As diverse microbial and dietary exposures are crucial for appropriate immune development in early life, which in turn may impact on allergy development, the use of probiotics and ω -3 fatty acids have become interesting targets for perinatal prevention.^{15,155,218} Furthermore, as allergen-specific immunotherapy has the potential to both alleviate symptoms of allergic disease, as well as prevent further manifestations from developing, it has become an attractive strategy in childhood allergy prevention.²¹⁹ In the coming sections, the putative halting of the allergic march by the above mentioned strategies will be elaborated.

Probiotics – amongst all microbes, who's a good bug?

Allergic diseases have in several aspects been related to microbial dysbiosis, possibly caused by modern-day habits decreasing both the amounts and total diversity of microbial exposures in our everyday lives.²⁷ Several studies corroborate this by revealing altered gut microbiomes in children developing allergic disease^{179,180,182,196,220–222}, which proposes that interventions that modulate gut microbiota could provide a meaningful approach in allergy prevention.²²³ To counteract this absence of crucial immunostimulatory microbes, the supplementation of beneficial bacteria – probiotics – has become an approach in possibly restoring both homeostatic microbiomes and immune responses in the host.

The World Health Organisation defines probiotics as “live microorganisms which when administered in adequate amounts confer a health benefit on the host”.²²⁴ Probiotics may have a positive impact on host immunity in several ways. The mucosal barrier function may be enhanced, for instance by strengthening the epithelial barrier and increasing the production of mucus and antimicrobial mediators by the mucosal epithelia.³⁵ Furthermore, probiotic bacteria may compete with pathogenic bacteria for necessary nutrients or niches, ultimately excluding them from the microbiome by inhibiting their life-cycle. More importantly, probiotics may additionally affect immune responses of the host by microbial components modulating Th cell differentiation and cytokine responses.³⁵ This is especially true for the small intestine, where the colonisation of commensals is relatively low, allowing for transient colonisation of the probiotics. The decisive factor for the initiation of tolerant or inflammatory immune responses is which type of pattern recognition receptor that recognises the pathogenic, commensal or probiotic antigen. The bacteria interact with different pathogen recognition receptors on innate immune cells, through cell wall components such as lipoteichoic acid, peptidoglycans and capsular polysaccharides.^{35,225} For instance, both peptidoglycan and lipoteichoic acid act as ligands for TLR2³⁵, and TLR9-signalling may be induced by sensing the high CG-content of bacterial DNA.¹⁶⁴ The subsequent production of retinoic acid and cytokines such as TGF- β and, the otherwise pro-inflammatory, TSLP from the activated innate immune cells (epithelia and resident APCs) may thereafter promote tolerance induction by influencing DCs within the lamina propria to modulate Th cell differentiation towards Tregs.¹⁶⁴

Several bacterial strains have shown beneficial effects as probiotic supplements, among them the *Lactobacillus* subspecies.²²⁶ Lactobacilli make up a group of Gram-positive bacteria, which are a natural part of the human microbiota in the skin, gastrointestinal tract, vagina and breast milk, to mention but a few.^{164,226} A vast variety of *Lactobacillus* strains exist, and although each strain exhibits strain-specific effects, the above described mechanisms of action for immunomodulation generally apply for the lactobacilli as well.²²⁷ *Lactobacillus reuteri* (*L. reuteri*) is one of the most studied lactobacilli, which has proven ability to bind to the mucosal epithelium of the gut and create biofilms.²²⁶ The bacterium produces several antimicrobial agents, such as reuterin, which may inhibit growth of other bacterial species, along with lactic acid, ethanol and acetic acid.²²⁶ Furthermore, *L. reuteri* is able to produce several essential vitamins, such as vitamins B9 and B12, and may also positively modulate the diversity and composition of microbiota in *e.g.* the gut.²²⁶ The maturation of DCs is also modulated by *L. reuteri*, by promoting the development of pro-tolerogenic APCs that drive differentiation towards immunoregulatory IL-10-producing Tregs.²²⁸ In these ways, lactobacilli may improve mucosal barrier function while at the same time induce tolerant immune responses.

The role of probiotics in restoring tolerance in allergy development has been extensively studied, especially in perinatal life. The combined pre- and postnatal treatment with probiotics seems the most favourable in terms of allergy prevention^{33,229}, since exclusive prenatal²³⁰ or postnatal²³¹⁻²³⁴ treatments have failed to show corresponding effects. In studies investigating pre- and postnatal supplementation with several different strains of lactobacilli, such as *L. rhamnosus*^{235,236}, *L. rhamnosus GG*^{235,237,238}, *L. paracasei*²³⁹, *L. acidophilus*^{238,240} and *L. reuteri*²⁴¹ alone and in combination with other probiotic strains, the prevention of eczema and/or IgE-associated eczema has repeatedly been shown. The allergy preventive effects of some of these studies also persisted later into childhood.²⁴²⁻²⁴⁵ Increased barrier function of the skin has also been proven in children with AD who were treated with *L. reuteri*.²⁴⁶ However, the evidence for general allergy preventive effects of probiotics on other allergic diseases such as asthma and ARC are still limited.²⁴⁷ Further studies in the above mentioned *L. reuteri* cohort²⁴¹ have demonstrated that faecal presence of *L. reuteri* during the first week of life associated with lower circulating CCL17 and CCL22 concentrations, and higher CXCL11 concentrations, at age 6 months.¹⁷⁹ Additionally, the responsiveness upon TLR2²⁴⁸ and allergen stimulation²⁴⁹ was decreased in children treated with *L. reuteri*.

Probiotics are beneficial microbes in the sense that they may modulate immune maturation and function, while also enhancing barrier integrity and preventing pathogens from intruding mucosal linings. The dose, timing and type of probiotics plays a crucial role for possible beneficial downstream effects. In children with a high propensity to develop allergic diseases, the pre- and postnatal use of probiotic strains such as Lactobacilli have proven partly effective in the prevention of allergic diseases. In Papers II and III, the possible modulation of L. reuteri on the epigenome will be studied alone or in combination with ω -3 fatty acids.

ω -3 fatty acids – modulators of inflammatory responses, as a matter of fat

Lipids comprise a diverse group of fatty acids and derivatives thereof²⁵⁰, which are major constituents of cellular membranes, involved in cellular signalling pathways as well as in energy storage for cellular metabolism.²⁵¹ In today's Westernised societies, fatty acids comprise a substantial proportion of ingested calories. The concurrent increase of diseases with inflammatory components suggest that fatty acids may provide important precursors to several mediators that initiate and sustain inflammation, apart from supplying dietary content. However, fatty acids are also contributing to resolution of inflammation, highlighting the significance of what type of fatty acids that are ingested.

The polyunsaturated fatty acids (PUFAs) ω -3 and ω -6 are termed essential for human health, due to the absence of the needed enzymes for endogenous production of related mediators.^{251,252} Hence, they need to be ingested through the diet, and proportions between ω -3 and ω -6 PUFAs dictate the nature of the downstream responses. In a succession of enzymatic steps, the enzymes cyclooxygenase, lipoxygenase and cytochrome P450 convert the PUFAs linoleic acid (ω -6) and α -linolenic acid (ω -3) to the intermediate long chain PUFAs (LCPUFA) mediators arachidonic acid (AA), and docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), respectively (Figure 5).^{76,253} The ω -3 LCPUFAs EPA and DHA are mainly contained in oily fish such as mackerel, salmon and herring, while ω -6 counterparts are found in amongst other things soybean and sunflower seed oils.²⁵³ EPA and DHA are usually present as soluble mediators in the circulation, and are transported to the site of inflammation through oedema²⁵⁴, whereas AA is

Figure 5. A brief overview of PUFAs and their downstream immune mediators.

principally integrated into the cellular membrane.⁷⁶ While AA generally gives rise to pro-inflammatory mediators such as prostaglandins and leukotrienes, EPA and DHA mainly contribute to the production of SPMs including protectins, resolvins and maresins (Figure 5).²⁵⁵ The AA-derived lipoxins also belong to this group of immunomodulatory mediators²⁵⁵, which restore homeostasis by resolving inflammation.²⁵⁴ Resolution of inflammation is achieved through the general enhancement of immune responses, by attenuating endothelial and epithelial permeability, and decreasing granulocyte recruitment to the inflammation site.²⁵⁵ Furthermore, the differentiation of plasma cells from B cells is promoted, as is also the phagocytosis of microbes and efferocytosis of apoptotic immune cells by MØs, while the production of pro-inflammatory

cytokines and chemokines is inhibited.^{254,255} In an allergic inflammation context, lipids are engaged in the maturation of APCs such as DCs, antigen processing and presentation, as well as in the process of transporting allergens across mucosal linings.²⁵⁶ Moreover, impaired SPM-responses have been observed in asthmatic individuals, both in the circulation as well as in the airways.²⁵⁵ In children with atopic asthma, the levels of total ω -6 fatty acids were elevated, whereas the levels of EPA seemed to have a protective effect of general asthma development.²⁵⁷ Furthermore, DHA inhibits B cells from isotype-switching to IgE-production in an IL-4/CD40 dependent manner²⁵⁸, and both EPA and DHA inhibit the release of IL-4 and IL-13 from mast cells, resulting in dampened production of reactive oxygen species.²⁵⁹ Altogether, this suggests that pathological features of allergic disease may originate from dysregulated lipid mediator responses. The intake of dietary fatty acids is in affluent societies mainly skewed in favour of the ω -6 LCPUFAs, thus supplying precursors for the production of pro-inflammatory mediators to sustain allergic inflammation. Indeed, ω -3 LCPUFAs make up just a small proportion of the phospholipids in the cell membranes of immune cells, which are dominated by ω -6 LCPUFAs in content.²¹⁸ The fatty acid composition in phospholipids of the cell membrane alters cellular function, as changes in the fluidity of the cell membrane modulates protein-protein and lipid-protein interactions, and hence influences signal transduction pathways within the cell.²¹⁸ Thus, by increasing the dietary intake of ω -3 LCPUFAs, the ratio could be modulated towards the production of the immunomodulatory SPMs, and thereby possibly modulate allergic inflammation.²¹⁸

Fatty acids are readily transported to the developing child via the placenta during gestation, and the preferential enrichment of LCPUFA in the foetus, especially throughout the last trimester, suggests crucial functions of these mediators for normal development.²⁶⁰ Studies of ω -3 fatty acids during pregnancy and after birth in allergy development have shown modulation of these mediators in allergic disease.¹⁵⁵ For instance, in a study supplementing ω -3 fatty acids as fish oil prenatally to the mothers, cord blood levels of IL-4 and IL-13 were indeed diminished.²⁶¹ Furthermore, the risk of asthma in the child was decreased after prenatal ω -3 fatty acid intervention during the last trimester of pregnancy.²⁶² Another study showed that the probability of having had asthma medication prescribed during the first 24 years of life were decreased in children whose mothers had been supplemented with ω -3 fatty acids in the last trimester of pregnancy²⁶³, suggesting these effects may be sustained in the long-term. In a study of postnatal ω -3 supplementation, treated children showed lower allergen-induced levels of IL-13, with concomitant increases in IFN- γ and TNF-responses in peripheral blood mononuclear cell (PBMC) cultures.²⁶⁴ In previous studies by our group, we demonstrated lower prevalence of IgE-associated eczema and sensitisation at 1 year of age upon combined pre-and postnatal ω -3 intervention²⁶⁵, which was accompanied by a lower risk of developing any type of IgE-mediated disease until 2 years of age in actively treated children.²⁶⁶

The ω -3 fatty acids EPA and DHA compose precursors for immunomodulatory mediators, such as the SPMs, which are of crucial importance in the resolution of inflammation. In the present-day world, the intake of ω -3 fatty acids is markedly decreased and may promote the presence of inflammatory diseases. In allergic diseases, the supplementation with ω -3 fatty acids has proven partly successful, but the mechanisms behind the effects remain incompletely elucidated. In Paper III, the possible epigenome modulating effects of prenatal ω -3 supplementation on cord blood CD4+ cells are studied.

Immunotherapy – re-creating tolerance by desensitisation

Allergen-specific immunotherapy (AIT) is a treatment method that has been around for over a century, and to date it still is the only treatment option that potentially may cure present allergic disease, and prevent further manifestations from developing.²⁶⁷ The prospect of AIT is to reconstruct tolerant immune responses towards the allergen against which an allergic individual was sensitised to begin with, thereby reversing IgE-mediated allergen sensitisation.²⁶⁸⁻²⁷⁰ This reinstatement of tolerance, essentially by the process of desensitisation, constitutes the remission of present allergen-specific immune responses, by means of anergy, apoptosis and inhibition of T cells, but also the induction of immunoregulatory innate and adaptive immune responses.

A variety of AIT treatments in different formulations are today available in clinical practice, mainly to alleviate symptoms of *e.g.* grass and birch pollen, bee venom and dust mite-induced allergic reactions²¹⁹, while several others are presently in clinical trials. AIT may be administered in different ways, depending on the characteristics of the allergen and the tissues the allergen affects the most. The dose of the allergen in AIT is predominantly a higher dose than normally encountered in everyday life for perennial and seasonal allergens.²⁷¹ This is important, as high doses of antigen exposures during Th cell differentiation favour Th1- over Th2-differentiation, while lower doses create the opposite constellation.^{51,52} Furthermore, in some AIT practices, a succeeding up-dosing regimen of the allergen is chosen, followed by a period of a maintenance dose, whereas others maintain steadily high doses throughout the treatment period.²⁷²

Subcutaneous immunotherapy (SCIT) was the first practically used method of AIT, which encompasses the regular administration of injections consisting of steadily increasing doses of the allergen into the skin over several years.²⁶⁷ As there is a risk for anaphylaxis in performing this procedure, the treatment needs to be administered at a hospital, which along with the invasiveness of the method provides drawbacks of the treatment. SLIT, comprising the daily administration of lyophilised tablets or droplets under the tongue during several consecutive years, has therefore in the past 30 years or so received attention in the AIT field.²⁷³ SLIT is considered a more patient-convenient administration route, as it may be administered at home after taking the initial dose under clinical surveillance. Furthermore, SLIT displays fewer and less severe side effects as compared to SCIT, owing to lower treatment doses, rendering SLIT superior to SCIT regarding tolerability and safety.²⁷⁴ However, while the compliance to SCIT treatment is higher than for SLIT treatment, children in general have a higher compliance irrespective of treatment.²⁷⁵ Both intervention types are efficient in the treatment of ARC^{275,276}, in favour of SCIT over SLIT²⁷⁴, and while SCIT is more efficacious in the prevention of both tree-and grass-induced asthma, SLIT alleviates asthma symptoms against grass only.²⁷⁵

The clinical efficacy of SCIT and SLIT have solid evidence in the treatment of ARC²⁷⁷ and promising results in prevention of asthma among children^{219,274,278,279}, but few studies have investigated the possibly allergy preventive effects of AIT treatment. This changed with the Preventive Allergy Treatment and the GRAZAX[®] Asthma Prevention (GAP) studies, where the prevention of asthma in children with ARC were studied, using SCIT²⁸⁰ and SLIT²⁸¹, respectively. In the Preventive Allergy Treatment trial, children with birch and/or grass pollen allergy were treated with SCIT injections for three years, and showed less symptoms of ARC and a lower risk of developing asthma at the end of treatment.²⁸⁰ These effects were also persisting five and ten years after treatment end.^{282,283} However, this study did not have a placebo-treated

control group to compare with²⁸⁰, in contrast to the GAP trial.²⁸¹ In the double-blind, placebo-controlled GAP trial, children with timothy grass pollen-induced ARC were treated with *Phleum pratense* (timothy grass pollen) SLIT for three years, and followed-up two years later.²⁸¹ The GAP study did not meet its primary outcomes of asthma prevention, which may be due to strict definitions of asthma, and that the test for evaluating lung function possibly was less appropriate for use in children.²⁸⁴ Notwithstanding, lower symptom and medication scores of both ARC and asthma were recorded, both during treatment and two years later, effects that were independent of season.²⁸⁴ Apart from these clinical improvements, long-term effects were seen in treated children on decreased SPT positivity, total and allergen-specific IgE levels, accompanied by augmented allergen-specific IgG₄ levels.

In the course of AIT, several immunological changes take place, involving both diminishing previous allergen-specific memory responses of the allergic individual, along with reconstructing tolerant immune responses towards the allergen, all in all resembling the state of a healthy individual.²⁶⁸⁻²⁷⁰ This is achieved by inhibiting effects of both the early and late phases of allergic reactions. An almost immediate effect is observed on the activity of basophils and mast cells, which becomes markedly decreased upon AIT initiation.²⁶⁸ This desensitisation results in less IgE-mediated degranulation and drives the probably most important effect of AIT: prevention of IgE-mediated anaphylaxis. The mechanisms behind this phenomenon are not completely elucidated, however, as the allergen-specific IgE-concentrations initially remain high and transiently even increase during the course of AIT, probably due to the exposures to large amounts of the sensitising allergen.²⁷⁰ However, later throughout treatment, there is a gradual decrease in the IgE levels, and potential seasonal increases are blunted.^{270,285} Interestingly, AIT also diminishes the frequencies of ILC2s²⁸⁶ as well as seasonal increases of these cells²⁸⁷, thus preventing further Th2-activation.

AIT affects DC maturation by modulating it towards tolerance, hence increasing the numbers of tolerogenic DCs.²⁷¹ This will in turn promote the differentiation of allergen-specific Th cells towards mainly Treg- and Th1-responses, while also initiating regulatory B (Breg)-responses.²⁷⁰ Throughout the course of AIT, these initial tolerant immune responses will be continually reinforced. Allergen-specific Bregs and Tregs mediate their effects upon activation by the secretion of the immunoregulatory cytokines IL-10 and TGF- β , as well as through the contact dependent actions of CTLA-4 and PD-1.²⁸⁵ The activity and function of pro-inflammatory Th2 cell subsets and downstream mediators are modulated through the inhibitory actions of IL-10 and TGF- β . Examples include the IL-10-mediated dampening of the production and effector functions of the Th2-associated cytokines IL-4, IL-5 and IL-13, and the inhibition of IgE-dependent activation of mast cells.^{36,74} Additionally, upon induction of Th1-responses and its associated cytokine IFN- γ , these immunomodulatory cytokines may induce isotype class-switch in B cells, enabling the production of allergen-specific IgA (TGF- β), IgG₁ (IFN- γ) and IgG₄ (IL-10), respectively.²⁷⁰ IgA-, IgG₁- and IgG₄-antibodies virtually act as competitive inhibitors of IgE-mediated allergen binding, thereby preventing further Fc ϵ RI-dependent sensitisation and mast cell degranulation from taking place. IgG₁- and IgG₄-antibodies may furthermore hinder present memory B cells from producing allergen-specific IgE by preventing IgE-allergen complexes from binding to the low-affinity Fc γ RIIb on the B cell.²⁷⁰ This attenuates the IgE-dependent activation of mast cells and basophils in two ways: firstly by decreasing the amount of allergen-specific IgE-antibodies, as fewer B cells produce them, and secondly the IgG- and IgA-antibodies may compete with IgE for allergen binding. This coincides with the decrease in SPT positivity that is evident later in the treatment, caused by the diminished expression of allergen-

specific IgE-antibodies.²⁸⁵ Later in the course of AIT, the otherwise season dependent increases of mast cells, eosinophils and basophils in the tissues are also absent, while the degranulation rate is maintained at low levels. Furthermore, the increasingly tolerant microenvironment in the treated individual inhibits the recruitment and cytokine-mediated functions of Th2 cells and eosinophils on the tissues. With the concurrent increases in Th1 cells, corresponding IFN- γ mediated effects and presence of IgG₁-antibodies, this along with the immunomodulatory responses of Bregs and Tregs, establishes a novel immune balance in the immune system.

There are treatment-specific effects on *e.g.* the humoral immune response during allergen-specific immunotherapy. While SLIT to a higher extent induces IgA- and IgG₁-production, SCIT mounts stronger IgG₄-responses, which can be explained by the different environments in which the initial immune responses has been initiated.^{270,285} Furthermore, in dust mite-allergic subjects both SCIT and SLIT treatment induced higher presence of CD4+CD25+FoxP3+ Tregs compared to both controls groups and base line levels.²⁸⁸ However, there was a correlation with clinical improvement only for SLIT.

Over the course of several years, AIT establishes a new balance in the immune system, by modulating the early and late phase allergic reactions. These changes consist of decreasing IgE-mediated immune responses, while at the same time inducing allergen-specific immunomodulatory Treg- and Breg-responses, and counterbalancing Th1-responses. Cytokines secreted by these cells shift humoral immune responses towards elevated IgG₁, IgG₄ and IgA-production, which may further prevent IgE-production, mast cell activation and inhibit the effects of present allergen-specific Th2 cells. Altogether, these events promote the maintenance of peripheral tolerance. However, the immune effects of SLIT in children with timothy grass pollen allergy have thus far not been described. In Paper IV, humoral and systemic aspects of P. pratense SLIT in children from the GAP study are the objective of the examinations.

Aims and hypotheses of the thesis

The general aim of this thesis was to study different perspectives of immune maturation from birth throughout adolescence by investigating systemic, mucosal and epigenetic immune mediators in relation to allergy development. The examinations of these mediators were performed both by studying the natural course of allergy development (Paper I), and by analysing how immunomodulating treatments in perinatal life (Papers II and III) and childhood (Paper IV) affect development of allergic disease.

Paper I

Aim: To assess circulating levels of the allergy-related chemokines CCL17, CCL18, CCL22, CXCL10 and CXCL11 at birth and in childhood (ages 1 and 8), in relation to development of asthma and sensitisation from infancy throughout adolescence (ages 1-16 years) in an observational birth cohort.

Hypothesis: Allergy development will be associated with and predicted by elevated levels of the Th2-related chemokines CCL17 and CCL22, as well as the Th2/Treg-associated chemokine CCL18. In contrast, decreased circulating levels of the Th1-associated chemokines CXCL10 and CXCL11 will relate to and precede development of allergic outcomes.

Paper II

Aim: To study epigenetic effects of pre- and postnatal *L. reuteri* intervention at birth, 1 and 2 years of age in relation to immune maturation and allergy development, by examining DNA methylation patterns in CD4+ cells both at four Th cell-related loci and on an epigenome-wide scale.

Hypothesis: Probiotic intervention will lead to increased transcriptional accessibility, reflected by hypomethylation, at immune-related sites on both the local and global DNA methylation level.

Paper III

Aim: To examine epigenome-wide effects of combined prenatal *L. reuteri* and ω -3 fatty acids on DNA methylation patterns in cord blood CD4+ cells.

Hypothesis: The probiotic intervention will lead to general hypomethylation in the epigenome, with possible pronounced effects compared to Paper II due to the earlier initiated treatment. Treatment with ω -3 fatty acids will show the same pattern in immune-related loci, and the combination of the two interventions may work synergistically to create the largest DNA methylation differences in immunological pathways.

Paper IV

Aim: To investigate systemic and mucosal immune responses upon a three-year treatment with grass pollen SLIT in timothy grass pollen-allergic children, by measuring circulating allergy-related cytokines and chemokines, as well as salivary IgA-responses.

Hypothesis: The active SLIT treatment will lead to the induction of tolerant immune responses by increasing Th1- and Treg-responses while simultaneously decreasing Th2-responses. The levels of timothy grass pollen-specific IgA-antibodies in saliva will also increase.

Study design and methodological considerations

In this thesis, the work on four different childhood allergy cohorts is described. Collectively, several parts of the mechanisms behind allergy development and possible allergy prevention strategies in childhood were studied. In Paper I, the role of chemokines as predictors of allergy development was investigated in an observational cohort of children. In Papers II and III possible allergy preventive effects of pre- and postnatal supplementation with either probiotics alone (Paper II) or in combination with ω -3 fatty acids (Paper III) were studied at the level of epigenetic regulation. In Paper IV, the effects of sublingual timothy grass pollen-specific immunotherapy on immune responses in grass pollen-allergic children were studied. In the sections below, details on each of the specific studies, and which analyses that were performed within each project are described.

Paper I – Chemokine studies in the Manchester Asthma and Allergy Study (MAAS)

In Paper I, the involvement of circulating chemokines in the natural course of allergy development was investigated (Figure 6). This was studied in children from the Manchester Asthma and Allergy Study (MAAS)²⁸⁹, an observational population-based birth cohort, which prospectively studies the development of allergies and asthma in childhood and adolescence. The study collected at total of 1184 mother-child pairs with eligible samples and variable information, from a mixed urban-rural population within the Manchester and Cheshire regions in the United Kingdom. The children have been followed up regularly for almost two decades, with available follow-up data at different intervals in the ages 1-19 years.

For the purpose of this study, information on asthma development and sensitisation from the clinical follow-ups performed by a paediatrician evaluating the development of allergic disease in the children at ages 1, 3, 5, 8, 11, and 16 years, were included. SPTs were also performed at all ages, for seven allergens (house dust mite (*Dermatophagoides pteronyssinus*), cat, dog, grass pollen, moulds, cow's milk, and egg white) until age 8, and nine allergens thereafter (addition of birch pollen (*Betula verrucosa*) and peanut). A test was considered positive when the mean wheal diameter was at least 3 mm larger than that for the corresponding negative control. Asthma was defined as fulfilling two of the following three criteria: current wheeze, current use of asthma medication, or physician-diagnosed asthma ever at any of the studied time points. Informed consent was provided in written form by parents or legal guardians, and the study was approved by the South Manchester Local Research Ethics Committee. Ashley Woodcock is the holder of the ethical permits for the whole study, for the start of the study (including the 1 year follow-up, Dnr ERP/94/032), the amendment for cord blood analyses (ERP/95/137) and the follow-up at age 8 years (Dnr 03/SM/400).

In order to study the association of chemokines with allergy development, the levels of five allergy-related chemokines were measured by means of ELISA (CCL18) and Luminex (CCL17, CCL22, CXCL10 and CXCL11) methodologies in 905 plasma samples from the MAAS cohort. Samples were available at three time points, from the time of birth (cord blood, n=376), 1 year (n=195) and 8 years (n=334) of age. The resulting chemokine concentrations were thereafter associated to outcomes such as asthma and sensitisation throughout childhood and adolescence,

Statistical analyses:

Association studies to development of asthma and sensitisation using cross-sectional logistic regression and longitudinal generalised estimating equation (GEE) models

Figure 6. An overview of the study protocol in Paper I. Blood samples were collected at birth (cord blood), 1 year and 8 years of age, and measurements of the five allergy-related chemokines were performed using Luminex (CCL17, CCL22, CXCL10 and CXCL11) and ELISA (CCL18) methodologies. The plasma levels of the chemokines were thereafter associated to the development of asthma and sensitisation between 1-16 years of age both cross-sectionally and longitudinally using logistic regression and GEE models, respectively.

as well as to previously machine learning-derived allergy clusters²⁹⁰⁻²⁹³, to assess involvement in allergy development. Subsequently, possible predictive and longitudinal effects on allergy outcomes of the circulating chemokine levels were investigated using logistic regression and generalised estimating equation (GEE) models.

Paper II – DNA methylation studies in the *Lactobacillus reuteri* (*L. reuteri*) cohort

In Paper II, both local and global epigenomic effects of pre- and postnatal probiotic intervention were surveyed in CD4⁺ T cells from children included in the *L. reuteri* study (Figure 7). The samples originated from a randomised, double-blind, placebo-controlled probiotic trial in which allergy prevention by the pre- and postnatal supplementation of the probiotic strain *L. reuteri* was studied (ClinicalTrials.gov-ID: NCT01285830).²⁴¹ In this study, the in total 188 included infants had a family history of allergic disease, consisting of at least one family member having

Figure 7. An overview of the study protocol in Paper II. Blood samples were collected at the time of birth (cord blood), 1 year and 2 years of age, and cord blood or peripheral blood mononuclear cells (CBMC/PBMC) were isolated from these samples. Upon purification of CD4⁺ lymphocytes, one part was used for flow cytometry determination of cell type proportions of naïve and memory cells. From the remaining cells, genomic DNA was extracted and utilised for the analyses of DNA methylation patterns using two different methods. A Th subset locus-specific method was used to inspect the DNA methylation status of four CpGs in the promoters of the Th cell-associated genes *IL-17*, *IFNG*, *FOXP3* and *IL-13*. Epigenome-wide DNA methylation patterns were studied using the Illumina 450K array. Results of the two methods were evaluated using both statistical and bioinformatic approaches.

eczema, asthma, gastrointestinal allergy, allergic urticaria or ARC. The treatment was initiated prenatally, from gestational week (GW) 36 in the pregnant women, who either received *L. reuteri* ATCC 55730 (5 droplets, 1×10^8 colony forming units (CFU), BioGaia®, Stockholm, Sweden) or placebo once daily until delivery. The postnatal treatment was started on day 1-3 after birth in the infants with the same product and dose daily throughout the first year of life. The probiotic preparation consisted of freeze-dried *L. reuteri* suspended in coconut and peanut oil containing cryoprotective components, while the placebo preparation only comprised the vehicle. The children were monitored by research nurses at multiple time points and follow-ups were performed by a paediatrician at 2 and 7 years of age. SPTs were performed at ages 6, 12 and 24 months using egg white, fresh skimmed cow's milk as well as standardised cat, birch (*B. verrucosa*) and timothy grass extracts. At 7 years of age SPTs included the mentioned allergens as well as dog, peanut and house dust mites (*Dermatophagoides pteronyssinus*). Evaluated allergic diseases included eczema, ARC, urticaria, asthma and gastrointestinal allergy. Informed consent was obtained from both parents before inclusion. The study was approved by the Regional Ethics Committee for Human Research in Linköping, Sweden. Bengt Björkstén holds the ethical permit for the initial analyses until the follow-up at age 2 years (Dnr 99323), while Maria Jenmalm holds corresponding responsibility for the ethical permit including the follow-up at age 7 years (Dnr M171-07).

From blood samples retrieved at three time points; at birth (cord blood), 1 and 2 years of age, CD4+ cells were isolated from CBMCs or PBMCs. On the CD4+ cells, flow cytometric studies were performed to examine the proportions of naïve (CD4+CD45RA+) and memory (CD4+CD45RA-) cells within the population. From the remaining CD4+ cells, genomic DNA was extracted, which in turn was utilised for studying whether treatment with *L. reuteri* affected epigenetic modifications in terms of DNA methylation. This was studied using two different methods: one surveying DNA methylation patterning at Th cell-specific loci and one studying the same on an epigenome-wide scale. The former approach assays DNA methylation patterns at four distinct CpGs in the promoters of *IFNG*, *IL-13*, *IL-17* and *FOXP3* using a Th subset locus-specific DNA methylation analysis method²⁹⁴, as a proxy to study polarisation into the Th1, Th2, Th17 and Treg cell subsets, respectively. For this analysis, samples (expressed as total (placebo/*L. reuteri*)) were included as follows: 32 (17/15) samples from the time of birth (cord blood), 24 (9/15) samples from age 1 and 35 (15/20) samples from age 2 years. Additionally, corresponding results from 8 healthy and 8 allergic adults (age- and sex-matched, mean age=28.9 years) were included for a comparison of immune maturation later in life. Associations to development of allergic disease as well as intervention groups were examined for all samples. Epigenome-wide DNA methylation analyses were carried out using the Illumina Infinium 450K array, with which more than 485 000 CpG sites spread over the entire genome were investigated.²⁹⁵ For these analyses, 19 (10/9) samples were available from cord blood; 10 (6/4) samples at age 1; and 19 (11/8) samples at age 2 years. Downstream statistical analyses of the DNA methylation data upon pre-processing included identifying DMPs and differentially methylated genes (DMGs), as well as network analyses and the development of a classifier.

Paper III – DNA methylation studies in the PRObiotics and OMega-3 (PROOM-3) study

In Paper III, the role of global DNA methylation patterning in CD4+ T cells upon prenatal combined *L. reuteri* and ω -3 fatty acid intervention were studied in new-borns from the

PROOM-3 study (Figure 8). In this experimental study, cord blood samples from 63 children in the presently on-going randomised double-blind placebo-controlled trial PROOM-3 (PRObiotics and OMega-3, ClinicalTrials.gov-ID: NCT01542970) were included. Included study subjects exhibited a family history of allergic disease, as described for the *L. reuteri*-study above. The study consists of four arms; double active treatment (*L. reuteri* and ω -3 fatty acids, L ω), single active treatments (active treatment for either *L. reuteri* (LP) or ω -3 fatty acids (P ω), placebo for the other) and double placebo treatment (PP). The pre- and postnatal probiotic treatment regimens comprise supplementation of the mothers from GW 20 until delivery with *L. reuteri* or the corresponding placebo, and the children from days 1-3 after birth throughout the first year of life. The active probiotic intervention consists of the probiotic strain *L. reuteri* DSM 17938 (BioGaia®, Stockholm, Sweden) suspended in refined coconut and peanut oil droplets, given twice daily to the mothers (2 $\times$ 20 droplets, 10⁹ CFU) and once daily to the children (1 $\times$ 5 droplets, 10⁸ CFU), whereas the placebo treatment consists of refined coconut and peanut oil droplets without probiotics.

Likewise, the ω -3 fatty acid treatment or corresponding placebo are supplemented from GW 20 throughout the first three months of lactation to the mothers. The mothers ingest either three capsules of Pikasol® (1 g capsules containing 640 mg ω -3 LCPUFA (35% EPA, 25% DHA), Orkla Health (previously Axellus), Lund, Sweden) or the placebo supplement olive oil twice daily. A variety of samples are collected from both mothers and children at several time points throughout pregnancy and infancy. Blood samples from children are collected at birth (cord blood), 6 months, 1 and 2 years of age. Allergy questionnaires surveying the child's health are filled out at ages 3 and 6 months, 1 and 2 years by the parents. Clinical follow-ups are performed on the children at ages 1, 2, 3, 4, 6 months and 1 year of age by a research nurse, and by a paediatrician at 2 years of age. As the study is still on-going, only cord blood samples were included in the present analysis, with follow-up data from parental reported questionnaires available from the 3-month time point. Hence, no information on subsequent allergy development could be included in this sub-study. Informed consent was obtained from both parents prior to inclusion in the study. Ethical permission for this study was granted by the Regional Ethics Committee for Human Research in Linköping, Sweden. Karel Duchén holds the ethical permit for the PROOM-3 study (Dnr 2011/45-31).

In Paper III, global analyses of DNA methylation patterns in CD4⁺ cells from CBMCs were investigated using the Illumina Infinium MethylationEPIC array, investigating more than 866 000 CpG sites in the epigenome.²⁹⁶ Samples were available from the time of birth from all four treatment groups in the following numbers: L ω = 18, LP = 16, P ω = 15 and PP=14. Upon isolation of CD4⁺ cells, the proportions of naïve (CD4⁺CD45RA⁺) and memory (CD4⁺CD45⁻) cells within the CD4⁺ cell population were determined using flow cytometry. DNA was extracted from the remainder of the CD4⁺ cells and was further utilised to study whole epigenome-wide DNA methylation patterns. Statistical and bioinformatic analyses of the DNA methylation data were carried out, including pre-processing, identification of DMPs and DMGs, pathway enrichment analyses and the identification of intervention modules using network approaches.

Figure 8. An overview of the study protocol in Paper III. Blood samples from the time of birth (cord blood) were included in this study, from all four intervention groups. Upon purification of CD4+ lymphocytes from CBMC, one part was used for flow cytometry determination of cell type proportions of naïve and memory cells. From the remaining cells, genomic DNA was extracted and subsequently used for whole epigenome investigations of DNA methylation patterns by means of the Illumina MethylationEPIC 850K array. Results were processed using both statistical and bioinformatic approaches to determine the biological importance of the interventions. L – *Lactobacillus reuteri*, ω – ω -3 fatty acids, P – placebo.

In Paper IV, diverse immunological responses were investigated in children with timothy grass pollen-related ARC upon treatment with SLIT in samples from the GRAZAX® Asthma Prevention (GAP) trial.²⁸¹ In this international randomised, double-blind, placebo-controlled multi-centre study, timothy grass pollen-allergic children were treated once daily with the sublingual timothy grass pollen immunotherapy tablet GRAZAX® for three consecutive years, followed by an open follow-up period of two years (Figure 9). The primary outcome of the study was time to development of asthma, as measured by reversibility of lung function.

Children in the ages 5–12 years of age with concurrent ARC and established sensitisation towards timothy grass, based on SPT positivity (wheal diameter ≥ 3 mm) towards timothy grass (*P. pratense*) and positive *P. pratense*-specific IgE levels (sIgE ≥ 0.7 kU/L) in serum, were included in the study. The active treatment consisted of lyophilised SQ sublingual tablets containing standardised timothy grass pollen-allergen extracts (75 000 SQ-T) from *P. pratense* (GRAZAX®, ALK, Hørsholm, Denmark). The placebo treatment consisted of a sublingual tablet without grass pollen-extract, in appearance non-distinguishable from the active treatment. The whole study included 812 children from several countries and study sites, and in Sweden 28 children were included from the two study sites, one in Stockholm (Sachs' children and youth hospital at Södersjukhuset) and one in Linköping (Allergy Center at Linköping University Hospital).

For the performance of this study, blood and saliva samples were collected from the children at three time points, following the grass pollen season in the particular year: at the randomisation visit (2010), after three years of treatment (2013) and two years after treatment cessation (2015). Data for SPTs (timothy grass, birch (*B. verrucosa*), mugwort, horse, dog, cat, house dust mites (*Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*) and moulds (*Alternaria alternata*, *Cladosporium herbarium*)), IgG₄ and IgE levels were measured at clinical follow-up visits throughout the treatment period. Information about ARC symptoms were collected at different time points using the visual analogue scale (VAS) score (records from visits 2, 9 and 14 in peak to end of respective grass pollen season, asking the question “How has your hay fever been last week?”) and the allergic rhinitis severity (ARS) score (daily diary records for 14 days in peak to end of grass pollen season 2015, asking the question “How has your hay fever been today?”). In the Swedish sites, a total of 17 children completed the study, with blood and saliva samples available from 25 unique individuals. Informed consent was obtained from parents or legal guardians of the children. The study was approved by the Regional Ethics Committee for Human Research in Lund, Sweden. Inger Winnergård is the holder for the ethical permits of the study, composed of a permit granted for the GAP study sites in Sweden (Dnr 2009/551) and an amendment for studies of Th cell-responses and IgA-measurements at our study centre (Dnr 2010/430). The study has further been approved by the Swedish Medical Product Agency (2009-011235-12) and the European Union Drug Regulating Authorities (EudraCT: 151:2010/48756).

The GAP study

Blood and saliva
sample collection

Treatment period
Clinical follow-up period

Inclusion

Blood
n = 20

Saliva
n = 17

Year 3

n = 15

n = 14

Year 5

n = 15

n = 16

Blood sample analyses:

Saliva sample analyses:

Statistical analyses:

Cross-sectional and longitudinal association and correlation studies of measured mediators and clinical outcomes

Figure 9. An overview of the study protocol in Paper IV. Blood and saliva samples were collected at the time of inclusion in the study, just after the three-year SLIT treatment was ended and two years after treatment cessation. In the blood samples, measurements of the allergy-related chemokines CCL17, CCL22, CXCL10, CXCL11 (Luminex) and CCL18 (ELISA) were performed in plasma. Additionally, allergen stimulations with timothy grass (*P. pratense*) and birch (*B. verrucosa*) were performed in isolated PBMC. In the collected saliva samples, levels of total IgA and secretory IgA, as well as Phl p 1-specific IgA were measured by ELISA and Luminex methodologies, respectively. Statistical analyses included both cross-sectional and longitudinal comparisons, in association and/or correlation to clinical and immunological outcomes.

Briefly, in Paper IV, peripheral blood and saliva samples were available in the following numbers at the respective time points (n=Blood/Saliva); inclusion (prior to treatment, n=20/17), year 3 (after three years with treatment, n=15/14), and year 5 (two years after treatment ended, n=15/16). From the blood samples, PBMCs were isolated and stimulated with timothy grass (*P. pratense*) and birch pollen (*B. verrucosa*) allergens. Supernatants from these experiments were analysed by means of ELISA (IL-10) and Luminex (IFN- γ , IL-5, IL-10, IL-13, CCL17 and CXCL10) methodologies, to examine content of allergy-related cytokines and chemokines. Additionally, plasma samples were retrieved from the blood samples and analysed regarding cytokine and chemokine content with ELISA (CCL18) and Luminex (CCL17, CCL22, CXCL10 and CXCL11). Collected saliva samples were isolated and heat-inactivated prior to analysis of total and secretory IgA (ELISA) as well as Phl p 1-specific IgA (1-plex Luminex) levels. Associations of the collected data to intervention group belonging, clinical outcomes such as ARC symptoms, as well as previously collected information on SPTs, IgG₄ and IgE levels were performed. Furthermore, the stimulated cytokine Th2/Th1-ratios were also investigated in relation to intervention group.

Ethical considerations

In the four described projects of this thesis, several ethical aspects of the performed research appear. As the research is performed in children, their parents or legal guardians have in all projects provided informed consent on their behalf. However, the children received age-adapted information about the study and, when feasible, also the opportunity to sign their own consents as a preparation for the parental consent. That means that older children got this opportunity, whereas smaller children did not have this choice. This could, especially when the children grow older, become a matter of discussion, as the decision to participate or not initially did not take into account the opinion of the child. However, it may be assumed that parents make these decisions in their children's best interest, and that no child is forced to participate against their will. Furthermore, as the families are allowed to withdraw from study participation at any time point without further explanation, this also provides a safety mechanism for the included children.

Retrieval of biological sample materials from these children provide another important ethical consideration. In the informed consents, the parents are also asked whether or not they agree to the long-term storage of these samples for other purposes than the particular study at hand, to which they have agreed to provide samples from their children. In the four projects, children from new-borns to adolescents have provided blood samples, which allows for studies of peripheral immune responses in the circulation. The process of retrieving peripheral blood

samples is an inevitably invasive procedure, which many children may find uncomfortable and even fearsome. In order to decrease suffering for the children, the younger children in particular have been given the opportunity for local anaesthesia at the blood withdrawal site. Moreover, the children have not been forced to multiple attempts of blood withdrawal, unless they consented to this at the occasion. When collecting blood samples from new-borns (Papers I-III), this has exclusively included the withdrawal of blood from the umbilical cord, hence not posing a painful invasive measure on neither the neonate nor the mother. This material mainly consists of cells and mediators of foetal origin^{297,298}, although possible maternal contamination could take place during sample collection.

Saliva samples comprise another interesting matrix to study in immunology, especially for examining mucosal immune responses. For the analyses in Paper IV, saliva samples were collected from children and adolescents by means of passive drooling using a vacuum pump connected to a so-called tracheal suction set. This procedure is non-invasive and the only restrictions on the children have been not to eat within the last half-an hour before the collection of the samples.

In the projects described in Papers II and III, the possible allergy preventive effects of pre- and postnatal *L. reuteri* alone (the *L. reuteri* study) or in combination with ω -3 fatty acids (the PROOM-3 study) were studied. In both projects, the interventions were initiated during pregnancy, in the last month of pregnancy in the *L. reuteri* cohort, and in the latter half of pregnancy in the PROOM-3 cohort. This poses additional ethical questions, as the prenatal supplementation presumably does not only affect the becoming mother upon ingestion, but also the developing foetus. Any effects of the prenatal treatment may consequently be upheld and potentiated by the postnatal treatment, and quite possibly may influence the long-term health of the infant as it grows up. The hypotheses of these studies compose the putative induction of tolerant immune responses through the interventions, preferably also with long-term effects. This could initially sound like a potentially dangerous experiment. However, the studied probiotic strains and ω -3 fatty acids given to the pregnant mothers, and later their babies, have been carefully studied and thoroughly scrutinised for potential toxicity and side effects in several studies over long periods of time.^{299,300} Bearing this in mind, it is also quite possible that the mothers nowadays themselves may have begun supplementations of other probiotic strains and ω -3 fatty acid supplements, as the putative immunomodulating benefits of these treatments during pregnancy have become more well-known. This is, however, an obvious exclusion criterion for these studies, as this would confound the outcomes of the projects.

Also, in Papers II and III, studies on epigenetic modifications were performed. Being the translating mechanism responsible for the adaptation of the principally rigid genome to ever-changing environmental factors, the question comes up whether these types of analyses pose a threat to the personal integrity of the included study participants. In general, although epigenetic modifications such as DNA methylation patterns are considered highly stable, they are also immensely dynamic to environmental stimuli. Epigenetic patterns studied in a particular moment provide a snapshot of how the transcriptome is regulated at that time, while in another it may look completely different. Although the exact mechanisms behind the temporal and spatial regulation of DNA methylation are not completely elucidated at present, DNA methylation could be used as a biomarker for health and disease. This could provide valuable information about ways in which life-style factors could be modified in order to improve health, with putative utility in personalised medicine. As in the *L. reuteri* and the PROOM-3 studies, possible alterations of pre- and postnatal interventions on epigenetic DNA methylation patterns were

sought as one of several possible mechanisms of action. This approach is on an informational level³⁰¹, as the interventions themselves thus far are not known to mainly act on an epigenetic level, although the aim of the studies in Papers II and III is to investigate such mechanisms. Furthermore, one may argue that the information retrieved from these children may neither be used to identify the specific children at the time point of sample retrieval nor later in life, due to the plasticity of epigenetic modifications. Although we in the *L. reuteri* study for instance developed a classifier for the outcome of the allergic response from the DNA methylation patterns in the children, this could not be used to identify the children on an individual level. Information on epigenomic patterns has from our point of view been handled the same way genomic information is handled, in a pseudonymised fashion with initial intervention blinding prior to statistical analyses, making it impossible to draw conclusions on samples origin or biasing possible treatment effects. As the epigenomic effects of the investigated interventions are just beginning to be clarified, the ethical aspects of personal responsibility in terms of health outcomes on both a personal level and future generations in *e.g.* dietary choices throughout pregnancy, will need further discussion.

In the GAP study, which is described in Paper IV, the possible asthma preventive effects of timothy grass pollen SLIT in children with timothy grass pollen-induced ARC was studied. SLIT has in recent years been noticed as a more patient attractive option compared to SCIT, due to its easy supplementation procedure along with its solid tolerability and safety profile. By the supplementation of standardised amounts of the timothy grass pollen allergen over a long period of time (three years), the primary aim of the main study was that the treatment will prevent development of asthma in children with timothy grass pollen-induced ARC, while secondary outcomes predicted the alleviation of ARC symptoms due to induction of tolerance against timothy grass.²⁸¹ Ethically speaking, supplying pure allergen towards which the children are sensitised may seem contradictory, as this intuitively would cause the patient harm. Although the administration of the tablet under the tongue in the initial treatment period may cause swelling or itching locally, severe adverse effects such as anaphylactic reactions rarely occur. Over time, the symptoms eventually subside, and during the study period, the children had access to standard rescue medication to ease their ARC symptoms. If the children would experience any asthma symptoms during their participation in the trial, they were encouraged to schedule additional visits to investigate and alleviate their symptoms. Hence, one may argue that the children in this study were under continual observation, as they had regular contact with the allergy clinics, with possible additional visits and medical treatments easily accessible in case of emergency.

Another ethical aspect of the performance of our studies lies in the sponsored collaborations, which provide the basis of the studies in Projects II-IV. In these studies, different types of interventions that are presently either experimentally validated or tested in clinical trials (*L. reuteri* and ω -3 fatty acids, Papers II and III) or are available on the market (GRAZAX[®], Paper IV) were investigated in collaboration with the manufacturers of the interventions. In Papers II and III, BioGaia[®] supplied the investigated probiotic strain, and partially supported the *L. reuteri* trial (Paper II) financially during the first two years of the study. The ω -3 fatty acids in the PROOM-3 study (Paper III) were provided by Orkla Health (previously Axellus), and in the GAP study (Paper IV), the grass pollen-specific SLIT tablets were supplied by ALK-Abelló A/S. In none of the mentioned sponsored collaborations did the sponsors have any influence on the research questions pursued, the statistical analyses performed or the interpretation of the experimental findings in preparation for publication. As the independence of the academic

researchers was ascertained and separated from the possible financial interests of the sponsoring companies, the ethical issue of working with profit-driven organisations may in this instance be considered negligible.

From an ethical point of view, the performance of the above described studies may be justified by several advantages. The first considers the knowledge gained by performing studies both studying the natural course of allergy development without intervention (as in Paper I) and the outcomes of using various putatively allergy preventive interventions (as studied in Papers II–IV). Understanding the background to why allergies develop at all in the first place will provide insights into how one may treat allergic disease and, in the best of cases, even prevent allergic disease from developing. Furthermore, studies on interventions with possible allergy preventive effects would add knowledge about how the immune system may be modulated in allergy prone children. This would in several ways promote the aspiration of the greater good on multiple levels. On one hand for the individual children, by alleviating their suffering in providing more effective medications against their symptoms and constrain negative effects on sleep, growth and school performance on a day-to-day basis. In a perfect scenario, the prevention of allergies in children with high propensity of doing so would also serve this objective. On the other hand, also the society as a whole would benefit from results of these studies, as sick-leave days with economic loss could be reduced and the halting of the allergic march in childhood would confer health and socioeconomic benefits not only in childhood, but possibly also in adulthood. Other advantages of performing these studies include the knowledge gained by performing these studies in children in particular. Alternative strategies could be to study the above effects in for instance appropriate animal models or even adult humans. However, as children are not simply small adults, the collected information from these studies may be invaluable for the development of safe and directed treatment options for children. Furthermore, as the allergic march begins in childhood, and the foundations thereof originate from alterations in immune maturation throughout perinatal life, this provides the window of opportunity for allergy preventive strategies. Indeed, efficacious allergy prevention interventions are supplemented both pre-and postnatally²⁰, indicating the importance of both pregnancy and early postnatal life for immunomodulation. Hence, this provides arguments against studying allergy development from an immune maturation perspective in adults. The performance of studies on animals, which comes with yet other ethical issues, could possibly provide information about shared physiological mechanisms at work. On the other hand, the applicability of such findings to humans would regardless of age be limited, as there are substantial differences between even phylogenetically related species. Limitations of performing these studies include the risk of causing unintentional harm to the children by either the interventions themselves, or during the process of collecting samples at multiple occasions. However, as the studies are based on solid scientific evidence, and they are further performed with the support of highly competent and caring clinical personnel, mindful of the well-being of these children, I believe that the benefits of performing these studies outweigh the risks of partaking in them.

For all the four projects, ethical permission has been applied for and approved by the local ethics committees, with reference to the respective ethical permission diary numbers in the project descriptions above.

Methodological foundations

In this thesis, several experimental methods have provided the framework for the presented results. Table I briefly summarises the used methods and in which project they were implemented. Below, brief theoretical aspects of each of the utilised methods are also described.

Enzyme-linked immunosorbent assay (ELISA)

Enzyme-linked immunosorbent assay (ELISA) is a standard immunoassay for the relative quantification of proteins, such as cytokines, chemokines or antibodies in different matrices.^{302,303} There are several variations to this method, and in Papers I and IV, the quantification of the cytokine IL-10 and chemokine CCL18 (Table I) in allergen-stimulated PBMC supernatants and plasma, respectively, were carried out using the in-house direct sandwich ELISA approach. In Paper IV, CCL18 was measured using ELISA as the concentrations in the circulation are markedly higher than that of other cytokines and chemokines, making it difficult to incorporate within the dynamic range of the multiplex bead assay with other mediators. In Paper IV, IL-10 was measured in allergen-stimulated supernatants using ELISA, to ascertain the early production by monocytes after 24 hours. Later production of the same mediator after 6 days of incubation was investigated using a multiplex bead assay approach. In the ELISA technology the initial step consists of binding a capture antibody to a solid phase, in this case a polystyrene ELISA plate. Upon addition of the chosen sample possibly containing the antigen of interest for measurement, a biotinylated detection antibody is added, thereby completing the antibody-antigen sandwich. Thereafter, the addition of a streptavidin-conjugated horse radish peroxidase enables the enzymatic conversion of a substrate into a colorimetric signal. By means of a standard curve, consisting of a dilution series of known protein concentrations, the colorimetric signal is translated into a concentration, which corresponds to the content of the antigen in the sample. In Paper IV, the detection of total IgA in saliva was performed in a similar fashion, although the antigens of interest in this assay were monomeric or dimeric IgA-antibodies. The indirect in-house ELISA approach was also used to detect secretory IgA-antibodies. A capture antibody capturing the secretory component (in this case a monoclonal anti-human secretory component antibody) was bound to the plate, followed by the same steps as above. This hence results in the measurement IgA-antibodies which are dimeric, as they are connected by the secretory component.

Multiplex bead assay (Luminex)

Whereas the ELISA methodology enables the measurements of single analytes, the multiplex bead assay or Luminex methodology offers the opportunity of measuring several analytes at once in the same volume of sample at a higher dynamic range and sensitivity compared to ELISA.^{302,303} In Papers I and IV, in-house Luminex assays were setup for the measurements of several cytokines and chemokines in both plasma and allergen-stimulated PBMC supernatants (Table I). In these Luminex assays, capture antibodies are coupled to beads containing a known proportion of red and infrared dyes, which makes them distinguishable from each other. These beads are then suspended in the sample of interest, thereby allowing for several bead sets with distinct

capture antibodies to bind their specific antigens in the matrix. Upon addition of a biotinylated detection antibody, with subsequent addition of a streptavidin-fluorophore complex, the measurement of the amount of antigens in the samples is facilitated by the detection of a fluorometric signal. Yet again, these signals are translated into a concentration of the antigens upon relating them to standard curves of the respective antigens. In Paper I, a 4-plex in-house Luminex assay measured plasma levels of CCL17, CCL22, CXCL10 and CXCL11, whereas a 6-plex in-house Luminex assay was used for the detection of IFN- γ , IL-5, IL-10, IL-13, CCL17 and CXCL10 in allergen-stimulated PBMC supernatants in Paper IV. Additionally, in Paper IV the quantification of timothy grass pollen-component Phl p 1-specific IgA-antibodies in saliva was performed as a single plex assay. In this case, the Phl p 1 allergen component was coupled to the beads, enabling an indirect approach for measurement of IgA-antibodies specific for this component.

Flow cytometry

By adding fluorophore-conjugated antibodies specific for antigens such as CD4 and CD45RA to CD4⁺ cells, as was done in papers II and III, the expression of these surface markers may be studied using one of the most common methods in immunology: flow cytometry.³⁰⁴ Simply described, the converging of the provided sample into a steady stream of single cells in a so-called flow cell enables the flow cytometer to measure several parameters. By measuring the scatter of visible light, flow cytometry enables the investigation of features such as size (forward scatter) and granularity (side scatter). Additional excitation of fluorophores bound to antigen-specific antibodies by laser light allows for the analysis of both surface bound and intracellular proteins through the emitted signals. Upon data collection, these signals are analysed by means of a process known as gating, in which positive and negative cell populations are distinguished from each other. Ultimately, statistics of the expression of the antigen of interest in the cell or on the cell surface are obtained, which may be used for further studies.

DNA methylation analyses - Th subset locus-specific and epigenome-wide approaches

In this thesis, features of the epigenome in CD4⁺ cells upon pre- and postnatal intervention with probiotics and/or ω -3 fatty acids were examined by surveying one of the most stable and widely studied epigenetic modifications: DNA methylation.⁷⁹ DNA methylation is mostly studied in CpG sites, and in this context consists of the presence of a methyl (-CH₃) group on the fifth carbon of the cytosine residue. The effect of this modification is context dependent, but DNA methylation mainly confers transcriptional silencing of the locus in which the CpG site is situated.

In Paper II, the DNA methylation patterns were studied both on a local and global scale. Using a Th subset locus-specific method²⁹⁴, the DNA methylation patterns of four CpGs in promoters of Th cell-related loci were examined. These CpGs are located upstream of the *IFNG*, *IL13*, *IL17* and *FOXP3* promoters, and the methylation status of these loci reliably predict the activation of the loci as well as the polarisation of naïve Th cells into the respective subsets (Th1, Th2, Th17 and Tregs, respectively). The procedure consists of several steps. From isolated CMBC or PBMC, CD4⁺ cells were isolated, from which DNA was subsequently extracted.

The DNA was then bisulfite converted to allow for distinction between methylated (unaltered cytosines) and unmethylated (turned into the nucleotide base uracil) cytosines. The subsequent locus-specific PCR reaction then translates the uracil residues into thymines. This is followed by a methylation sensitive single nucleotide polymorphism primer extension step, in which unlabelled primers bind next to the studied CpG sites, and extend the nucleotide sequences with fluorescently labelled dideoxynucleotides. As each locus exhibits a unique length of the primers, the methylation status of each locus may be identified specifically using capillary electrophoresis.

The epigenome-wide examination of the DNA methylation status of several hundred thousand CpG sites spread across the genome may be investigated using high-throughput DNA methylation analyses such as the Infinium platforms provided by the Illumina Corporation. In papers II and III, the Illumina 450K³⁰⁵ and the Illumina MethylationEPIC 850K²⁹⁶ arrays were utilised, respectively, to study DNA methylation patterns on a global scale. Both arrays implement the Illumina Bead Chip technology comprising two different methods for identification of unmethylated and methylated cytosines: the Infinium I and Infinium II assays.²⁹⁵ As described above, the DNA for study initially needs bisulfite conversion to enable the distinction between methylated and unmethylated sites. Thereafter, the two different technologies are applied on the bisulfite converted DNA. A common feature of the two Infinium assays is the use of beads coupled to long target-specific probes for the detection of the DNA methylation status at a particular CpG site. The Infinium I assay exhibits two probes for each interrogated locus, one specific for the unmethylated C's, which are read as T's after whole genome amplification of the bisulfite treated DNA. The other one is specific for the methylated condition, the C's that have been left unchanged upon bisulfite conversion. Specificity of the identification is achieved by matching the probes at their 3' ends to either of these two conditions. In contrast, the Infinium II assay holds only one bead type for the probe-specific identification of the DNA methylation status. In this case, the added complementary bases at the 3' end of the probe differentiate between the methylated sites, with G's corresponding to C's, and unmethylated sites, with A's corresponding to T's at the opposite strand. This allows for identification of a larger number of CpG sites in the epigenome, without making assumptions about neighbouring CpG sites, which is the case in the Infinium I assay. Regardless of assay, results from these analyses are presented as beta values, representing a mean of the methylation status (designated as 0 or 1 for the presence of methylation at any given CpG site) in the sample at hand. In the Illumina 450K array, more than 485 000 sites are examined, whereas over 866 000 sites are inspected using the Illumina MethylationEPIC 850K approach.²⁹⁶ The latter includes more than 90% of the sites included in the Illumina 450K array³⁰⁶, with additional information on enhancer regions. Prior to high-end statistical analyses, pre-processing of these large amounts of data include the correction of the use of the two Infinium assays along with various quality control measures (described in detail below).

Allergen stimulation of peripheral blood mononuclear cells

In allergic individuals, the immune response is often skewed towards the Th2-response along with diminished Th1- and Treg-responses. A means to study this immune imbalance in a functional manner is to isolate PBMCs from allergic individuals and stimulate this mixture of monocytes and lymphocytes with allergens and thereafter determine the secretion of cytokines and chemokines in the supernatants. This was done in Paper IV, where isolated PBMCs were stimulated *in vitro* with birch (*B. verrucosa*) and timothy grass (*P. pratense*) allergens, as well as

a positive control (phytohaemagglutinin), for 24 hours and 6 days, respectively. The stimulations were performed in serum-free medium, as this enhances the sensitivity of detecting antigen-specific immune responses.^{307,308} In the supernatants, IL-10 was measured from the 24-hour time point using ELISA as described above, whereas IFN- γ , IL-5, IL-10, IL-13, CCL17 and CXCL10 were measured after 6 days using a 6-plex in-house Luminex assay. This enabled the study of early monocyte-derived responses, as well as later mainly T lymphocyte-derived responses.

Table I. A tabular overview of the utilised methods in the papers included in this thesis.

Methods, sample types and analytes		Paper I (MAAS)	Paper II (<i>L. reuter</i>)	Paper III (PROOM-3)	Paper IV (GAP)
CBMC/PBMC isolation			X	X	X
CD4 magnetic bead isolation			X	X	
PBMC allergen stimulation (24h and 6 days with birch, grass and PHA)					X
In-house ELISA					
Plasma	CCL18	X			X
Allergen-stimulated PBMC supernatant	IL-10				X
Saliva supernatant	Total IgA				X
Saliva supernatant	Secretory IgA				X
In-house Luminex					
Saliva supernatant	1-plex (<i>Phleum pratense</i> -specific IgA)				X
Plasma	4-plex (CCL17, CCL22, CXCL10, CXCL11)	X			X
Allergen-stimulated PBMC supernatant	6-plex (IFN- γ , IL-5, IL-10, IL-13, CCL17 and CXCL10)				X
DNA extraction			X	X	
Flow cytometry			X	X	
T helper subset locus-specific DNA methylation analysis			X		
Illumina 450K array			X		
Illumina 850K array				X	

Methodological considerations

Throughout the years of working with the four described projects, technological advancements were made in the fields of study used in this thesis. For instance, the use of the immunological workhorses ELISA and Luminex has now increasingly been replaced by state-of-the-art methods such as mesoscale discovery (MSD) and proximity extension assay (PEA). Being an electrochemiluminescent, plate-bound, sandwich immunoassay similar to ELISA, MSD still possesses the multiplexing capabilities of the Luminex methodology, albeit with comparable higher sensitivity and less time consuming performance.^{309,310} PEA takes this a step further, as minute amounts of sample are needed for proteomic studies of circulating mediators.³¹¹ By means

of amplifying signals of oligonucleotide-labelled antibodies binding to the target proteins of interest, either using qPCR or next-generation sequencing approaches, the method circumvents issues of cross-reactivity in multiplexed immunoassays, thereby providing higher specificity. The previous cost-effectiveness of the in-house ELISA and Luminex assays is nowadays sometimes marginal, for which reason similar analyses to those performed in this thesis may in future become, at least partly, replaced by MSD and PEA. The in-house technologies are not obsolete, however, as few if any commercial ready-to-use kits for either ELISA or Luminex analyses in saliva are available for ascertaining levels of *e.g.* allergen-specific IgA-antibodies. The advantages of measuring *e.g.* salivary allergen-specific IgA using a 1-plex Luminex (as described in Paper IV), include the possibility to immobilise the allergen on the bead surface, instead of on a plate, thereby increasing the surface area for antibody-antigen interactions. Furthermore, as the beads are in solution, this presumably facilitates easier binding of the antibodies to the bead-bound allergen, which may enable the otherwise challenging measurement of these mediators in saliva.

In the epigenetic field, similar changes have taken place in methodological area. Our primary epigenetic studies in Paper II consisted of studying the DNA methylation status of four Th cell-related loci using a method first described by Janson *et al.*²⁹⁴ The initial idea was to perform these analyses in papers III as well, but with time the high-throughput technologies surveying hundreds of thousands of sites all over the epigenome entered the field, providing a more comprehensive view of the DNA methylome. Illumina's 450K array was employed in paper II, and should have been performed in Paper III as well, if it were not for the introduction of the MethylationEPIC 850K array shortly thereafter. Hence, with time the analyses rapidly became much broader, while the methodologies still complement each other, as all the investigated CpGs are not included in any single one of the methods. The implementation of the advanced statistical and bioinformatic methods downstream of epigenome-wide screening of CpGs was initially a very time-consuming and complex task, as the field was new and unexplored. However, as more knowledge was gained and shared in the scientific community over time, the performance of the analyses has become easier to handle, but still requires skilled bioinformaticians conducting the work. In the present setting, other methods such as whole genome-wide bisulfite sequencing could have provided an alternative for the above conducted analyses. In Papers II and III, validation on the protein or transcription level are missing. In Paper II, this was due to the lack of available samples for those types of analyses. From the PBMC samples used in Paper III, RNA has been collected for later analyses using RNA sequencing, but were not considered part of the included manuscript due to time limitations. However, merging of the epigenetic and transcriptomic data using a multiomics approach has been considered for future analyses.

Statistical and bioinformatic analyses

In this thesis, a large variation of statistical methods has been implemented to identify, ascertain, and verify the findings of the four included papers. Brief descriptions of the utilised methods will follow, and a comprehensive overview of the applied methods within each project is given in Table II. Initially, demographic comparisons were performed in all four papers to inspect inherent differences in the study populations that could possibly confound downstream findings. Utilised tests for these analyses were Chi-square tests or Fisher's exact test for categorical variables with sample sizes above and below five observations, respectively. Continuous variables were surveyed using the parametric Student's t-test. To determine whether to proceed with parametric

or non-parametric univariate statistical analyses, the Shapiro-Wilk's or D'Agostino-Pearson tests of normality were carried out. Cross-sectional comparisons of distributions between two groups were carried out by using the non-parametric Mann-Whitney U test. Similar comparisons between more than two groups were performed either with a one-way ANOVA for parametric data, or Kruskal-Wallis tests for non-parametric data, followed by appropriate *post hoc* tests. Longitudinal comparisons over time were performed using the non-parametric Friedman test with Dunn's multiple comparisons test. To study correlations between variables, the non-parametric Spearman's rank test was applied. The predictive abilities of particular variables on outcomes of interest were explored using binomial logistic regression models. The resulting coefficients of these analyses provide an odds ratio of the respective outcomes per log-unit increase in the studied variable. GEE models are used for the longitudinal study of population-averaged effects of a variable of interest on a particular outcome over time.

DNA methylation analyses comprise a series of analysis steps including pre-processing, followed by high-level statistical analyses and accompanying validation analyses. In this thesis, the analyses were mainly performed within the R programming environment, which is excellent for handling large data matrices. Firstly, the raw DNA methylation data were pre-processed, including a filtering step, which removed CpG probes that were non-significant, multi-hit probes, non-CpG probes or overlapping with single nucleotide polymorphisms. Thereafter, the sex chromosomes were either removed, or retained, depending on the distribution of the sexes in the sample groups. Thereafter, a normalisation procedure, known as the beta mixture quantile (BMIQ) method, was carried out to adjust for the two different implemented Infinium assays used in the Illumina DNA methylation arrays. To study and potentially adjust for differences in cell proportions within the utilised isolated CD4+ samples, the Houseman method³¹² was used. In the case cord blood samples were studied, the samples were tested for maternal contamination by inspecting 10 CpG sites according to Morin *et al.*²⁹⁷ If threshold values of methylation were exceeded for at least half of these sites, the samples were considered contaminated by maternal cells. Singular value decomposition (SVD) analyses were carried through to determine significant components of variations within the data, that would need correction prior to proceeding with high-level analyses.

The performance of multidimensional scaling (MDS) analyses allows for unbiased investigation and visualisation of inherent differences with the now cleaned data sets, which may indicate differences due to *e.g.* intervention groups. Thereafter, analyses of differentially methylation within these data sets were performed, by fitting linear models to the data using the limma package.³¹³ By comparing treatment groups to the placebo group, these linear models provide mean methylation difference (MMD) and p-values for each probe. The p-values of are corrected for multiple comparisons using the Benjamini-Hochberg false discovery rate (FDR) procedure. To ascertain that the differential methylation analyses have not resulted in inflated significances among the tested probes, tests for genomic inflation using the BACON R package³¹⁴ were performed. Other validation approaches included bootstrapping and permutation analyses. All in all, chosen thresholds of MMD and FDR-corrected p-values provide the basis for the definition of DMPs and indirectly also DMGs. Pathway enrichment analyses using the Gene Ontology and/or KEGG databases were performed for the DMGs, providing initial information about pathways in which the genes are involved. However, as genes often act in concert with a multitude of other genes in their biological context, network analyses provide insights into the interaction of the DMGs with other genes. Thereby it is possible to establish the possible biological importance of DNA methylation changes induced by the investigated interventions.

Table II. A tabular summary of implemented statistical and bioinformatic methods in the thesis.

Statistical approach	Paper I (MAAS)	Paper II (<i>L. reuteri</i>)	Paper III (PROOM-3)	Paper IV (GAP)
Demographic comparisons				
Chi-square test (normal or Pearson)		X	X	
Fisher's exact test	X	X		X
Student's t-test	X			X
Assessments of data distribution				
Shapiro-Wilk's test of normality	X		X	
Levene's test of equal variances			X	
D'Agostino-Pearson omnibus normality test		X		X
Univariate statistical methods				
Parametric (>2 groups): One-way ANOVA (incl. <i>post hoc</i> test)		X		
Non-parametric (2 groups): Mann Whitney <i>U</i> test	X			X
Non-parametric (>2 groups): Kruskal-Wallis test (incl. <i>post hoc</i> test)	X	X	X	
Non-parametric: Friedman's test (incl. Dunn's <i>post hoc</i> test)				X
Correlation analyses				
Non-parametric: Spearman's rank correlation	X			X
Multivariate statistical methods				
Binomial logistic regression	X			
Generalised estimating equation (GEE) model	X			
DNA methylation quality control				
Pre-processing (filtering incl. removal of non-CpG, multihit, SNP-related and low quality probes)		X	X	
Batch correction		X	X	
Sex chromosomes retained (Yes/No)		Yes	No	
BMIQ normalisation		X	X	
Cell type composition analysis		X	X	
Test of maternal blood contamination		X	X	
Singular value decomposition (SVD)		X	X	
Differential DNA methylation analyses				
Multidimensional scaling (MDS)		X	X	
Linear model (moderated t-test, limma)		X	X	
FDR correction (Benjamini-Hochberg)		X	X	
Test for genomic inflation (BACON)		X	X	
Permutation and bootstrapping		X		
Network analysis				
Module identification (MODifierR)		X	X	
Pathway enrichment analysis		X	X	
Classifier construction		X		

In this thesis, intervention modules were created using four module identifying methods (Clique SuM, DIAMOnD, MCODE and Module Discoverer) from the MODifierR package.³¹⁵ Genes within the networks were either defined as nodes within their modules if they were identified by at least two methods (Paper II) or all four methods (Paper III). Hereafter, pathway enrichment analyses were performed again, for entire networks and clusters thereof, to give clues about the biological significance of the findings. In Paper II, we also performed statistical analyses in relation to development of allergic disease, and further established relevance for allergy through the development of a classifier.

Results and discussion

Paper I

Circulating chemokine levels in infancy and childhood predict outcomes of allergic disease later in life

Chemokines are important regulators of immune cell recruitment³⁶, and a number of chemokines have been related to the development of allergic diseases, such as CCL17, CCL18, CCL22, CXCL10 and CXCL11.^{113-120,124,125} The study of these chemokines in the natural course of allergy development has, however, not been previously performed in a larger population of children. Hence, the circulating levels of the above-mentioned allergy-related chemokines were measured in blood samples from children included in an observational birth cohort.²⁸⁹ Plasma concentrations of the chemokines from the time of birth (cord blood) and at 1 and 8 years of age were related to development of sensitisation and asthma at ages 1, 3, 5, 8, 11 and 16 years. Furthermore, as allergic manifestations such as asthma and sensitisation are heterogeneous in nature, the role of chemokines in allergic disease heterogeneity was examined, by relating the chemokines to previously machine learning-derived allergy clusters.

Childhood CXCL10 and CCL18 levels influence present and future development of sensitisation

Sensitisation, which in essence describes the propensity of an individual to produce IgE-antibodies, constitutes the prerequisite of developing allergic symptoms caused by IgE-mediated mast cell degranulation.³ As we previously have associated chemokines to development of sensitisation^{113,124,125}, corresponding comparisons were performed in Paper I. Levels of the Th1-associated chemokines CXCL10 and CXCL11 increased throughout childhood (Paper I: Figure 1 D-E), in accordance with previous observations.^{113,124,125} Relating the chemokine levels to the development of sensitisation, decreased levels of CXCL10 and CXCL11 at birth and/or 1 year of age associated with development of sensitisation later in childhood (summary in Figure 10A, Paper I: Figure 3A-B, Figure S3A-B). This is in line with findings from a previous study, in which children later becoming sensitised revealed lower levels of CXCL11 at birth and age 2 years.¹¹³ An inverse relationship between high levels of the same chemokine and sensitisation was also observed¹¹³, in keeping with the predicted lower odds of developing sensitisation at age 11 if the levels of CXCL10 were high at age 8 (Figure 10B, OR=0.4, 95% CI 0.2-0.8, p=0.012). This suggests that impaired Th1-responses could contribute to the propensity of becoming sensitised in susceptible individuals.

Regarding trajectories of the Th2-associated chemokines, the levels of CCL17 decreased from birth throughout childhood, whereas the opposite was true for CCL22 (Paper I: Figure 1A-B). The augmented concentrations of CCL17 in cord blood associated to being sensitised at age 8 (Figure 10A, Paper I: Figure S3D), but no significances were observed for CCL22. None of the Th2-associated chemokines demonstrated further relationship in predicting sensitisation. This finding was rather unexpected, as earlier studies have shown that increased cord blood levels of CCL22 predicted sensitisation, whereas enhanced CCL17 concentrations predicted development of allergic symptoms.^{124,125} Possibly, this could be attributed to the observed increase in the levels of CCL22 throughout childhood, which contrasts to previous results from our group.^{113,124,125} Elevated CCL22 levels have been demonstrated in house dust mite-sensitised

Figure 10. Associations and predictive value of chemokines to development of sensitisation. Panel A provides a graphical summary of findings from the associative analyses of the circulating chemokine levels at birth (in cord blood), age 1 and age 8 years, to the development of sensitisation at ages 1, 3, 5, 8, 11 and 16 years. Displayed significant findings originate from comparisons between sensitised and non-sensitised individuals (Mann-Whitney *U* test, $p < 0.05$). Panel B displays a forest plot visualising the probabilities of developing sensitisation, derived using logistic regression models adjusted for sex, parental atopy and parental smoking. Odds ratios are denoted with crosses and the corresponding values below them, and 95% confidence intervals are illustrated by the error bars. All displayed models have an adjusted p -value of < 0.05 .

children at birth until 1.5 years of age¹²⁷, and whereas house dust mite sensitisation is uncommon in Sweden, it poses the main allergen towards which children are sensitised in the UK.³¹⁶ Substantial differences in sample size, as well as differences in prevalence of sensitisation could, however, also confound this result. The Th2/Treg-induced chemokine CCL18 showed consistently elevated levels at 1 and/or 8 years of age, and associated both with present sensitisation as well as development at multiple time points later in childhood and adolescence (Figure 10A, Paper I: Figure 3C-E, Figure S3C). Interestingly, the odds of becoming sensitised were elevated at ages 8 (OR=3.3, 95% CI 1.5-7.1, $p=0.002$), 11 (OR=3.0, 95% CI 1.3-6.7, $p=0.008$) and 16 years (OR=4.2, 95% CI 1.7-10.5, $p=0.002$) if the concentrations of CCL18 were increased at age 8 (Figure 10B). Furthermore, when evaluating the role of the chemokines in allergic disease heterogeneity, higher CCL18 levels at age 8 tended to associate with clusters related to development of sensitisation towards multiple allergens compared to sensitisation to few or no allergens (Paper I: Figure 6A-B).

Neonatal CXCL11 and childhood CCL18 levels predict different outcomes in terms of asthma development

Asthma is a chronic respiratory inflammatory disease, presenting with both remodelling and hyperresponsiveness of the airways³¹⁷, and manifesting with characteristic symptoms such as

recurrent episodes of exacerbations including wheeze, chest tightness as well as breathing difficulties.³¹⁸ In previous studies by us¹¹³ and others¹¹⁴⁻¹¹⁹, elevated levels of the Th1-associated chemokines have been associated with development of wheeze and asthma. In Paper I, we could corroborate these results for CXCL10, as increased circulating concentrations at ages 1 and 8 years associated with later development of asthma (Figure 11A, Paper I: Figure 2A-B, Figure S2A) However, unlike the findings for CXCL10, decreased CXCL11 levels at birth associated with asthma development at age 16 years, and elevated neonatal levels seemed protective on the outcome (Figure 11B, Paper I: Figure S2B, OR=0.4, 95% CI 0.2-0.8, p=0.012). Hence, there seems to be differences in temporal effects of these chemokines, despite the fact that both are induced by IFN- γ .¹⁰⁷ Possibly, the higher affinity of CXCL11 in relation to CXCL10 to the shared receptor CXCR3, and its superior chemotactic capabilities, could explain this fact.³¹⁹

Figure 11. Associations and predictive value of chemokines to development of asthma. Panel A provides a graphical summary of findings from the associative analyses of the circulating chemokine levels at birth (in cord blood), age 1 and age 8 years, to the development of asthma at ages 1, 3, 5, 8, 11 and 16 years. Displayed significant findings originate from comparisons between asthmatic and non-asthmatic individuals (Mann-Whitney U test, $p < 0.05$). Panel B displays a forest plot visualising the probabilities of developing asthma, derived using logistic regression models adjusted for sex, parental atopy and parental smoking. Odds ratios are denoted with crosses and the corresponding values below them, and 95% confidence intervals are illustrated by the error bars. All displayed models have an adjusted p -value of < 0.05 .

Another possibility is that the diminished Th1-response this represents could contribute to dysregulation in immune responses early in life, amplifying the typical Th2-skewing of the newborn. Thereby this could pave way for allergic disease, and induction of Th1-responses later in childhood could be compensatory in nature, to outweigh Th2-skewing. Furthermore, exacerbations caused by viral respiratory infections are common among asthmatic children³²⁰, possibly also explaining the induction of these Th1-associated chemokines.

No significant findings could be reported for the Th2-associated chemokines, but yet again elevated systemic levels of the dually regulated chemokine CCL18 at 8 years of age associated with development of asthma at ages 8 and 16 years (Figure 11A, Paper I: Figure 2 C-D). The probability of developing asthma at ages 8 (OR=3.8, 95% CI 1.5-9.8, $p=0.006$) and 16 years

(OR=5.3, 95% CI 1.7-16.1, $p=0.003$) also increased substantially with presence of high concentrations of CCL18 at age 8 (Figure 11B). These findings match previous results, showing elevated levels in recurrent wheeze¹¹³ and asthma^{133,134}, and are further supported by the association of elevated CCL18 concentrations at age 8 in children presenting with asthma exacerbations compared to non-wheezers (Paper I: Figure 6D).

Infant and childhood levels of CCL18 predict long-term development of sensitisation and asthma

As some of the chemokines both associated with development of sensitisation and asthma, and predicted the development of both outcomes later in life, we proceeded with GEE models to assess whether these findings were also solid in longitudinal analyses. All chemokines at each age were included in the models, and no consistent longitudinal patterns were evident for any of the chemokines at any age, except for CCL18 at ages 1 and 8 years (Figure 12). In relation to sensitisation, the longitudinal models included SPT results from five time points throughout childhood (ages 3, 5, 8, 11 and 16 years). We chose to exclude age 1 data, as the prevalence of sensitisation at that age was generally lower than later in life, and the sample size was limited compared to the other ages. The odds of becoming sensitised in the time span from 3-16 years of age increased with every unit increase in CCL18 levels at age 1 (OR=3.1, 95% CI 1.2-8.3, $p=0.022$). Similarly, examinations of the effects on later childhood exclusively (ages 8, 11 and 16 years), showed that elevated circulating CCL18 levels at age 1 (OR=3.5, 95% CI 1.3-9.9, $p=0.018$) and age 8 (OR=3.0, 95% CI 1.5-5.9, $p=0.002$) increased the odds of becoming sensitised by more than threefold. In a similar manner, longitudinal effects of the chemokines were performed in relation to subsequent asthma development at ages 5-16 years (four time points) and 8-16 years (three time points). We excluded development of pre-school wheeze as a surrogate for asthma development, hence motivating our use of the 5-16 years interval. In line with the previous findings in Paper I, augmented CCL18 levels at age 1 year posed higher odds of developing asthma from early school age through adolescence (OR=2.9, 95% CI 1.1-7.6, $p=0.028$). Likewise, elevated CCL18 levels at ages 1 (OR=3.5, 95% CI 1.3-9.8, $p=0.01$) and 8 years (OR=3.0, 95% CI 1.6-5.9, $p=0.001$) associated with higher odds of developing asthma between 8 and 16 years of age.

CCL18 is constitutively expressed in several tissues, including by alveolar MOs in the lung, and may influence Th cell differentiation by poising development towards immunoregulatory Treg subsets.¹²⁸ This indicates that CCL18 in the steady state supplies immunoregulatory functions. Indeed, the regulatory features of CCL18 is diminished in allergic individuals,^{138,139} and elevated levels have been observed in several allergic conditions such as AD^{136,137} and allergic rhinitis.¹³⁵ As CCL18 furthermore may stimulate collagen-production in the lung¹⁴⁰, a key event in remodelling of the lung in asthma development³¹⁷, this possibly explains the findings in relation to asthma in Paper I by us and others.^{133,134}

- Limitations of the analyses in Paper I include the lack of experimental evidence looking at functional relevance of the presented findings. Furthermore, the generalisability of the findings may be an issue, as also shown by discrepancies to previous studies, as the study population was recruited in a rather confined area rendering it fairly homogenous.

Figure 12. A forest plot displaying odds ratios from generalised estimating equation (GEE) models predicting longitudinal allergy development from the measured CCL18 levels. The predictive ability of the chemokine CCL18 on longitudinal development of asthma was assessed at ages 5-16 years and 8-16 years, whereas sensitisation was evaluated between 3-16 years and 8-16 years of age. Odds ratios are denoted with crosses and the corresponding values below them, and 95% confidence intervals are illustrated by the error bars. All displayed models showed an adjusted p-value of <0.05.

+ The strengths of Paper I include the substantial sample size as well as the performance of both cross-sectional and longitudinal analyses, which have not been performed for chemokines in this kind of setting previously.

In conclusion, the studies in Paper I revealed that elevated CCL18 levels predicted the development of asthma and sensitisation, with longitudinal effects of childhood concentrations on later allergy outcomes. The Th1-associated chemokines CXCL10 and CXCL11 levels also preceded later development of sensitisation and asthma, albeit with differential regulation at distinct time points. Altogether, this supports further studies on chemokines as putative biomarkers in allergy development.

Paper II

Prenatal intervention with *L. reuteri* modulates epigenetic landscape in cord blood CD4+ cells

In foetal life, maternal exposures originating from *e.g.* the diet and living environment influence the interpretation of the genetic setup of the developing child, hence preparing the foetus for postnatal life.³²¹ Epigenetic regulation makes up this link between the virtually static genome and ever-changing environmental exposures, thereby modifying and adapting the genome to appropriate interaction and reaction to its surroundings.⁷⁹ As epigenetic modifications such as DNA methylation regulate the differentiation of Th cells, and Th cells are key players in allergy development, this implies a role for epigenetic regulation in the development of allergies.^{79,83} Furthermore, as reduced diversity of microbial exposures is believed to underlie the allergy epidemic²⁷, probiotic supplementation has evolved as a means to increase diversity. However, the role of DNA methylation patterns in CD4+ Th cells early in life upon perinatal probiotic intervention, in the attempt to prevent childhood allergies, has not been well-established. Therefore, we in this study explored local and global DNA methylation patterns, in CD4+ Th cells from our pre- and postnatal probiotic intervention study.²⁴¹ Children included in this study provided samples at birth (cord blood) and ages 1 and 2 years, CD4+ Th cells were isolated from cord blood and peripheral blood samples, and were examined by means of two different approaches. Using a Th subset-specific DNA methylation analysis method, four CpGs in the promoters of the Th cell-related loci *IFNG*, *IL-13*, *IL-17* and *FOXP3* were investigated.²⁹⁴ Epigenome-wide DNA methylation analyses were performed by means of Illumina's Infinium 450K array³⁰⁵, and the epigenomic outcomes were for both methods investigated in relation to age, intervention group and development of allergic diseases.

Probiotic intervention leads to greatest epigenome-wide changes at the time of birth and affects immune-related pathways

Studies of whole epigenome-wide DNA methylation patterns upon prenatal treatment with probiotics throughout the last month of pregnancy, revealed intrinsic differences between treated and untreated children at the time of birth (Figure 13A), but not at the other time points (Paper II: Figure 3B-C). This was also reflected in the differential DNA methylation patterns, where the largest amount (1629) of DMPs (defined as MMD $>\pm 5\%$ and FDR <0.1) was observed (Figure 13B) in cord blood, while few or no DMPs were evident at 1 and 2 years of age (Paper II: Figure 3E-F, Tables S1, S3-4). This suggests that, in our setting, the probiotic intervention had a more prominent impact throughout foetal life than early postnatal life, which could be explained by the considerably higher antigenic load of exposures from other origins after birth compared to *in utero*. Moreover, this points at the importance of prenatal exposures, including possibly disease preventive interventions, for later immune maturation and later putative development of disease.^{33,223} Furthermore, investigating the direction of methylation of the 1629 DMPs, more than two thirds of the sites were hypomethylated, indicating transcriptional permissiveness of the sites. As the DNA methylation differences were the clearest at the time of birth, we proceeded with further analyses on these sites. Mapping the DMPs to their corresponding DMGs, 1015 genes containing at least one DMP were found, and most were hypomethylated (Paper II: Figure S5). Gene enrichment analyses showed involvement of the

genes affected by probiotic intervention in *e.g.* cell migration, cell signalling and TGF- β signalling (Paper II: Table S11). This shows that probiotic intervention leads to a general permissive transcriptional state in genes related to the immune-related pathways. Although DNA methylation generally is considered to convey transcriptional silencing, context dependent differences in other genomic regions are also apparent.⁸⁷ Either way, as we neither could confirm transcriptional activation on an mRNA or protein level, nor perform functional studies on the isolated CD4⁺ cells, further studies on the subject would have to clarify the relevance of these findings.

No significant findings could be revealed when inspecting possible differential DNA methylation patterns between children later developing allergic manifestations or not. In contrast, when examining separate treatment group comparisons of DNA methylation patterns throughout the first 2 years of life in relation to the time of birth, we found that shared genes between treatment groups were mainly hypermethylated and involved in processes such as IFN- γ signalling and T cell activation (Paper II: Figure S2B and S3A-B). This was corroborated by findings from the Th subset locus-specific DNA methylation analyses, where the *IFNG* promoter locus was most highly methylated at birth, and significantly less methylated at the other measured time points (Paper II: Figure 2A). These observations are in line with previous studies, revealing a lower degree of DNA methylation when comparing children^{322,323} and adults³²²⁻³²⁴ with neonates. Altogether, these findings presumably mirror normal immune development throughout the first year of life, as the Th2-skewed immune system of foetal life due to environmental exposures slowly but steadily is accompanied by viral-induced Th1-responses. No further differences were seen for comparisons of intervention groups or outcomes of allergy development in the Th subset-specific DNA methylation analyses.

Figure 13. Visualisation of MDS analysis and volcano plot of differential DNA methylation at birth. In panel A, an MDS plot illustrates inherent differences in the DNA methylation data (upon filtering, BMIQ-normalisation and cell-type correction). Red dots indicate probiotics, blue dots denote placebo. Panel B visualises results from the differential DNA methylation analyses in a volcano plot, based on the comparison between probiotics and placebo at birth. The differential methylation ($-\log_{10}$ p-values) is plotted against Mean $\log_2$ methylation difference (MMD) for each probe. The differential methylation analysis included fitting a linear model to the obtained data and FDR-correction for multiple testing. The horizontal dashed line represents a cut-off of FDR < 0.1 and the vertical dashed lines corresponds to MMD < -0.05 and > 0.05, respectively. Red dots indicate DMPs fulfilling both criteria for FDR and MMD described above.

Network analyses reveal relationships between DMGs, immunological pathways and allergy development upon probiotic intervention

As the identified DMGs above were related to a number of immune-related pathways, this encouraged us to further examine the role of these genes in interaction with other genes, using network analyses. By the identification of modules using four module discovery methods from the MODiER package³¹⁵, a consensus module was created consisting of genes predicted by at least two of the four methods. The resulting intervention module consisted of 91 genes with 361 protein-protein interactions (Figure 14A), of which most (n=53) originated from the differential DNA methylation analyses, and the rest from the interaction network (Paper II: Table S12). Genes enriched within the whole network were represented in pathways such as PI3K-signalling and leukocyte migration (Paper II: Figure 4B). Closer inspection of the structure of the network showed four clusters related to general cellular processes such as transcription and translation, but also TGF- β , PI3K-Akt and MAPK signalling as well as chemokine signalling (Figure 14A). Although we in the initial analyses could not show any specific differential methylation patterns between children who did and did not develop allergic manifestations, we were interested in whether the genes from the network were related to cumulative allergy development until age 7 years. By means of extracting DMPs from all the 91 genes within the module, including those extracted from the interaction network, 2734 DMPs were used for the comparison between children developing allergic manifestations compared to those who did not. Indeed, we could show that more genes were significantly different than expected by chance (Figure 14B, n = 61 genes, OR = 1.43, p = 2.4×10^{-6}), hence suggesting that the genes found in the intervention module were indeed related to allergy development as well. In line with the previous findings, most genes were yet again hypomethylated, and some genes also displayed mixed methylation patterns (Figure 14B). The classifier developed from these data also showed decent ability to discriminate between allergic and healthy children (AUC = 0.78, p = 3.0×10^{-3} , Paper II: Figure 5B), providing solidity to these findings.

Examining the connectivity of genes within the network, a number of genes possessing more than 10 connections were considered highly connected (Paper II: Table S12). Of these genes, five had a MMD of at least 5%, and were also previously related to development of allergic disease (Paper II: Table S13). *PRKCA* and *ITGA4* were both hypermethylated, while *CCR10*, *OXGR1* and *GNAO1* were hypomethylated. *PRKCA*, the gene coding for protein kinase C alpha, was most highly differentially hypermethylated among these genes. It has previously been related to asthma development^{325,326}, and allergen-stimulated PBMC IL-5 levels at 1 year of age have been observed to correlate with the expression of protein kinase C alpha at birth.³²⁷ Similarly, the also hypermethylated gene *ITGA4*, which codes for integrin subunit alpha 4, is differentially expressed in children developing asthma.³²⁸ The heterodimeric complex of integrin $\alpha 4$ with the integrin $\beta 7$ subunit is highly important for the homing mechanisms of imprinted T cells in re-circulating back to the gut mucosa.³²⁹ As the probiotic intervention most probably executes its function in the gut mucosa, and the expression of $\alpha 4\beta 7$ is elevated in allergic subjects³³⁰, this is of notable interest for our study. Another protein of importance in the homing of T cells is the CC chemokine receptor 10 (CCR10), which upon binding of epidermally produced CCL27 recruits T cells to the skin.³³¹ In Paper II, this gene was hypomethylated, and interestingly, in individuals with AD and asthma the expression of CCR10 on circulating CD4+ cells was increased.³³² In the light of our findings, this is particularly intriguing, as active probiotic treatment led to less development of IgE-associated eczema among children in this cohort²⁴¹ The

Figure 14. Graphical representations of the intervention module, and the relation of the intervention module to allergy development. Panel A illustrates the module identified based on genes containing significant DMPs (MMD $>\pm 5\%$, FDR <0.1) from the differential DNA methylation comparison between the probiotic and placebo groups at birth. Circumscribed yellow areas display intra-connected subclusters within the network, as examined by bioinformatic gene enrichment analyses for pathways. Panel B shows the same module, where genes containing probes that are significantly differentially methylated, in the comparison between children that do and do not develop allergic manifestations until age 7 years ($p < 0.05$), are highlighted from the background. Blue denotes the presence of only hypo-methylated probes, red indicates the presence of only hyper-methylated probes, and colouring with both blue and red indicates presence of both hypo- and hyper-methylated probes. Nodes represent genes and connecting lines represent protein-protein interactions (representing a STRING combined score > 0.7).

two other hypomethylated genes, the G-protein-related mediators *OXGR1* and *GNAOI*, have been implicated in allergy development in mouse models³³³ and humans³³⁴, respectively.

- Limitations of the analyses in Paper II include the small sample size, which may have impacted on the presented results, as well as the inability to perform longitudinal analyses in this data set. However, to compensate for this drawback, we have by means of several statistical approaches ensured the robustness of our data. This included assessing genomic inflation of the differential DNA methylation analyses, further bootstrapping and permutation analyses to confirm the resulting DMPs and the criterion of the DMGs in the networks to be present in at least two of the four module-identifying methods. As our findings are in line with what others have previously found^{93,94,97,335}, this suggests that the statistical approaches were statistically sound and scientifically relevant. Furthermore, the lack of data on actual transcriptional activation provides another limitation for these analyses, as the functional relevance of the differential DNA methylation could not be evaluated.

+ Strengths of Paper II include the performance of two different types of DNA methylation analyses, which complement each other in the knowledge they produce. Furthermore, performing these analyses in purified CD4+ cells, which are crucial in development of allergic disease, provides another advantage of our study. The samples collected in this double-blind randomised placebo-controlled trial are also unique, supporting the credibility and importance of studying probiotic intervention in foetal life and infancy. Another strength of the study consists of the performance of network analyses, which provide clues to the interactions between DMGs and other genes in a wider biological context.

In conclusion, the findings of Paper II show that the epigenome was poised towards enhanced immune maturation and activation at the time of birth in CD4+ T cells from children supplemented with prenatal L. reuteri. Hence, this suggests that probiotic intervention may provide the foetal immune system with a head start.

Paper III

Differential epigenomic effects of combined prenatal intervention with *L. reuteri* and ω -3 fatty acids in cord blood CD4+ cells

During pregnancy dietary exposures provide not only nutrients for the developing foetus, but also important cues for immune maturation and precursors for mediators involved in inflammation. The essential ω -3 and ω -6 fatty acids constitute parent fatty acids giving rise to inflammation resolving SPMs such as resolvins and maresins in the former case, and pro-inflammatory mediators such as prostaglandins and leukotrienes in the latter.²⁵³ In the present day context, there is a dominance of ω -6 fatty acids in our diets, hence driving the production of mediators maintaining an allergic inflammatory state. This imbalance may, however, be altered through the supplementation with ω -3 fatty acids modulating the inflammatory response towards resolution.²¹⁸ As an extension of the studies in Paper II, Paper III encompasses epigenetic studies in children enrolled within the PROOM-3 trial. This study investigates the role of pre-and postnatal combined probiotic and ω -3 fatty acid supplementation on allergy development, and whether there may be synergistic effects of combining approaches to stimulate immune responses and modulate inflammation. Investigations of the epigenome-wide DNA methylation patterns of this combined approach have not previously been performed in the setting of neonatal CD4+ T cells. In the PROOM-3 study, four treatment arms were studied using the Illumina Infinium MethylationEPIC 850K array²⁹⁶: double active treatment (*L. reuteri* and ω -3 fatty acids, L ω), single active treatments (active treatment for either *L. reuteri* (LP) or ω -3 fatty acids (P ω), placebo for the other) and double placebo treatment (PP).

Combined *L. reuteri* and ω -3 fatty acid treatment leads to greatest differential methylation effects

Initial MDS analyses of the DNA methylation data showed clear differences between the treated and untreated groups (Figure 15A). The greatest effects on differential methylation were seen in the combined treatment group (L ω) compared to the double placebo group (PP), where 1659 DMPs (Paper III: Table II) were identified as significant (Figure 15B, MMD $> \pm 5\%$, FDR < 0.1). Of these DMPs, a majority were hypermethylated (Paper III: Table S5). These DMPs were subsequently mapped to 794 genes, which were mainly involved in processes such as Phospholipase D signalling, PI3K-Akt and MAPK signalling as well as TCR signalling (Paper III: Figure S6A). As in Paper II, further network analyses were pursued to examine the role of the DMGs in putative interactions with other genes. In contrast to Paper II, the genes identified within the modules in Paper III were identified by all four chosen module discovery methods from the MODifier package.³¹⁵ In this way, the number of DMGs were narrowed down to 170 genes (Paper III: Table S5), which actually all derived from the differential methylation analyses. Of these genes, six genes had more than 10 connections within the network, and were considered highly connected, namely: *FYN*, *GNB5*, *SNAP23*, *LAMC1*, *IL6* and *RAF1*. Of these genes, all but *IL6* were hypermethylated. IL-6 is a cytokine responsible for inducing both innate and adaptive inflammatory responses, and it modulates Th cell effector function by activation of pathways such as MAPK, JAK-STAT and PI3K signalling.³³⁶ Interestingly, in asthmatic adolescents, hypomethylation of *IL6* associated with decreased lung function³³⁷ and in a perinatal probiotic intervention trial, the *IL6R* gene was hypomethylated in the first six to

Figure 15. Visualisation of MDS analysis and differential DNA methylation analysis for the double active (L ω) treatment group in comparison to the double placebo group (PP). In Panel A, red dots represent the active treatment group, while blue dots represent the placebo group. Panel B illustrates the distribution of DMPs in a volcano plot. DMPs are defined as having a MMD of $>\pm 5\%$ (vertical dotted lines) along with an FDR-corrected p-value of <0.1 (horizontal dotted line) and are depicted as red dots.

twelve months of life.³³⁸ Although we have neither assessed the development of allergic disease in these children yet, nor evaluated transcriptional relevance of our findings using other methods, it is tempting to speculate that treated neonates become equipped with inflammatory abilities, that may outweigh remnants of foetal Th2-skewing. Another gene of interest was *FYN*, which was hypermethylated in the combined treatment group. This non-receptor tyrosine-protein kinase acts downstream of the TCR, with key functions in T cell activation and differentiation³³⁹, indicating that the treatment may inhibit these events. The implication of this is, however, not certain. The genes within the network formed several subclusters, involved in processes such as antigen processing and presentation, TCR signalling and TGF- β signalling (Figure 16).

Single treatment with *L. reuteri* or ω -3 fatty acids show fewer but interesting findings

In line with the findings from the L ω group, the single active probiotic (LP) and ω -3 fatty acid (P ω) groups showed inherent differences in the DNA methylation data (Figures 17A and 19A). Similarly, substantial numbers of DMPs were evident for the LP (1246) and P ω (984) groups (Figure 17B and Figure 19B, Paper III: Table II). Mapping these DMPs to genes, 620 and 494 genes were discovered for the LP and P ω groups, respectively. As in the L ω group, the majority of the DMPs, and consequently DMGs, were hypermethylated in both single treated groups (Paper III: Table II, Table S6-S7). Subsequent module identification resulted in a condensation of the number of genes in the networks to 100 and 87 genes for the LP and P ω groups, respectively. While no genes had more than ten connections within the P ω module (Figure 20), two genes coding for adenylyl cyclases (*ADCY7* and *ADCY9*) were considered highly connected within the LP network (Figure 18).

Figure 16. Graphical representation of the consensus module created with the DMGs from the differential methylation comparison between the double treated *L.reuteri*/w-3 (Lw) and the double placebo (PP) group. Genes are represented as nodes with protein-protein interactions (STRING combined score > 0.7) illustrated as connecting lines within the network. Red nodes illustrate hypermethylated genes, blue nodes indicate hypomethylated genes and both colours denote mixed methylation patterns. Highly connected genes (>10 connections) are designated as nodes with bold black lines. Biologically relevant clusters identified by bioinformatic pathway enrichment analyses are enclosed by black lines.

Shared genes between networks reveal common pathways, while unique genes provide insights into treatment-specific effects

As the respective networks for LP (Figure 18) and Pw (Figure 20) displayed subclusters found within the Lw network, such as antigen presentation and signalling pathways downstream of the TCR, further investigations into shared and unique network genes and pathways were pursued. Ten genes were shared between all treatment groups (Paper III: Figure 6A, Table S8), being involved in general cell signalling pathways such as extracellular matrix–receptor interactions and cellular adhesion. While the same number of genes were shared between the single treated LP and Pw groups, no pathways could be revealed for this comparison. 24 genes involved in Ca²⁺ signalling and O-glycan biosynthesis were common between the Lw and LP comparisons, whereas the 14 genes shared between the Lw and Pw groups related mainly to TCR signalling and downstream pathways. This suggests that genes that are affected by all treatment constellations involve pathways which are generally important for T cell signalling and function. However, the direction of the methylation plays a role as well in the possible implication of these

findings. In essence, the direction of methylation is stable in a majority of the DMGs when comparing the L ω group to the single treated groups (LP, P ω) and the single treated groups with each other (Paper III: Figure 7). For DMGs shared between all comparisons, about half the common genes retain the same DNA methylation status, while the rest display different patterns depending on the treatment group. Hence, in general most genes common between all or some comparisons maintain identical DNA methylation patterns, suggesting treatment common effects on the epigenome.

Investigating the unique genes for each comparison, indications for synergistic effects of the double treated L ω group appears. While the number of genes between the LP and P ω group were similar (56 vs. 53 genes), the L ω group revealed a striking 122 unique genes (Paper III: Figure 6B). The argument for synergy of the double treatment could furthermore also be supported by the number of DMPs and DMGs extracted from the differential methylation analyses prior to network module creation, and the shared genes and pathways between the comparisons. The P ω comparison showed no relevant enriched pathways, whereas the L ω and LP groups, despite the uniqueness of the genes for the different comparisons, showed enrichment for similar pathways involved in TCR signalling, Th cell differentiation, antigen processing and presentation as well as asthma. As we presently do not have any information on allergy development, future studies will have to show whether these differential methylation changes associate to the development of *e.g.* asthma.

The most striking finding of Paper III was the general hypermethylation upon treatment with any constellation of perinatal interventions. This contrasts to findings from other studies mainly showing differential hypomethylation upon perinatal *L. rhamnosus GG/Bifidobacterium lactis* (in obesity-related genes)³³⁸, and ω -3 fatty acid treatment³⁴⁰, while yet others have revealed no differential DNA methylation upon prenatal fish oil supplementation.³⁴¹ These differences could originate from the different investigated cell types (whole blood at 6–12 months³³⁸, cord blood leukocytes³⁴⁰ and cord blood CD4+ cells³⁴¹), as well as methodological and statistical differences.

Figure 17. Visualisation of MDS analysis and differential DNA methylation analysis for the single active probiotic (LP) treatment group in comparison to the double placebo group (PP). In Panel A, red dots represent the active treatment group, while blue dots represent the placebo group. Panel B illustrates the distribution of DMPs in a volcano plot. DMPs are defined as having a MMD of $>\pm 5\%$ (vertical dotted lines) along with an FDR-corrected p-value of <0.1 (horizontal dotted line) and are depicted as red dots.

Figure 18. Graphical representation of the consensus module created with the DMGs from the differential methylation comparison between the single treated *L.reuteri* (LP) and the double placebo (PP) group. Genes are represented as nodes with protein-protein interactions (STRING combined score > 0.7) illustrated as connecting lines within the network. Red nodes illustrate hypermethylated genes, blue nodes indicate hypomethylated genes. Highly connected genes (>10 connections) are designated a nodes with bold black lines. Biologically relevant clusters identified by bioinformatic pathway enrichment analyses are enclosed by black lines.

More interestingly, this stands in opposition to the general hypomethylation revealed upon prenatal probiotic intervention in Paper II, upon which the stated hypothesis of Paper III was founded. However, several apparent differences between the two papers could explain this discrepancy. Firstly, the treatment was initiated at an earlier stage in the studied cohort in Paper III (GW 20) compared to previous pre- and postnatal studies on separate *L. reuteri* (GW 36)²⁴¹ and ω -3 fatty acids (GW 25).^{265,266} Furthermore, the maternal dose of the probiotic intervention was increased by an order of magnitude (10^9 compared to 10^8 CFU previously²⁴¹), which possibly could potentiate prenatal effects of the *L. reuteri* intervention *in utero*. The platform used for the epigenome-wide DNA methylation analyses also differs between the papers, as the 450K array was used in Paper II and the MethylationEPIC 850K array was employed in Paper III. Whereas about 485 000 sites were studied in Paper II, more than 866 000 sites were evaluated in Paper III, rendering the data sets considerably different in size from the start. Furthermore,

Figure 19. Visualisation of MDS analysis and differential DNA methylation analysis for the single active ω -3 fatty acid (P ω) treatment group in comparison to the double placebo group (PP). In Panel A, red dots represent the active treatment group, while blue dots represent the placebo group. Panel B illustrates the distribution of DMPs in a volcano plot. DMPs are defined as having a MMD of $>\pm 5\%$ (vertical dotted lines) along with an FDR-corrected p-value of <0.1 (horizontal dotted line) and are depicted as red dots.

various statistical adjustments, as for instance the exclusion of sex chromosomes in Paper III, may also have impacted on differences in the outcomes between the papers.

- Limitations to the studies in Paper III include the absence of additional experimental data on transcriptional activity, which would have provided valuable validation on the possible influence of the respective interventions on transcriptional regulation and gene expression. These analyses are in pipe-line for the project, but were excluded from the work presented in this thesis due to time restrictions. Another drawback of this work is presented by the lack of information on later allergy development in the children. This information is only available for some of the children as the study is presently on-going, and associations of allergy development to altered DNA methylation patterns will be evaluated in future analyses. Additionally, DNA methylation analyses on later time points in childhood, such as at ages 6 months, 1 and 2 years, could also come in question in up-coming studies.

+ Strengths of the studies in Paper III include the study of purified CD4⁺ cells, which are key players in the development of allergic inflammation. Furthermore, although the sample size was seemingly limited, we managed to get reasonable numbers in all four arms of this double-blind randomised placebo-controlled study, enabling us to study how the methylome reacts to the different intervention combinations in a unique material. The performance of network analyses to investigate gene-gene interactions also offers advantages in terms of the statistical power of the analyses.

Figure 20. Graphical representation of the consensus module created with the DMGs from the differential methylation comparison between the single treated ω -3 fatty acid (P_w) and the double placebo (PP) group. Genes are represented as nodes with protein-protein interactions (STRING combined score > 0.7) illustrated as connecting lines within the network. Red nodes illustrate hypermethylated genes, blue nodes indicate hypomethylated genes and both colours denote mixed methylation patterns. Biologically relevant clusters identified by bioinformatic pathway enrichment analyses are enclosed by black lines.

*In conclusion, the studies in Paper III revealed that regardless of treatment group, interventions induced general hypermethylation in DMPs. The pathways affected by the treatment with *L. reuteri* and/or ω -3 fatty acids were mainly involved in T cell signalling as well as other immune-related pathways. However, further studies evaluating gene expression and later allergy development will be important for the elucidation of the impact of the interventions in regulating Th cell function at the epigenomic level.*

Paper IV

Three years of SLIT alters immune responses towards tolerance in grass pollen-allergic children

Immunotherapy distinguishes itself from other treatments by not only alleviating symptoms of present allergic disease, but also by possibly inducing tolerance towards the allergen causing the symptoms, and hence preventing further allergic diseases from developing. As present ARC predisposes for the development of allergic asthma in the future^{280,282,283}, the general aim of the GAP study was to examine the possible asthma preventive effects of *P. pratense* SLIT in children with ARC induced by the same allergen.²⁸¹ Several *P. pratense* allergen components induce sensitisation in children, but Phl p 1 is often the initial³⁴² and main component towards which children with timothy grass pollen-induced ARC are sensitised.^{343,344} While studies of the immunomodulatory effects of SLIT in adults have revealed effects on both innate and adaptive immune responses^{345,346}, corresponding effects in children and adolescents are largely unexplored. Therefore, we in this study set out to determine the immunomodulatory effects of *P. pratense* SLIT in children with timothy grass pollen-induced ARC before (at inclusion), during (after three years with treatment) and after intervention (two years after the treatment was ended). The studies comprised both *in vitro* and *in vivo* measurements of cytokines and chemokines, to study possible alterations in immune responses. Effects on the local immune milieu in the oral cavity were also studied, by analysing salivary levels of total, secretory and Phl p 1-specific IgA-antibodies.

P. pratense SLIT modulates immune responses in the oral mucosa by inducing timothy grass pollen-specific IgA-antibodies in saliva

The main finding of this study was that the levels of Phl p 1-specific IgA-antibodies in saliva were significantly increased after three years with active SLIT treatment (Figure 21A), with sustained elevated levels two years after the treatment had ended. Although circulating levels of timothy grass pollen-specific IgA₁ and IgA₂ in sera have been observed upon SLIT treatment in adults³⁴⁵, corresponding elevations in saliva have previously not been shown in neither children²⁸⁴ nor adults.³⁴⁷ This makes it a unique finding, as it is the first time anyone has successfully measured grass pollen-specific IgA-antibodies in saliva, which poses certain challenges as a study matrix. Moreover, although other studies have shown elevations of house dust mite-specific salivary IgA-antibodies upon SCIT in adults³⁴⁸, this is the first study to show corresponding elevations of timothy grass pollen-specific IgA in saliva from children and adolescents upon SLIT treatment. Upon further examination of these level over time in children treated with GRAZAX®, a significant difference was observed between the time of inclusion and after three years of treatment, but not later (Figure 21B). Bearing in mind that the number of children included in these analyses were few, and individual trajectories may look different, this still poses interesting scenarios. Possibly, upon reaching elevated levels of the Phl p 1-specific IgA-antibodies after three years of treatment, these threshold levels were maintained after two years, even without daily exposure to the allergen. The uptake of the tablet in the mucosal immune environment of the oral cavity, which by default is a pro-tolerogenic environment, may have facilitated the induction of tolerance in several ways. The persisting long-term effect of grass pollen-specific IgA-production indicates that a part of the induced effector B cells from the treatment period have remained as memory B or plasma cells in *e.g.* the salivary glands.

Figure 21. Associations of salivary Phl p 1-specific IgA-antibody levels in saliva with treatment in children from the GAP study. The mucosal concentrations of timothy grass pollen-specific IgA-antibodies in saliva were investigated at inclusion, after three years of treatment and two years after treatment ending (five years from inclusion time point). Panel A illustrates the comparison between GRAZAX®-treated (grey circles) and placebo treated (white circles) children. The data are shown as medians with interquartile ranges. The numbers of studied individuals are indicated below each time point as n (P=Placebo/G=GRAZAX®). * $p < 0.05$, ** $p < 0.01$, Mann-Whitney U test. In Panel B, the longitudinal association of salivary Phl p 1-specific IgA-antibodies in the active treatment group is visualised over time. The total number of individuals in the groups at the respective time points are indicated below each time point. A Friedman test with Dunn's multiple comparisons test was performed, * $p < 0.05$.

Additionally, it is plausible that the seasonal exposures maintained the levels, while local tolerance had been reached at that point. No differences in the levels of salivary total IgA or secretory IgA could be found (Paper IV: Figure S2), which we attributed to the fact that mucosal IgA is directed to an immense plethora of antigens, which would not be highly affected by the allergen-specific levels of one single allergen.

Alterations in IgE-mediated mast cell degranulation and presence of circulating IgG₄ after SLIT treatment

The presence of high levels of allergen-specific IgG₄-antibodies is considered a reliable biomarker for induction of tolerance in the context of allergen-specific immunotherapy, as the levels normally are low in the circulation.³⁴⁹ Its competitive allergen binding ability, and lack of immune complex forming and complement activating capacities yields the IgG₄-antibodies immunomodulatory features in this context. Indeed, the induced and prolonged immune responses were also visible in the periphery, with elevated grass pollen-specific IgG₄ levels measured three and five years after the treatment started (Paper IV: Figure 1C), in line with findings from the main study.²⁸⁴ As class switch to IgG₄-production requires the presence of IL-10,³⁵⁰ this suggests that Bregs and Tregs in the periphery provide this immunomodulatory environment upon induction of tolerance. We did, however, not see any differences in the allergen induced levels of IL-10 between the treatment groups (Paper IV: Table S1). Possibly

this could be explained by the fact that IL-4 may also drive IgG₄ class-switching in B cells.³⁵¹ Although IL-10 addition to IL-4 supplemented B cells restored IgA- and IgG-synthesis, allergic individuals exhibited significantly lower Ig-production compared to healthy controls.³⁵² Hence, one may not exclude that over the course of SLIT, the contribution of inductive signals may originate from a changing proportion of the Th2- and Treg/Breg-responses, indicating that further studies would have to elaborate on the induction of these antibodies *in vivo*. Furthermore, it has been shown that saliva induces production of IL-10 by autologous isolated circulating monocytes³⁵³, suggesting that studies of cytokine secretion in saliva and its effects on *e.g.* PBMCs might have added insights into further local immunomodulating effects of SLIT.

In addition to these findings, decreased wheal diameters upon SPTs with the allergen were seen at three and five years after treatment initiation (Paper IV: Figure 1A), indicating less IgE-dependent mast cell degranulation by inhibition through *e.g.* the competition of allergen-specific IgG₄ for allergen binding.³⁴⁹ Surprisingly, these differences were not reflected in the circulating grass pollen-specific IgE levels (Paper IV: Figure 1B). This could be due to use of different allergen components in the measurements performed by us and the measurements performed by other investigators within the GAP study. Furthermore, increased levels of grass pollen-specific IgE over time may originate from molecular spreading processes, proceeding from mono- to oligosensitisation towards several grass pollen-components over time, of which the initiating Phl p 1 component then makes up a smaller proportion.³⁴² Interestingly, we observed a concomitant significant decrease in the SPT reactivity towards birch after three years with treatment. Although this was not the one of the primary outcomes of our study, the presence of these types of spill-over effects has not been observed thus far in adults.^{347,354} Speculatively, the immune systems of children and adolescents may be more dynamic and adaptable in response to the treatment.

SLIT re-balanced Th1- and Th2-responses, with maintained shift after treatment ending

A shift in the immune responses upon active treatment was also measurable *in vivo* in plasma, revealing augmented levels of the Th1-associated chemokine CXCL10 (Figure 22A) both after treatment with GRAZAX® for three years, and two years later. A similar pattern was seen for CXCL11, with a trend towards higher levels in treated compared to untreated subjects at the three-year time point (Paper IV: Figure 3B). None of the other measured allergy-related chemokines showed any significant differences, which was also the case for the cytokine or chemokine concentrations from *in vitro* allergen-stimulated PBMC supernatants (Paper IV: Table S1). However, the ratio between mediators originating from the Th1- and Th2-responses may be used as a proxy for changes in the respective immune responses. Indeed, the ratios of the Th2-associated cytokines IL-5 (Figure 22B) and IL-13 (Paper IV: Figure 3D) over the Th1-associated cytokine IFN- γ were decreased in children who had been treated for three years with GRAZAX®. This proposes, although no differences in the allergen induced levels of cytokines could be seen, that the immune responses in these individuals were more Th1-prone upon treatment, corroborating the finding from the chemokines. The longevity of the effect could, however, only be measured in chemokines, which could be due to the fact that the levels in supernatants were induced upon allergen challenge, whereas the levels of the chemokines in plasma were not. Of additional note, there was a strong inverse correlation after three years of treatment between the IL-5/IFN- γ ratio in the allergen-stimulated supernatants and the allergen-specific IgA levels in saliva (Paper IV: Figure 4), suggesting that reductions in systemic

Th2-responses and concomitant induction of local immunoregulatory responses are related. In line with findings from the entire GAP study²⁸⁴, decreases in the reported symptoms of allergic rhinitis were evident after three years with treatment as shown by significantly lower records of VAS and allergic rhinitis scores (ARS) (Paper IV: Figure S3). Similarly, significantly lower VAS scores were recorded two years after treatment end, but only when the score was reported daily for 14 days compared to the incident symptom evaluation during the last week before the visit (Paper IV: Figure S3A and S3C). No correlations between the clinical outcomes and the measured immunological markers could be shown. Taken together, apparent long-term clinical and immunological effects corroborates findings from adults treated with GRAZAX[®].^{346,347}

Figure 22. Effects on spontaneous chemokine and induced cytokine levels upon SLIT treatment. In Panel A, the plasma levels of the Th1-associated chemokines CXCL10 are shown. In Panel B, the induced levels of IL-5 and IFN- γ in supernatants from grass pollen-stimulated PBMCs are shown. All measurements were performed at inclusion, after three years of treatment and two years after treatment ending (five years from inclusion time point). White circles indicate placebo treatment, grey circles GRAZAX[®] treatment. The numbers of individuals analysed are indicated below each time point as n (P=Placebo/G=GRAZAX[®]). The data are illustrated as medians with interquartile ranges. * $p < 0.05$, ** $p < 0.01$, Mann-Whitney U test.

- Limitations of the study include the relatively small sample size, and the limited generalisability of the findings on this rather homogenous group of children from two study centres in Sweden. However, the latter may be only partly true, as the findings in Paper IV that were also investigated in the main study reflect the findings of the entire study.

+ Strengths of the analyses in Paper IV include the investigations of both local mucosal immune responses at the site of treatment uptake, as well as peripheral immune responses on a systemic scale. Furthermore, the analysis of clinical outcomes as well as innate, and both cellular and humoral adaptive immune responses, allows for a broader insight into mechanisms behind the development of tolerance in SLIT. The longitudinal analyses performed on several samples also provide another strength of these studies. Moreover, the analysis of samples originating from this first double-blind, randomised, placebo-controlled trial of SLIT in children and adolescents with grass pollen-induced ARC provides another advantage of the performed studies.

In conclusion, the studies in Paper IV revealed that a three-year treatment with timothy grass pollen SLIT induced tolerant immune responses in children with timothy grass pollen-induced ARC, partly in the long-term. Mucosal increases in Phl p 1-specific IgA-antibodies were accompanied by induction of systemic IgG₄ levels and circulating Th1-associated chemokines both at treatment end and two years later in treated children. Furthermore, reductions in SPT reactivity and symptom scores indicate clinically meaningful reductions in Th2-associated immune responses. These findings altogether propose that SLIT treatment indeed promoted tolerance with maintained partial long-term effects.

Concluding remarks and future perspectives

In this thesis, several aspects of immune maturation and modulation in allergic disease among children have been investigated, from prenatal life throughout childhood up until adolescence (Figure 23). This was done by examining different immune mediators in the natural course of allergic disease (Paper I) and in the modulation of future (Papers II-IV) and present (Paper IV) allergic manifestations. The mediators were studied in the circulation (Papers I and IV) and in saliva (Paper IV), as well as by examining epigenetic modifications of Th cells (Papers II-III).

In Paper I, the role of chemokines in the spontaneous development of allergic disease in childhood and adolescence was investigated. These studies revealed that elevated levels of the dually Th2/Treg-regulated chemokine CCL18 preceded the development of sensitisation and asthma, which also persisted longitudinally until early adulthood. While the finding is in line with that of several other studies investigating allergy development^{113,133-137}, none has previously shown long-term influence on allergic outcomes throughout this time frame. Furthermore, although a large number of immunological mediators have hitherto been studied as potential biomarkers for allergic disease in the MAAS cohort, none has previously shown this strong and consistent associations to allergy outcomes. This suggests that CCL18 should be further investigated in other cohorts to study its applicability as a predictor of allergies in other demographic populations. In Paper I, we also observed associations to both asthma and sensitisation for the levels of the Th1-associated chemokines CXCL10 and CXCL11, which despite the common upstream regulation of the two chemokines showed different temporal effects on the different outcomes. We thereby partly confirmed our stated hypothesis, with exception for the observed differences in the directionality of the Th1-associated chemokines. These results should be investigated more thoroughly in other materials, to disentangle the different roles these chemokines may have in allergy development, and particularly how viral-induced wheeze may modulate these chemokine levels in asthma. Also, whether the fluctuations over time in the chemokine levels are a cause or a consequence of allergic disease remain to be uncovered. Taken together, the collected findings in this study encourages further examinations of the role of chemokines as biomarkers for allergy development as well as in a clinical setting.

In Papers II and III, the effects of pre- and postnatal *L. reuteri* intervention alone or in combination with ω -3 fatty acids on the epigenome of CD4+ T cells were examined. As the greatest epigenome-wide effects of *L. reuteri* treatment were seen at the time of birth in Paper II, the studies in Paper III were likewise performed in cord blood. While the prenatal *L. reuteri* supplementation in Paper II lead to general hypomethylation, in line with the stated hypothesis, any combination of active treatments in Paper III gave rise to general hypermethylation when compared to the placebo groups, in opposition to the stated hypothesis. Despite these discrepancies, differential methylation in both papers affects T cell-related and immune-related pathways. Studies on the epigenetic effects of probiotic or ω -3 fatty acid intervention on childhood allergies are scarce. To our knowledge no-one has previously shown genome-wide differential DNA methylation upon probiotic treatment in childhood allergies in CD4+ cells, and while effects of prenatal ω -3 fatty acid treatment in another study showed changes in histone modifications of Th cells³⁵⁵, no differences could be seen on the DNA methylation level.³⁴¹ This points out the uniqueness of the findings in Papers II and III. The effects of the altered DNA methylation patterns as a result of any of the investigated interventions would, however, have to be ascertained on a transcriptional or expression level to draw any firm conclusions about the

biological significance of the findings. It would also be interesting to perform lipidomic analyses in the PROOM-3 cohort, and merge it with epigenetic and transcriptional data in a multiomics approach, to receive a broad perspective of interventional effects. Furthermore, in line with the hypothesis stated in Paper III, there seemed to be synergistic effects upon double active treatment with *L. reuteri* and ω -3 fatty acids, as reflected by the amount of differentially methylated sites and genes compared to any of the single treated groups. However, how this comes about mechanistically still needs further elucidation. Besides this, studies of the relevance of these findings in relation to allergy development still need addressing, and might provide further insights into how pre- and postnatal combined interventions may alter the course of allergy development. Altogether, the studies of Papers II and III show that prenatal intervention with probiotics, whether alone or along with ω -3 fatty acids, alters the DNA methylation patterns of cord blood CD4⁺ T cells, showing that maternal supplementation during pregnancy may affect immune development of the neonate.

In Paper IV, the immunological effects of *P. pratense* SLIT in children with timothy grass pollen allergy were studied. The treatment modulated both the mucosal oral immune response, as well as systemic immune responses towards a more tolerant state. In saliva, the levels of Phl p 1-specific IgA were elevated both at the end of treatment and two years later. These results were particularly intriguing, as it has previously from a methodological point of view been difficult to measure allergen-specific IgA-antibodies in saliva. Hence, this finding is unique in its own right, and validation of these findings in a larger cohort would provide insights into the applicability of this non-invasive measurement method in *e.g.* monitoring treatment success. The long-term effects in saliva were also mirrored in the circulation by augmented concentrations of grass pollen-specific IgG₄, a find well in line with previous research.^{284,356,357} Further effects in the circulation were represented by the high levels of the Th1-associated chemokines CXCL10 and CXCL11 upon treatment, which were maintained at these levels at follow-up. This suggests that the Th1-associated chemokines could play important roles in the creation and maintenance of tolerance. As these two chemokines also showed interesting findings in Paper I in allergy development, further studies should elaborate on the different inflammation promoting and immunomodulating roles these chemokines may have in these processes. Additionally, a skewing of the balance between the Th2/Th1-response towards the latter was observed in ratios of IL-5 and IL-13 over the IFN- γ levels. The collected findings of Paper IV were in line with the stated hypothesis, and showed induction of Th1-responses along with diminished Th2-responses, and elevated immunoregulatory responses. Taken together, this suggests that the SLIT treatment has induced tolerance, with partial persisting effects after treatment end, in treated children.

From where we stand right now in the field of allergy development and modulation, I believe that the future of allergy treatment and research investigations lies in combinations; the combination of different treatment regimens and the combination of several diagnostic tools for allergy development, to a much larger extent than has been previously the case. The realisation that allergic diseases are much more heterogeneous than previously believed provides explanation for the lack of treatment efficacy in some endotypes of allergic diseases.³⁵⁸ Additionally, as many of the treatments available today in clinical practice come with more or less troublesome side effects, the future of allergy treatments will, as treatments for other diseases alike, gravitate towards personalised medicine. In this context, it is possible that the findings in this thesis could be utilised in the quest for the optimal prediction, treatment and prevention of allergic diseases at different time points in the allergic march.

Figure 23. A summarising overview of the findings in this thesis.

Although the evidence to date for general allergy preventing effects of probiotics and ω -3 fatty acids are limited, the prevention of IgE-mediated eczema and allergic disease has repeatedly been shown in different studies.^{218,223} Future studies in relation to allergy development in the combined PROOM-3 intervention study will show whether these effects persist, and possibly are potentiated by each other, as well. From a wider perspective, in the case of parental history of allergies, it may then be possible to promote the perinatal intervention with probiotics and/or ω -3 fatty acids as a measure of possible immunomodulation away from allergic diseases. Prebiotics could also be implemented in this setting, as they may promote healthy commensal microbiomes.²⁴⁷ Speculatively, DNA methylation signatures derived from Th cells could in the future possibly be used as monitoring biomarkers for development of allergic disease, or even for treatment success. Implementing these strategies at the time of birth or early in childhood could also show how different environmental exposures (such as the supplementation with probiotics or ω -3 fatty acids) may modify the epigenetic patterns of inherited genetic predisposition. Thereby, it would be possible to give advice to patients on which particular exposures to avoid and which to promote.

Furthermore, circulating chemokine levels could also be interesting to monitor during infancy and childhood, as a way to predict development of allergic asthma and sensitisation. As previous studies showed predictive abilities for CCL17 and CCL22^{124,125}, and the work in this thesis showed the same for CCL18, CXCL10 and CXCL11, this convincingly points out the importance of these mediators in the allergy development and their potential as predictive immune biomarkers. Supposed this could be validated in another large cohort of children with allergic propensity, the measurement of all these allergy-related chemokines, along with maybe a Th17-related chemokine, could provide helpful knowledge about possible development of sensitisation and allergic manifestations. It would have been very interesting to measure the chemokines at several of the other time points as well in the MAAS cohort, and future studies should extend these measurements to early adulthood, maybe even in a double-blind randomised placebo-controlled allergy prevention trial, to disentangle the role of these mediators on allergy development later in life.

By the monitoring of chemokine or DNA methylation biomarkers throughout childhood, it would be possible to initiate appropriate treatments, even according to disease severity, at an early stage, thereby personalising the treatment while at the same time decreasing the risk for further allergy development. If allergic disease would still develop at some time during childhood and adolescence, immunotherapy might in the future be the primary long-term treatment choice, as desensitisation is a key factor on the route back towards tolerance. More so, this could be combined with *e.g.* biologics targeting Th2-mediators²⁷², probiotics and ω -3 fatty acids, as a means to promote immunomodulatory effects on different levels of the allergic inflammatory response. Furthermore, the combination of different allergen targets of immunotherapeutic treatments, either different components within the same type of allergen or combinations of different allergens, could provide possible wider allergen-preventive effects, as multi-sensitisation is common among allergic individuals. Studies using component-resolved diagnostics may in the coming years aid the choice of the right components to target, on a regional and individual specific level. Another interesting combination would be the merging of SLIT with SCIT intervention. For instance, with early SLIT treatment the foundations for oral tolerance towards the allergen may be laid, followed by SCIT treatment, which would maintain the treatment effects as seasonal boosters of pro-tolerant immune responses, or vice versa. Newer

immunotherapy approaches such as intralymphatic and epicutaneous immunotherapy may also be promising candidates for these combinations in a clinical setting.

Upon validating the findings on allergen-specific IgA-antibodies in saliva in a larger cohort, and at more time points throughout and after immunotherapy, it would be possible to use this as a marker of tolerance induction. By evaluating the levels of allergen-specific antibodies in saliva during the treatment period and at later follow-ups, it may be possible to discriminate between responders and non-responders at different stages of immunotherapy. If low levels are detected along with re-emerged presence of symptom, directed stimulation of tolerant immune responses could be commenced in non-responding patients at an early stage. This would not only be cost-effective in relation to treatment outcome, but also cause less suffering for children possibly being treated for many years with no benefits. The implementation of this method is feasible even during childhood, as it is reliable, and the collection of samples is non-invasive. Furthermore, the measurement method could also possibly be multiplexed, measuring different components within the same allergen, or different allergens as well, giving the most information about how IgA-responses change over time to seasonal and/or perennial allergens. Further interesting findings could have originated from studies of SLIT-treated saliva on *e.g.* circulating CD4+ cells and monocytes, to give further insights into how tolerance is induced by salivary mediators in the oral mucosa.

In conclusion, the four studies presented in this thesis have uncovered several perspectives of immune maturation and immune modulation in allergy development throughout childhood and adolescence. Circulating chemokine levels in childhood predict later allergy development. The immune system of allergic children, while poised towards allergic inflammation, is demonstrably dynamic in response to perinatal immunomodulating treatments such as probiotics and ω -3 fatty acids, and DNA methylation patterns reflect these changes. Likewise, SLIT treatment in childhood and adolescence modulates circulating and salivary immune responses, towards partial long-term tolerance. These collected results provide promising foundations to build further upon, in the quest for better understanding and treatments of allergic disease in childhood.

Epilogue

Never too late – a break in the atopic march?

The flames of the candle lights on the table flicker as several people enter the dining room to get seated for some Advent festivities. The atmosphere is vibrant – hugs, gifts and greetings replace each other at a fast pace. One by one, friends and family take their seat at the long table set with pepparkakor, glögg and lussekatter. It has been a while since I saw Linus and his family. He just recently turned 12 and has really grown from being a little boy to a youngster. A brief hello with a smile on his behalf at least shows that he is still not too cool to talk to his godmother. As everyone starts digging in on the treats, I ask his mother about how Linus is doing. After discussing school and his newly found love for padel, I cannot help, as a good friend and scientist, but ask her about his allergies. ‘His hay fever is a lot better, although he is not completely without symptoms’, his mother says. ‘so, the immunotherapy has helped to some extent. He finished the treatment about two years ago and gets through pretty well, with less medication than before. Although, some seasons it is still rough, depending on the pollen load of course’, she adds. ‘The same goes for his asthma, although he will probably have to accept living with that for the rest of his life.’ I nod in understanding, and tell her I am glad that he is doing better. I pour her another cup of glögg, and we join the intense conversations about the up-coming Christmas vacations.

The fire in the stove slowly dies out, and as I set to refill it with some new firewood, Linus comes and asks if he can help. ‘Of course’, I tell him, and hand him the brim-full basket that I just brought inside. Carefully, he refills the stove, and the fire slowly begins to light up again. We sit down in the couch in front of the stove, and I offer him some nuts from a bowl on the table. He lifts up his hand to decline. ‘I am still allergic’, he says, ‘and would not dare to try my luck although my allergies are better.’ With the sincere curiosity only found in children, he asks if what I do in my profession will ever make him completely free from his allergies. I tell him honestly that there are of course no guarantees. ‘But if we look at all the treatments you have had access to until today, like your antihistamines, the asthma medications and the immunotherapy, they all originated from work done by researchers. What might come in the future are treatments that are targeted to fit you as an individual better’. He nods in silence, waiting for me to elaborate. ‘Maybe some markers in your blood can be measured’, I continue, ‘to tell which types of allergies you might get later, or looking at the on and off switches of genes in the DNA we may better understand how this comes about. This could inform doctors, so that you get treatments that are more appropriate for your type of allergies. Do you follow?’ He nods again, reaching for some candy in another bowl on the table. ‘I also think we will see more treatments really early in life, like in pregnancy and shortly thereafter, in families with allergies. You know like the good bugs in yoghurts and the good fatty acids in salmon and mackerel? Then we could possibly prevent some allergies from developing.’ He looks genuinely surprised, and I briefly explain the background to him. I also tell him that I believe immunotherapies will be more common in allergy treatment, maybe in combination with the yoghurt bugs and fish fats. He interrupts me and asks excitedly, with his mouth full of candy, if he will ever be able to own any furry pets when he grows older. ‘It is possible that future treatments will be able to come around this issue, but only time will tell.’ The discussion ends abruptly as someone calls from the living room, asking whether anyone is up for board games. Me and Linus look at each other, as we answer in unison that we are coming. ‘Game on’, he says with a cheeky smile as we half-run to join the others.

Acknowledgments

On the origins of doctor Huoman – or the people thanks to whom she got there at all

As I am sitting down to write this part, I to my surprise find it the most difficult one of the whole thesis to write. Not because of the content itself – I could honestly write a book about the marvellous ways in which so many people have made life just extraordinarily worth living – but because I really feel so in awe over the immense support I have had. How do I express my gratitude to all those people who have supported me through thick and thin, and believed in me when I myself did not? If nothing else, this is an attempt at honouring all of you who have whole-heartedly supported me, in small and big deeds, throughout this journey. An homage to those who encouraged a very ordinary girl like me to fulfil the dream of becoming a researcher. I hope you all feel you have a share of my PhD. Because this is not a one-woman's work – this is about collaboration and mutual support, total interdependence with others. It is about learning from each other in work and in life, be it from family or friends, supervisors or students, colleagues or reviewers. And it has been one hell of a journey, where I have grown so much both personally and professionally. This thanks goes out to all of you, some mentioned, but non forgotten!

I feel compelled to start with those thanks to whom I exist in this very moment and place at all. A big great kiitos to my family, internally also known as the Finnish Mafia, who really are my Finnish juuret rooted into Swedish soil. *Mamma & pappa*, jag är oändligt tacksam för att ni alltid stöttat mig i min resa mot att bli forskare, från skolåldern hela vägen fram till idag. För alla värderingar ni lärt mig dela, för alla matlådor, bastubad och stugan, platsen på jorden som jag nog tycker bäst om. För att ni alltid står på min sida. Men kanske framförallt, tack så jättemycket för giving me my two siblings: *lillasyster Jessica* and *lillebror Jonathan*. Always on the same wavelength with jokes, you make me laugh from the heart, and you are the funniest people I know. *Jessi*, I cannot remember a life without you, you give so much of yourself to others, and your humour is the basis around which all our family gatherings revolve. The life of the party, the idéspruta for creativity, the longest hair and, since recently, also creative designer for PhD thesis covers (highly recommended!). *Joppe*, being the youngest of the three of us, I admire how you tune into your own way of doing things and learn things from scratch on your own, because you just put your mind to it. Also, you are the best training coach! You both inspire me so much, and I cannot express how much you both mean to me. You have also more recently extended the Finnish Mafia. It now also includes *Sofia G*, the one and only swagerska with the most impressive voice, both figuratively and physically speaking. I do not know anyone with a more solid belief in herself than you, and only you understand how awesome it is to sing loud to Queen Bey in the shower. And *Aleksander*, I so much appreciate the silent counterweight you add to our chatty family. Often less is more, and your calm ways are something I aspire to. And when you do say something, it is sprängstoff!

Thanks also to the whole Finnish mafia, for the family gatherings with my uncles and aunts *Jouni & Åsa* and *Jouko & Åsa*. Ni har alltid nyfiket frågat hur det går med forskningen och stöttat mig genom att ge mig möjlighet att göra och tänka på annat, som att pimpla, fika och umgås. Jag uppskattar det massor! Until recently, our *mummo & ukki* also took part in our gatherings. *Ukki* left the world before I started researching, and *minun kulta rakas mummo* sadly did not see me finish it off. I also wonder what *farmor & farfar*, whom I did not get to know, would have thought about this. Kaipaen teitä, *mummo & ukki*, och jag önskar att jag lärt känna er *farmor &*

farfar. I hope you feel my pride and joy of this achievement, which would not have been possible without you.

In den förlängda familjer har jag också haft turen att ha **farmor Ulla**. Tack för alla dina frågor om hur det går på labbet, mössen mår bara bra. När allt mest har känts som "Det går aldrig" på ren närkingska, har våra småresor och fikastunder gett mig perspektiv på vad som är viktigt i livet. Jag uppskattar våra telefonsamtal massor och tycker så mycket om dig! Och påminnelsen om att alla är tokiga, men att det uttrycker sig olika känns extra viktig nu. Till **Kalle & Marita**, ett innerligt tack för alla våra gemensamma utflykter och resor, fikor och middagar med spännande diskussioner om högt och lågt, nära och långt bort. Det är alltid lika trevligt och något att se fram emot när vardagen känts tung. Jag hoppas på många fler tillfällen framöver!

My own little family has a huge part in the making of this thesis. In order of importance, as we all know, the cat comes first. I would like to thank Putte, mattes urururururu, for putting up with me throughout the writing of this thesis. Literally no one has spent as much time with me during this time as you have, and no one has cared less about what I did while staring at that screen. Tack för att du alltid uppmuntrar mig med din totala närvaro, som får mig att släppa oron och istället tjänstgöra som första tjänstehjon i din stab. Allt gos, allt bus, all vila tillsammans när det varit tungt och svårt. Du är världens mest knasiga kisse, men också den mest älskvärda. Then comes, the husband. **Luky**, mitt hjärtas allt, mein Held, möj medved'. My wisest critic, my biggest supporter, my best friend in life to wake up next to every morning. I am convinced you are the better researcher of the two of us. Ohne dich hätte diese Reise wahrscheinlich nie begonnen, it would surely not have been as meaningful to go through, och jag hade definitivt inte klarat det utan dig vid min sida. Allt vi har gått igenom före och under min doktorandtid gör mig hoppfull om många år tillsammans framöver, även med en doktor i huset (som vi vill köpa snart!). Ja t'a veľmi ľubim, nekonečne veľmi. And thanks to you, I also have a Slovak rodina as well. Za túto moju slovenskú rodinu som veľmi vďačná. Chýbate mi **mamka, Lenka a Pa' o, Vladka a Maroš, spolu s Monikou a Melisou**. Teším sa na naše ďalšie stretnutie.

My first memories of the idea of becoming a researcher come from late högstadiet, when I at some point met **Paula Starbäck**. You taught biology and scientific subjects at gymnasiet in the neighbouring town, and after choosing that school, I got to learn some really awesome pre-university-lab-stuff. Your endless inspiration and passion for science in all shapes got me motivated to study at the university and pursue a PhD, and now you inspire me in seeing possibilities of working outside academia. Your impression made all the difference! Another important contributor to my scientific career choice comes from an evening course in basic immunology that I took in my Erasmus exchange year in Würzburg, Germany in 2009–2010. The lecturer, professor **Thomas Herrmann** introduced me to the field of immunology with a way of teaching that made even the most complicated processes seem extraordinarily fascinating and rather simple. Thinking back, this feeling still stays with me, and this was when I decided that I wanted to do research in immunology. Ich bin sehr dankbar dafür, dass ich den Abendkurs bei Dir belegt habe. Ohne dessen stünde ich ganz sicher heute nicht hier. Vielen Dank!

When coming back to Sweden from my exchange studies in Germany, I had decided upon two things: I will throughout my master studies work in a group with a female professor, and I will not pursue a PhD afterwards. Well, as it turns out, I liked working with **Maria Jenmalm** and her research so much that I ended up staying. **Maria**, tack för att du, med din enorma kunskap och ditt brinnande forskningsintresse inspirerade mig att vilja stanna i akademien trots allt. Din klokhet och styrka, ditt varma inkännande och din ousinliga tillit till min förmåga har fått mig att växa

som människa och forskare under doktorandtiden. I am forever grateful for letting me do things at my own pace, to find my way back to why I started with research in the first place. A big thanks also for choosing **Lennart Nilsson** as my co-supervisor. **Lennart**, you are the most down-to-earth person I know. Under sista halvan av doktorandtiden har vi lärt känna varandra bättre, och jag kan inte nog tacka dig för alla trevliga luncher, mysiga stugbesök och intressanta diskussioner vid middagar två timmar innan bjudning. Fantastiskt fint att få ha lärt känna din **Lena** också! Hoppas på många fler träffar framöver. I also owe **Maria** a great big thanks, for building up such a nice research group to work in, and I have had some amazing colleagues over the years. Thanks to previous and present colleagues of the Allergy group and the Allergy Center for all the nice meetings with summer and Christmas lunches. **Camilla Janefjord**, there is literally no one in the world I would rather spend a day, or a week, or more in the lab with than you! Tack för att du svarat på funderingar om labbet och livet när det behövts, och fyllt ut mina inkubationstider och pauser med idel roligheter. Och för att du är så fin kattvakt till Putte. I will miss you so much when I no longer work with you. **Anni Fornander**, tack för allt du lärde mig om labbet, och för att du tog ner mig på jorden som student. Jag kan fortfarande sakna ditt nynnande på labbet. Tack också till **Martina Abenius**, för att du hjälpte mig att lära mig stå för mig själv i början av doktorandtiden. Det är inte lätt att vara sig själv och agera i linje med sina värderingar när man vill anpassa sig till en ny roll. A warm thanks also goes to the students I have had the chance to work (along) with; **Annelie Castell**, **Carolina Gunhardson**, **Anna Pap**, **Simon Söderholm** and **Elin Olsson**. I learned so much about myself and the research we are doing by seeing it through your eyes, and growing with your motivation and skills. Also, I thank the participants of the studies and their parents for their participation, without which I would not have had anything to build my research on. I would also like to thank all co-authors on the papers for nice collaborations, especially **Mika Gustafsson** for sharing your expertise in advanced statistics and bioinformatics. I have also throughout my studies had the opportunity to visit **Adnan Custovic** both in Manchester and in London. Thank you for fruitful collaborations, and very nice study visits in the UK! A warm thanks also to **Sadia Haider**, for your patience in explaining statistics and very exhilarating discussions about the world in the big and the small things. Please, feel free to come visit some time!

Another reason I stayed in Linköping to pursue my PhD, was the welcoming and friendly atmosphere at AIR, and later at II. The fikaroom is a place of discussions of everything mellan himmel och jord, and I have had so many laughs with my colleagues. I am really grateful for everyone I have gotten to know there over the years, and would like to thank some in particular. **Karin Söderman**, dearest korsordsvän and kulturfaterworkare, thank you for being you! I admire you for standing your ground so solidly, and for reminding me that there is always another perspective to see. **Maria Tärnberg**, for being a great basgruppsbuddy with the students, and for being such a frisk fläkt in this conformative world. Both your soul and your clothes brighten up my day with bubbly joy. I miss having you around! **Petra Cassel**, thanks for sharing my love for liquorice, all your stories of cats (and their Instagram accounts) with me, and for being totally on my level when it comes to humour. A warm thanks also goes to **Jan Ernerudh** and **Christina Ekerfelt** for being so generous with your humour and good advice in life and science. A heartfelt thanks also goes to **Thomas Skogh**, flitens lampa lyser, fortfarande! **Jonas Wetterö**, for being such a funny and caring colleague, and for understanding what it means when things are så jävla gott! Good friends may also be found at work. **Linda**, **Liselotte** and **Daniela**, my TFD-ladies – thank you for sharing the love for good food and drinks, books and hiking. Our trip to Bristol was really amazing, and I still look for those cosy fleece pjs wherever I go. I miss you loads, and hope for a reunion soon! To **Georgia**, ευχαριστώ my dear friend, I will never forget your lovely

morning Helloooooo-hugs in the office. And I still look forward to learning more Greek. To *Johanna*, die bäste Kollegin vong Office her und 1 koole Freundin sowiso! Deine Struktur und dein Vermögen Sachen geschaffen zu bekommen sind beeindruckend, und wie du immer die gute Seite von Allem siehst inspiriert mich. Zusammen haben wir ja auch den tollsten *Stammtisch* gegründet: ein herzliches Dankeschön an *Annabel*, *Hansi* und *Lies*, der harte Kern, für all die schönen Momente zusammen. I would also like to thank *Valentina Bruno* and *Maria Svenvik*, for the sharing our office from time to time – you both lit up the everyday life of our lab with your presence. Grazie and Grüezi!

Friends are the family that you choose yourself, a good friend told me once. A large bunch of friends from all over the world have also made this journey both more joyful and interesting to take on. Thanks to my study buddies from the Uppsala times, when it all began. *Mona*, we have really followed each other's tracks over the years, and our conversations, whether live or digital always leave me with a smile on my face. *Cecilia*, a warm kiitos for the nice letters and conversations over the screen. Your laugh at the story about your mother's colleague on the Sweden ferry still brings me so much joy. And all my love to my Mädels aus den Würzburger Zeiten. *Maren*, *Uli*, *Joa* und *Steffi*. Es war einfach megaschön damals! Und auch wenn wir uns nicht so oft hören, finde ich es immer wieder beeindruckend, wie natürlich wir immer dort unsere Gespräche aufnehmen, wo wir zuletzt aufgehört haben. Ich vermisse euch sehr, und freue mich auf ein baldiges Wiedersehen. My gratitude also goes to my study mates from the master studies in Linköping. To *Pia*, my best friend and fellow Potter Head. I liked you so much already from the start, and without your eternal support I would not have allowed myself to fall when I needed to, and rise again when I was ready for it. Thank you for being the best cat-sitter, board game buddy and for sharing my love for puns and dad jokes. It is so nice to spend time with you, *Anders*, *Svante* and your recent family addition *Elsa*. I look forward to teaching her all I know about the world! *Petros* and *Amina*, thanks for all the good times during the studies, you made those two years some of the best in my life. I really miss talking to you guys! And *Karin Wählén*, you shine in your own pink glow! Can you imagine that we will have our dissertations two days apart?! It has been a crazy journey from our master studies until now. I am so proud of you, and feel happy to have had such a good friend by my side throughout these past few months. Hang in there, soon we will both be doctors!

As with many things in life, coincidence takes you many places and lets you meet many wonderful people. If it would not have been for the rain in the summer of 2012, I would never have met my Slovak family in Sweden, the *Ruskos*. *Katka & Martin*, u vás sa vždy cítim ako doma a veľmi si vás vážim. Naše spoločné stretnutia s vami znamenajú pre mňa aj pre Lukáša veľa. Keď ideme od vás domov, tak vždy s plným bruchom, veľkým úsmevom na perách a príjemným pocitom v srdci. And if it were not for you, we would not have gotten to know *Avdāns*, and shared all those great times together. *Nazlı & Tayfun*, we could not have wished for better neighbours in Lamböhov. I imagined this, and I did it! I miss you a lot, and hope to meet up soon again. And soon, *Martin and Tayfun*, *Luky* will also understand how it feels to have a doctor in the house! Another happy coincidence was that *Edith* moved into my room when I left the shared apartment I lived in in the beginning of my time in Linköping. That was the beginning of a lovely friendship, including many game nights with you and *Travis*, but also travels, good food and drinks, and conversations about the small and big things in life. And what a joy it is for me to be *Lukas's* godmother! I hope we can catch up on lost game nights soon, when I return from my research cave. And speaking of game nights, thanks to *Alva* and *Anders* for nice chats and introducing us to new games! Yet another wonderful coincidence, was that *Anthony & Joanitah*

whom I had gotten to know separately at university, actually were married! We enjoy so much spending time with you and the boys, and I hope to catch up soon! Tack till **Ahlam**, kära vän, för alla stunder med dig och din familj! Jag uppskattar vår vänskap jättemycket, och ser fram emot mer tid med dig framöver. A heartfelt thank you goes to **Johanna Lundin** and **Lina Jonsson**, who like me moved from our small town to bigger cities and explored the world for new perspectives and challenges. Your friendship means loads to me, knowing where we come from ("Sluta bjäla, börja gö"). **Johanna**, you are such an inspiration, totally self-made and competent, humble and kind at the same time! I admire your courage to follow your passion, and try new things. **Lina**, who would have thought we would end up working for the same employer, in the same institution so many years later? Without you, I would not have had any place to stay when I moved to Linköping. I enjoy our lunches a lot, and hope to hang around Friskis more with you soon! To **Amanda & Senad**, gracias and hvala for wonderful dinners, exciting trips and our long conversations about life. **Dotti & Krille**, wie toll, dass wir uns kennen gelernt haben! Wie ihr euren eigenen Weg findet, und ihn folgt, dass finde ich total schön und erfrischend, vor allem hier in Schweden. Ein riesiges Dankeschön geht auch an **Nele**. Es ist unglaublich wozu eine Brieffreundschaft führen kann. Auch wenn wir allermeist heute übers Telefon Kontakt haben, freue ich mich drüber immer wieder von dir zu hören. Du hast meine Reise von der Ferne mit grossem Interesse gefolgt, und deine Unterstützung bedeutet viel. Hoffentlich können wir uns bald wiedersehen, mit deiner kleinen Familie zusammen! **Linnea**, tack för ditt härliga nere-på-jorden-du, som du delar med dig av så generöst när vi ses. Vad kul vi haft på disputationer, middagar, luncher och promenader. Och vad klok du är, fina människa! An **Andrea & Roland**, für tolle Zeiten zusammen, für Katzenliebe, und für die Einsicht, dass es nicht nur einen Weg gibt, dass es auch anders geht. **Blažko & Julka**, mojí skevlí priatelja na Slovensku. Velmi vám ďakujem za pekné chvíle strávené na Slovensku a aj vo Švédsku. Mám vás veľmi rada a veľmi sa teším keď sa zase uvidíme, tentokrát aj s malou **Hankou**.

Things that have made life easier throughout these past years are also the spare time activities. A warm thanks to the book worms in the book club starring **Anna L**, **Anna K**, **Camilla J**, **Camilla S**, **Ammi**, **Kristina**, **Martina** and **Ingela**. It is just so enjoyable to eat, drink and discuss books with you. And our Christmas games are really the big happening every year. Then, I would like to swing my next thanks to **Agnes & Johan**, for unforgettable times in Lindy hop classes, especially in Herräng, but also for really nice after works and hikes. While dancing, we also got to know **Hannah & Stellan**. Thanks for nice dinners, for being such open-hearted people and for sharing what is yours with others! I would like to say Namasté to **Eva**, who introduced me to yoga and inspired me to look further within, to **Jenny** for teaching me to relax and let go, and to **Frida** for amazing yoga classes and reminding me that this too shall pass. And a sincere thank you to **Helena Yellow**, för din hjälp att hitta tillbaka till mig själv efter utbrändheten. För att du fått mig att lita på mitt sätt att göra saker på. För att du bit för bit fått mig att tro på mig själv igen och för sättet som du får mig att gå från våra samtal med ett lättare hjärta, en stärkt själ och en himla massa livsglädje. Du har gjort enorm skillnad för mig!

And finally, my warmest, biggest, most sincere tack, kiitos, danke, thank you, ďakujem goes to my past self. To **Hanna**, **mummon pikkunen Muru**, that young girl, aspiring at this big dream of becoming a scientist. Thank you, for allowing yourself to take on such a challenging journey, to grow beyond your own borders and for doing it your own way. I am so proud of you, for persisting when things got rough, for letting go when rest was needed and for keeping on doing the right thing all the way through. You did it, little one. I did it!

References

1. Asher MI, Montefort S, Björkstén B, *et al.* Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet (London, England)*. 2006;368:733-743.
2. Abrahamsson TR, Wu RY, Jenmalm MC. Gut microbiota and allergy: the importance of the pregnancy period. *Pediatr Res*. 77.
3. Thomsen SF. Epidemiology and natural history of atopic diseases. *European Clinical Respiratory Journal*. 2015;2:24642.
4. Dharmage SC, Perret JL, Custovic A. Epidemiology of Asthma in Children and Adults. *Frontiers in Pediatrics*. 2019;7:246.
5. Igea JM. The history of the idea of allergy. *Allergy*. 2013;68:966-973.
6. Galli SJ, Tsai M, Piliponsky AM. The development of allergic inflammation. *Nature*. 2008;454:445-454.
7. Bunyavanich S, Schadt EE. Systems biology of asthma and allergic diseases: A multiscale approach. *Journal of Allergy and Clinical Immunology*. 2015;135:31-42.
8. Ferreira MA, Vonk JM, Baurecht H, *et al.* Shared genetic origin of asthma, hay fever and eczema elucidates allergic disease biology. *Nature Genetics*. 2017;49:1752-1757.
9. Suaini NHA, Wang Y, Soriano VX, *et al.* Genetic determinants of paediatric food allergy: A systematic review. *Allergy*. 2019;74:1631-1648.
10. Moffatt MF, Gut IG, Demenais F, *et al.* A Large-Scale, Consortium-Based Genomewide Association Study of Asthma. *The New England Journal of Medicine*. 2010;363:1211-1221.
11. Martino DJ, Ashley S, Koplin J, *et al.* Genomewide association study of peanut allergy reproduces association with amino acid polymorphisms in HLA-DRB1. *Clinical & Experimental Allergy*. 2017;47:217-223.
12. Liang Y, Chang C, Lu Q. The Genetics and Epigenetics of Atopic Dermatitis—Filaggrin and Other Polymorphisms. *Clinical Reviews in Allergy & Immunology*. 2016;51:315-328.
13. Narożna B, Hoffmann A, Sobkowiak P, Schoneich N, Bręborowicz A, Szczepankiewicz A. Polymorphisms in the interleukin 4, interleukin 4 receptor and interleukin 13 genes and allergic phenotype: A case control study. *Advances in Medical Sciences*. 2016;61:40-45.
14. Pinto L, Stein R, Ribeiro J. Genetic associations with asthma and virus-induced wheezing: a systematic review. *Jornal Brasileiro de Pneumologia*. 2009;35:1220-1226.
15. Renz H, Skevaki C. Early life microbial exposures and allergy risks: opportunities for prevention. *Nature Reviews Immunology*. 2020:1-15.
16. Custovic A, Custovic D, Kljajic-Bukvic B, Fontanella S, Haider S. Atopic phenotypes and their implication in the atopic march. *Expert Review of Clinical Immunology*. 2020.
17. Oksel C, Custovic A. Development of allergic sensitization and its relevance to paediatric asthma. *Current Opinion in Allergy and Clinical Immunology*. 2018;18:109-116.
18. Selene BK, Zhou Z, Tao Z. The Atopic March: Progression from Atopic Dermatitis to Allergic Rhinitis and Asthma. *Journal of Clinical & Cellular Immunology*. 2014;05.
19. Paller AS, Spergel JM, Mina-Osorio P, Irvine AD. The Atopic March and Atopic Multimorbidity: Many Trajectories, Many Pathways. *Journal of Allergy and Clinical Immunology*. 2018;143:46-55.
20. Jenmalm MC, Duchén K. Timing of allergy-preventive and immunomodulatory dietary interventions - are prenatal, perinatal or postnatal strategies optimal? *Clin Exp Allergy*. 43.
21. Gerrard JW, Geddes CA, Reggin PL, Gerrard CD, Horne S. Serum IgE levels in white and metis communities in Saskatchewan. *Annals of allergy*. 1976;37:91-100.
22. Strachan DP. Hay fever, hygiene, and household size. *British Medical Journal*. 1989;299:1259.
23. Barker DJ. The fetal and infant origins of adult disease. *British Medical Journal*. 1990;301:1111.
24. Wadhwa PD, Buss C, Entringer S, Swanson JM. Developmental Origins of Health and Disease: Brief History of the Approach and Current Focus on Epigenetic Mechanisms. *Seminars in Reproductive Medicine*. 2009;27:358-368.
25. Rook GAW. Hygiene Hypothesis and Autoimmune Diseases. *Clinical Reviews in Allergy & Immunology*. 2012;42:5-15.
26. Hanski I, von Hertzen L, Fyhrquist N, *et al.* Environmental biodiversity, human microbiota, and allergy are interrelated. *Proceedings of the National Academy of Sciences*. 2012;109:8334-8339.

27. Lambrecht BN, Hammad H. The immunology of the allergy epidemic and the hygiene hypothesis. *Nature immunology*. 2017;18:1076-1083.
28. Blaser MJ. The theory of disappearing microbiota and the epidemics of chronic diseases. *Nature Reviews Immunology*. 2017;17:461-463.
29. Simon KA, Hollander GA, McMichael A. Evolution of the immune system in humans from infancy to old age. *Proceedings Biological sciences / The Royal Society*. 2015;282:20143085.
30. Gollwitzer ES, Marsland BJ. Impact of Early-Life Exposures on Immune Maturation and Susceptibility to Disease. *Trends in immunology*. 2015;36:684-696.
31. Bennett JM, Reeves G, Billman GE, Sturmborg JP. Inflammation–Nature's Way to Efficiently Respond to All Types of Challenges: Implications for Understanding and Managing “the Epidemic” of Chronic Diseases. *Frontiers in Medicine*. 2018;5:316.
32. Gusev YE, Zhuravleva YA, Zotova NV. Correlation of the Evolution of Immunity and Inflammation in Vertebrates. *Biology Bulletin Reviews*. 2019;9:358-372.
33. Jenmalm MC. The mother-offspring dyad: microbial transmission, immune interactions and allergy development. *Journal of internal medicine*. 2017;282:484-495.
34. Ashley NT WZaNR. Inflammation: Mechanisms, Costs, and Natural Variation. *Annual Review of Ecology, Evolution, and Systematics*. 2012; 43:385-406.
35. Bron PA, van Baarlen P, Kleerebezem M. Emerging molecular insights into the interaction between probiotics and the host intestinal mucosa. *Nature Reviews Microbiology*. 2012;10:66-78.
36. Commins SP, Borish L, Steinke JW. Immunologic messenger molecules: Cytokines, interferons, and chemokines. *Journal of Allergy and Clinical Immunology*. 2010;125.
37. Mjösberg J, Spits H. Human innate lymphoid cells. *Journal of Allergy and Clinical Immunology*. 2016;138:1265-1276.
38. Saravia J, Chapman NM, Chi H. Helper T cell differentiation. *Cellular & Molecular Immunology*. 2019;16:634-643.
39. Nguyen QP, Deng TZ, Witherden DA, Goldrath AW. Origins of CD4+ circulating and tissue-resident memory T-cells. *Immunology*. 2019;157:3-12.
40. Phan T, Tangye SG. Memory B cells: total recall. *Current Opinion in Immunology*. 2017;45:132-140.
41. Pieper K, Grimbacher B, Eibel H. B-cell biology and development. *Journal of Allergy and Clinical Immunology*. 2013;131:959-971.
42. Bournazos S, Ravetch JV. Diversification of IgG effector functions. *International Immunology*. 2017;29:303-310.
43. Novak N, Haberstick J, Bieber T, Allam J-PP. The immune privilege of the oral mucosa. *Trends in molecular medicine*. 2008;14:191-198.
44. Ortega-Gómez A, Perretti M, Soehnlein O. Resolution of inflammation: an integrated view. *EMBO Molecular Medicine*. 2013;5:661-674.
45. DuPage M, Bluestone JA. Harnessing the plasticity of CD4+ T cells to treat immune-mediated disease. *Nature Reviews Immunology*. 2016;16:149-163.
46. Pawlak M, Ho AW, Kuchroo VK. Cytokines and transcription factors in the differentiation of CD4+ T helper cell subsets and induction of tissue inflammation and autoimmunity. *Current Opinion in Immunology*. 2020;67:57-67.
47. Bhaumik S, Basu R. Cellular and Molecular Dynamics of Th17 Differentiation and its Developmental Plasticity in the Intestinal Immune Response. *Frontiers in Immunology*. 2017;8:254.
48. Palm NW, Rosenstein RK, Medzhitov R. Allergic host defences. *Nature*. 2012;484:465-472.
49. Rivas M, Chatila TA. Regulatory T cells in allergic diseases. *Journal of Allergy and Clinical Immunology*. 2016;138:639-652.
50. Hwang J-R, Byeon Y, Kim D, Park S-G. Recent insights of T cell receptor-mediated signaling pathways for T cell activation and development. *Experimental & molecular medicine*. 2020;52:750-761.
51. Brogdon JL, Leitenberg D, Bottomly K. The Potency of TCR Signaling Differentially Regulates NFATc/p Activity and Early IL-4 Transcription in Naive CD4+ T Cells. *The Journal of Immunology*. 2002;168:3825-3832.
52. Constant S, Pfeiffer C, Woodard A, Pasqualini T, Bottomly K. Extent of T cell receptor ligation can determine the functional differentiation of naive CD4+ T cells. *The Journal of Experimental Medicine*. 1995;182:1591-1596.

-
53. Zhu J. T Helper Cell Differentiation, Heterogeneity, and Plasticity. *Cold Spring Harbor Perspectives in Biology*. 2018;10.
 54. Hammad H, Lambrecht BN. Barrier Epithelial Cells and the Control of Type 2 Immunity. *Immunity*. 2015;43:29-40.
 55. Hong H, Liao S, Chen F, Yang Q, Wang DY. Role of IL-25, IL-33, and TSLP in triggering united airway diseases toward type 2 inflammation. *Allergy*. 2020.
 56. Gaffen SL, Jain R, Garg AV, Cua DJ. The IL-23–IL-17 immune axis: from mechanisms to therapeutic testing. *Nature Reviews Immunology*. 2014;14:585-600.
 57. Yasuda K, Takeuchi Y, Hirota K. The pathogenicity of Th17 cells in autoimmune diseases. *Seminars in Immunopathology*. 2019;41:283-297.
 58. Aujla SJ, Alcorn JF. TH17 cells in asthma and inflammation. *Biochimica et Biophysica Acta (BBA) - General Subjects*. 2011;1810:1066-1079.
 59. Palomares O, Akdis M, Martín-Fontecha M, Akdis CA. Mechanisms of immune regulation in allergic diseases: the role of regulatory T and B cells. *Immunological reviews*. 2017;278:219-236.
 60. Ansel MK, Lee DU, Rao A. An epigenetic view of helper T cell differentiation. *Nature Immunology*. 2003;4:616-623.
 61. Martín-Orozco E, Norte-Muñoz M, Martínez-García J. Regulatory T Cells in Allergy and Asthma. *Frontiers in pediatrics*. 2017;5:117.
 62. Jutel M, Akdis CA. T-cell Subset Regulation in Atopy. *Current Allergy and Asthma Reports*. 2011;11:139-145.
 63. Hellings PW, Steelant B. Epithelial barriers in allergy and asthma. *Journal of Allergy and Clinical Immunology*. 2020;145:1499-1509.
 64. Bulek K, Swaidani S, Aronica M, Li X. Epithelium: the interplay between innate and Th2 immunity. *Immunology and Cell Biology*. 2010;88:257-268.
 65. Krohn KI, Shikhagaie MM, Golebski K, *et al*. Emerging roles of innate lymphoid cells in inflammatory diseases: Clinical implications. *Allergy*. 2018;73:837-850.
 66. Walker JA, McKenzie ANJ. TH2 cell development and function. *Nature Reviews Immunology*. 2018;18:121-133.
 67. Kucuksezer UC, Ozdemir C, Cevhertas L, Ogulur I, Akdis M, Akdis CA. Mechanisms of allergen-specific immunotherapy and allergen tolerance. *Allergology International*. 2020.
 68. Komi D, Wöhrl S, Bielory L. Mast Cell Biology at Molecular Level: a Comprehensive Review. *Clinical Reviews in Allergy & Immunology*. 2020;58:342-365.
 69. Foster PS, Maltby S, Rosenberg HF, *et al*. Modeling TH2 responses and airway inflammation to understand fundamental mechanisms regulating the pathogenesis of asthma. *Immunological Reviews*. 2017;278:20-40.
 70. Smith M, Tourigny MR, Noakes P, Thornton CA, Tulic MK, Prescott SL. Children with egg allergy have evidence of reduced neonatal CD4+CD25+CD127lo/- regulatory T cell function. *Journal of Allergy and Clinical Immunology*. 2008;121:1460-1775098112.
 71. Tulic MK, Andrews D, Crook ML, *et al*. Changes in thymic regulatory T-cell maturation from birth to puberty: Differences in atopic children. *Journal of Allergy and Clinical Immunology*. 2012;129:199-2060000.
 72. Grindebacke H, Wing K, Andersson AC, Suri-Payer E, Rak S, Rudin A. Defective suppression of Th2 cytokines by CD4+CD25+ regulatory T cells in birch allergics during birch pollen season. *Clinical & Experimental Allergy*. 2004;34:1364-1372.
 73. Karlsson MR, Rugtveit J, Brandtzaeg P. Allergen-responsive CD4+CD25+ Regulatory T Cells in Children who Have Outgrown Cow's Milk Allergy. *The Journal of Experimental Medicine*. 2004;199:1679-1688.
 74. Palomares O, Martín-Fontecha M, Lauener R, *et al*. Regulatory T cells and immune regulation of allergic diseases: roles of IL-10 and TGF- β . *Genes and immunity*. 2014;15:511-520.
 75. Thirion D, Morianos I, Xanthou G, Samitas K. Innate immunity as the orchestrator of allergic airway inflammation and resolution in asthma. *International Immunopharmacology*. 2017;48:43-54.
 76. Miyata J, Arita M. Role of omega-3 fatty acids and their metabolites in asthma and allergic diseases. *Allergology international : official journal of the Japanese Society of Allergology*. 2015;64:27-34.
 77. Waddington CH. The Epigenotype. *International Journal of Epidemiology*. 2012;41:10-13.
 78. Ambrosi C, Manzo M, Baubec T. Dynamics and Context-Dependent Roles of DNA Methylation. *Journal of Molecular Biology*. 2017;429:1459-1475.

79. Schmid C, Delacher M, Huehn J, Feuerer M. Epigenetic mechanisms regulating T-cell responses. *The Journal of allergy and clinical immunology*. 2018;142:728-743.
80. Alisch RS, Barwick BG, Chopra P, *et al*. Age-associated DNA methylation in pediatric populations. *Genome Research*. 2012;22:623-632.
81. Zhao M, Qin J, Yin H, *et al*. Distinct epigenomes in CD4(+) T cells of newborns, middle-ages and centenarians. *Scientific reports*. 2016;6:38411.
82. Martino DJ, Tulic MK, Gordon L, *et al*. Evidence for age-related and individual-specific changes in DNA methylation profile of mononuclear cells during early immune development in humans. *Epigenetics*. 2011;6:1085-1094.
83. Alhamwe B, Alhamdan F, Ruhl A, Potaczek DP, Renz H. The role of epigenetics in allergy and asthma development. *Current Opinion in Allergy and Clinical Immunology*. 2020;20:48-55.
84. Deaton AM, Bird A. CpG islands and the regulation of transcription. *Genes & Development*. 2011;25:1010-1022.
85. Smith ZD, Meissner A. DNA methylation: roles in mammalian development. *Nature Reviews Genetics*. 2013;14:204-220.
86. Baribault C, Ehrlich KC, Ponnaluri VKCKC, Pradhan S, Lacey M, Ehrlich M. Developmentally linked human DNA hypermethylation is associated with down-modulation, repression, and upregulation of transcription. *Epigenetics*. 2018;13:275-289.
87. Baubec T, Schübeler D. Genomic patterns and context specific interpretation of DNA methylation. *Current Opinion in Genetics & Development*. 2014;25:85-92.
88. Schübeler D. Function and information content of DNA methylation. *Nature*. 2015;517:321-326.
89. Nugent BM, Bale TL. The omniscient placenta: Metabolic and epigenetic regulation of fetal programming. *Frontiers in Neuroendocrinology*. 2015;39:28-37.
90. Morales-Nebreda L, McLafferty FS, Singer BD. DNA methylation as a transcriptional regulator of the immune system. *Translational Research*. 2018;204:1-18.
91. Rodriguez RM, Suarez-Alvarez B, Mosén-Ansorena D, *et al*. Regulation of the transcriptional program by DNA methylation during human $\alpha\beta$ T-cell development. *Nucleic Acids Research*. 2015;43:760-774.
92. Popovic M, Fiano V, Fasanelli F, *et al*. Differentially methylated DNA regions in early childhood wheezing: An epigenome-wide study using saliva. *Pediatric Allergy and Immunology*. 2019;30:305-314.
93. DeVries A, Wlasiuk G, Miller SJ, *et al*. Epigenome-wide analysis links SMAD3 methylation at birth to asthma in children of asthmatic mothers. *J Allergy Clin Immunol*.
94. Lund RJ, Osmala M, Malonzo M, *et al*. Atopic asthma after rhinovirus-induced wheezing is associated with DNA methylation change in the SMAD3 gene promoter. *Allergy*. 2018;73:1735-1740.
95. Xu C-J, Söderhäll C, Bustamante M, *et al*. DNA methylation in childhood asthma: an epigenome-wide meta-analysis. *The Lancet Respiratory medicine*. 2018;6:379-388.
96. Reese SE, Xu C-J, den Dekker HT, *et al*. Epigenome-wide Meta-analysis of DNA Methylation and Childhood Asthma. *The Journal of allergy and clinical immunology*. 2018.
97. Martino D, Neeland M, Dang T, *et al*. Epigenetic dysregulation of naive CD4+ T-cell activation genes in childhood food allergy. *Nature Communications*. 2018;9:3308.
98. Martino D, Joo JE, Sexton-Oates A, *et al*. Epigenome-wide association study reveals longitudinally stable DNA methylation differences in CD4+ T cells from children with IgE-mediated food allergy. *Epigenetics*. 2014;9:998-1006.
99. Martino D, Dang T, Sexton-Oates A, *et al*. Blood DNA methylation biomarkers predict clinical reactivity in food-sensitized infants. *J Allergy Clin Immunol*. 135.
100. Quraishi BM, Zhang H, Everson TM, *et al*. Identifying CpG sites associated with eczema via random forest screening of epigenome-scale DNA methylation. *Clin Epigenetics*. 2015;7.
101. Zhang H, Kaushal A, Merid S, *et al*. DNA methylation and allergic sensitizations, a genome-scale longitudinal study during adolescence. *Allergy*. 2019;74:1166-1175.
102. Peng C, Meel ER, Cardenas A, *et al*. Epigenome-wide association study reveals methylation pathways associated with childhood allergic sensitization. *Epigenetics*. 2019.
103. Everson TM, Lyons G, Zhang H, *et al*. DNA methylation loci associated with atopy and high serum IgE: a genome-wide application of recursive Random Forest feature selection. *Genome Med*. 7.
104. Knudsen T, Rezwan FI, Jiang Y, Karmaus W, Svanes C, Holloway JW. Trans- and inter-generational epigenetic inheritance in allergic diseases. *Journal of Allergy and Clinical Immunology*. 2018.

105. Hughes CE, Nibbs RJB. A guide to chemokines and their receptors. *The FEBS Journal*. 2018;285:2944-2971.
106. Pease J, Williams T. Chemokines and their receptors in allergic disease. *Journal of Allergy and Clinical Immunology*. 2006;118:305-318.
107. Groom JR, Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunology and cell biology*. 2011;89:207-215.
108. Lacotte S, Brun S, Muller S, Dumortier H. CXCR3, Inflammation, and Autoimmune Diseases. *Annals of the New York Academy of Sciences*. 2009;1173:310-317.
109. Romagnani P, Annunziato F, Lazzeri E, *et al*. Interferon-inducible protein 10, monokine induced by interferon gamma, and interferon-inducible T-cell alpha chemoattractant are produced by thymic epithelial cells and attract T-cell receptor (TCR) $\alpha\beta$ +CD8+ single-positive T cells, TCR $\gamma\delta$ + T cells, and natural killer-type cells in human thymus. *Blood*. 2001;97:601-607.
110. Porter JC, Falzon M, Hall A. Polarized Localization of Epithelial CXCL11 in Chronic Obstructive Pulmonary Disease and Mechanisms of T Cell Egression. *The Journal of Immunology*. 2008;180:1866-1877.
111. Cole KE, Strick CA, Paradis TJ, *et al*. Interferon-inducible T Cell Alpha Chemoattractant (I-TAC): A Novel Non-ELR CXC Chemokine with Potent Activity on Activated T Cells through Selective High Affinity Binding to CXCR3. *Journal of Experimental Medicine*. 1998;187:2009-2021.
112. Fenwick PS, Macedo P, Kilty IC, Barnes PJ, Donnelly LE. Effect of JAK Inhibitors on Release of CXCL9, CXCL10 and CXCL11 from Human Airway Epithelial Cells. *PLOS ONE*. 2015;10.
113. Abrahamsson TR, Sandberg Abelius M, Forsberg A, Björkstén B, Jenmalm MC. A Th1/Th2-associated chemokine imbalance during infancy in children developing eczema, wheeze and sensitization. *Clinical and experimental allergy: journal of the British Society for Allergy and Clinical Immunology*. 2011;41:1729-1739.
114. Reubsaet LL, Meerding J, de Jager W, *et al*. Plasma chemokines in early wheezers predict the development of allergic asthma. *American journal of respiratory and critical care medicine*. 2013;188:1039-1040.
115. Tworek D, Kuna P, Młynarski W, Górski P, Pietras T, Antczak A. MIG (CXCL9), IP-10 (CXCL10) and I-TAC (CXCL11) concentrations after nasal allergen challenge in patients with allergic rhinitis. *Archives of Medical Science*. 2013;5:849-853.
116. Southworth T, Pattwell C, Khan N, *et al*. Increased type 2 inflammation post rhinovirus infection in patients with moderate asthma. *Cytokine*. 2020;125:154857.
117. Ghebre MA, Pang PH, Desai D, *et al*. Severe exacerbations in moderate-to-severe asthmatics are associated with increased pro-inflammatory and type 1 mediators in sputum and serum. *BMC pulmonary medicine*. 2019;19:144.
118. Wark PAB, Bucchieri F, Johnston SL, *et al*. IFN- γ -induced protein 10 is a novel biomarker of rhinovirus-induced asthma exacerbations. *Journal of Allergy and Clinical Immunology*. 2007;120:586-593.
119. Moskwa S, Piotrowski W, Marczak J, *et al*. Innate Immune Response to Viral Infections in Primary Bronchial Epithelial Cells is Modified by the Atopic Status of Asthmatic Patients. *Allergy, Asthma & Immunology Research*. 2018;10:144.
120. Gauthier M, Chakraborty K, Oriss TB, *et al*. Severe asthma in humans and mouse model suggests a CXCL10 signature underlies corticosteroid-resistant Th1 bias. *JCI Insight*. 2017;2.
121. Yoshie O, Matsushima K. CCR4 and its ligands: from bench to bedside. *International Immunology*. 2014;27:11-20.
122. Scheu S, Ali S, Ruland C, Arolt V, Alferink J. The C-C Chemokines CCL17 and CCL22 and Their Receptor CCR4 in CNS Autoimmunity. *International Journal of Molecular Sciences*. 2017;18:2306.
123. Wirnsberger G, Hebenstreit D, Posselt G, Horejs-Hoeck J, Duschl A. IL-4 induces expression of TARC/CCL17 via two STAT6 binding sites. *European Journal of Immunology*. 2006;36:1882-1891.
124. Sandberg M, Frykman A, Ernerudh J, *et al*. Cord blood cytokines and chemokines and development of allergic disease. *Pediatric allergy and immunology: official publication of the European Society of Pediatric Allergy and Immunology*. 2009;20:519-527.
125. Abelius MS, Ernerudh J, Berg G, Matthiesen L, Nilsson LJ, Jenmalm MC. High cord blood levels of the T-helper 2-associated chemokines CCL17 and CCL22 precede allergy development during the first 6 years of life. *Pediatric research*. 70.

126. Björkander S, Carvalho-Queiroz C, Hallberg J, *et al.* Childhood allergy is preceded by an absence of gut lactobacilli species and higher levels of atopy-related plasma chemokines. *Clinical & Experimental Immunology*. 2020.
127. Chiu C-Y, Su K-W, Tsai M-H, *et al.* Low Mother-to-Child CCL22 Chemokine Levels Are Inversely Related to Mite Sensitization and Asthma in Early Childhood. *Scientific reports*. 2018;8:6043.
128. Chenivesse C, Tscopoulos A. CCL18 – Beyond chemotaxis. *Cytokine*. 2018;109:52-56.
129. Kodelja V, Müller C, Politz O, Hakij N, Orfanos CE, Goerd S. Alternative macrophage activation-associated CC-chemokine-1, a novel structural homologue of macrophage inflammatory protein-1 alpha with a Th2-associated expression pattern. *Journal of immunology (Baltimore, Md : 1950)*. 1998;160:1411-1418.
130. Vulcano M, Struyf S, Scapini P, *et al.* Unique Regulation of CCL18 Production by Maturing Dendritic Cells. *The Journal of Immunology*. 2003;170:3843-3849.
131. Tiemessen MM, Jagger AL, Evans HG, van Herwijnen MJC, John S, Taams LS. CD4+CD25+Foxp3+ regulatory T cells induce alternative activation of human monocytes/macrophages. *Proceedings of the National Academy of Sciences*. 2007;104:19446-19451.
132. Islam SA, Ling MF, Leung J, Shreffler WG, Luster AD. Identification of human CCR8 as a CCL18 receptor. *The Journal of Experimental Medicine*. 2013;210:1889-1898.
133. de Nadai P, Charbonnier A-S, Chenivesse C, *et al.* Involvement of CCL18 in Allergic Asthma. *The Journal of Immunology*. 2006;176:6286-6293.
134. Yang IV, Tomfohr J, Singh J, *et al.* The Clinical and Environmental Determinants of Airway Transcriptional Profiles in Allergic Asthma. *American Journal of Respiratory and Critical Care Medicine*. 2012;185:620-627.
135. Peterson S, Poposki JA, Nagarkar DR, *et al.* Increased expression of CC chemokine ligand 18 in patients with chronic rhinosinusitis with nasal polyps. *Journal of Allergy and Clinical Immunology*. 2012;129:119.
136. Nomura I, Gao B, Boguniewicz M, Darst MA, Travers JB, Leung D. Distinct patterns of gene expression in the skin lesions of atopic dermatitis and psoriasis A gene microarray analysis. *Journal of Allergy and Clinical Immunology*. 2003;112:1195-1202.
137. Kim HO, Cho SI, Chung BY, Ahn HK, Park CW, Lee CH. Expression of CCL1 and CCL18 in atopic dermatitis and psoriasis. *Clinical and Experimental Dermatology*. 2012;37:521-526.
138. Chang Y, de Nadai P, Azzaoui I, *et al.* The chemokine CCL18 generates adaptive regulatory T cells from memory CD4 + T cells of healthy but not allergic subjects. *The FASEB Journal*. 2010;24:5063-5072.
139. Azzaoui I, Yahia S, Chang Y, *et al.* CCL18 differentiates dendritic cells in tolerogenic cells able to prime regulatory T cells in healthy subjects. *Blood*. 2011;118:3549-3558.
140. Luzina IG, Todd NW, Nacu N, *et al.* Regulation of pulmonary inflammation and fibrosis through expression of integrins alphaVbeta3 and alphaVbeta5 on pulmonary T lymphocytes. *Arthritis and rheumatism*. 2009;60:1530-1539.
141. Abeli MS, Janefjord C, Ernerudh J, *et al.* The Placental Immune Milieu is Characterized by a Th2- and Anti-Inflammatory Transcription Profile, Regardless of Maternal Allergy, and Associates with Neonatal Immunity. *American Journal of Reproductive Immunology*. 2015;73:445-459.
142. Svensson-Arvelund J, Ernerudh J, Buse E, *et al.* The Placenta in Toxicology. Part II. *Toxicologic Pathology*. 2014;42:327-338.
143. Dominguez-Bello M, Godoy-Vitorino F, Knight R, Blaser MJ. Role of the microbiome in human development. *Gut*. 2019;68:1108-1114.
144. Park J-E, Jardine L, Gottgens B, Teichmann SA, Haniffa M. Prenatal development of human immunity. *Science*. 2020;368:600-603.
145. Mirpuri J. Evidence for maternal diet-mediated effects on the offspring microbiome and immunity: implications for public health initiatives. *Pediatric Research*. 2020:1-7.
146. Albrecht M, Arck P. Vertically Transferred Immunity in Neonates: Mothers, Mechanisms and Mediators. *Frontiers in Immunology*. 2020;11:555.
147. Vuillermin PJ, Macia L, Nanan R, Tang MLK, Collier F, Brix S. The maternal microbiome during pregnancy and allergic disease in the offspring. *Seminars in Immunopathology*. 2017;39:669-675.
148. Blewett HJ, Cicalo MC, Holland CD, Field CJ. The immunological components of human milk. *Advances in food and nutrition research*. 2008;54:45-80.

149. Aagaard K, Ma J, Antony KM, Ganu R, Petrosino J, Versalovic J. The placenta harbors a unique microbiome. *Science translational medicine*. 2014;6.
150. Jiménez E, Marin ML, Martin R, *et al*. Is meconium from healthy newborns actually sterile? *Research in Microbiology*. 2008;159:187-193.
151. McDonald B, McCoy KD. Maternal microbiota in pregnancy and early life. *Science*. 2019;365:984-985.
152. Jakobsson HE, Abrahamsson TR, Jenmalm MC, *et al*. Decreased gut microbiota diversity, delayed Bacteroidetes colonisation and reduced Th1 responses in infants delivered by Caesarean section. *Gut*. 2014;63:559.
153. Francino PM. Birth Mode-Related Differences in Gut Microbiota Colonization and Immune System Development. *Annals of Nutrition and Metabolism*. 2018;73:12-16.
154. Macpherson AJ, de Agüero MG, Ganai-Vonarburg SC. How nutrition and the maternal microbiota shape the neonatal immune system. *Nature reviews Immunology*. 2017;17:508-517.
155. Julia V, Macia L, Dombrowicz D. The impact of diet on asthma and allergic diseases. *Nature Reviews Immunology*. 2015;15:308-322.
156. Dzidic M, Boix-Amorós A, Selma-Royo M, Mira A, Collado M. Gut Microbiota and Mucosal Immunity in the Neonate. *Medical sciences (Basel, Switzerland)*. 2018;6:56.
157. Renz H, Brandtzaeg P, Hornef M. The impact of perinatal immune development on mucosal homeostasis and chronic inflammation. *Nature Reviews Immunology*. 2012;12:9.
158. Tokuhara D, Kurashima Y, Kamioka M, Nakayama T, Ernst P, Kiyono H. A comprehensive understanding of the gut mucosal immune system in allergic inflammation. *Allergy International*. 2018;68:17-25.
159. Hooper LV, Macpherson AJ. Immune adaptations that maintain homeostasis with the intestinal microbiota. *Nature Reviews Immunology*. 2010;10:159-169.
160. Hooper LV, Littman DR, Macpherson AJ. Interactions Between the Microbiota and the Immune System. *Science*. 2012;336:1268-1273.
161. Kimura S. Molecular insights into the mechanisms of M-cell differentiation and transcytosis in the mucosa-associated lymphoid tissues. *Anatomical Science International*. 2018;93:23-34.
162. Rios D, Wood MB, Li J, Chassaing B, Gewirtz AT, Williams IR. Antigen sampling by intestinal M cells is the principal pathway initiating mucosal IgA production to commensal enteric bacteria. *Mucosal Immunology*. 2016;9:907-916.
163. Mora RJ, Iwata M, Eksteen B, *et al*. Generation of Gut-Homing IgA-Secreting B Cells by Intestinal Dendritic Cells. *Science*. 2006;314:1157-1160.
164. Wells JM. Immunomodulatory mechanisms of lactobacilli. *Microbial cell factories*. 2011;10 Suppl 1.
165. Niess J, Brand S, Gu X, *et al*. CX₃CR1-Mediated Dendritic Cell Access to the Intestinal Lumen and Bacterial Clearance. *Science*. 2005;307:254-258.
166. Denning TL, Wang Y-c, Patel SR, Williams IR, Pulendran B. Lamina propria macrophages and dendritic cells differentially induce regulatory and interleukin 17-producing T cell responses. *Nature Immunology*. 2007;8:1086-1094.
167. Islam SA, Luster AD. T cell homing to epithelial barriers in allergic disease. *Nature Medicine*. 2012;18:705-715.
168. Atarashi K, Umesaki Y, Honda K. Microbial influence on T cell subset development. *Seminars in immunology*. 2011;23:146-153.
169. Goto Y, Panea C, Nakato G, *et al*. Segmented Filamentous Bacteria Antigens Presented by Intestinal Dendritic Cells Drive Mucosal Th17 Cell Differentiation. *Immunity*. 2014;40:594-607.
170. Ivanov II, Atarashi K, Manel N, *et al*. Induction of Intestinal Th17 Cells by Segmented Filamentous Bacteria. *Cell*. 2009;139:485-498.
171. Jonsson H, Hugerth LW, Sundh J, Lundin E, Andersson AF. Genome sequence of segmented filamentous bacteria present in the human intestine. *Communications Biology*. 2020;3:485.
172. Furusawa Y, Obata Y, Hase K. Commensal microbiota regulates T cell fate decision in the gut. *Seminars in Immunopathology*. 2015;37:17-25.
173. Wu R-QQ, Zhang D-FF, Tu E, Chen Q-MM, Chen W. The mucosal immune system in the oral cavity-an orchestra of T cell diversity. *International journal of oral science*. 2014;6:125-132.
174. Moingeon P, Mascarell L. Differences and similarities between sublingual immunotherapy of allergy and oral tolerance. *Seminars in Immunology*. 2017;30:52-60.

175. Allam J-P, Novak N, Fuchs C, *et al.* Characterization of dendritic cells from human oral mucosa: A new Langerhans' cell type with high constitutive FcεRI expression. *Journal of Allergy and Clinical Immunology*. 2003;112:141-148.
176. Carlier FM, Sibille Y, Pilette C. The epithelial barrier and immunoglobulin A system in allergy. *Clinical & Experimental Allergy*. 2016;46:1372-1388.
177. Goleva E, Berdyshev E, Leung D. Epithelial barrier repair and prevention of allergy. *Journal of Clinical Investigation*. 2019;129:1463-1474.
178. West CE, Jenmalm MC, Prescott SL. The gut microbiota and its role in the development of allergic disease: a wider perspective. *Clinical & Experimental Allergy*. 2015;45:43-53.
179. Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm MC. Low diversity of the gut microbiota in infants with atopic eczema. *The Journal of allergy and clinical immunology*. 2012;129:434.
180. Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm MC. Low gut microbiota diversity in early infancy precedes asthma at school age. *Clinical & Experimental Allergy*. 2014;44:842-850.
181. Wang M, Karlsson C, Olsson C, *et al.* Reduced diversity in the early fecal microbiota of infants with atopic eczema. *Journal of Allergy and Clinical Immunology*. 2008;121:129-134.
182. Ling Z, Li Z, Liu X, *et al.* Altered Fecal Microbiota Composition Associated with Food Allergy in Infants. *Applied and Environmental Microbiology*. 2014;80:2546-2554.
183. Bäckhed F, Roswall J, Peng Y, *et al.* Dynamics and Stabilization of the Human Gut Microbiome during the First Year of Life. *Cell Host & Microbe*. 2015;17:690-703.
184. Weng M, Walker WA. The role of gut microbiota in programming the immune phenotype. *Journal of Developmental Origins of Health and Disease*. 2013;4:203-214.
185. Normand A-C, Sudre B, Vacheyrou M, *et al.* Airborne cultivable microflora and microbial transfer in farm buildings and rural dwellings. *Occupational and Environmental Medicine*. 2011;68:849.
186. Azad MB, Konya T, Maughan H, *et al.* Infant gut microbiota and the hygiene hypothesis of allergic disease: impact of household pets and siblings on microbiota composition and diversity. *Allergy, Asthma & Clinical Immunology*. 2013;9:15.
187. Jatzlauk G, Bartel S, Heine H, Schlöter M, Krauss-Etschmann S. Influences of environmental bacteria and their metabolites on allergies, asthma, and host microbiota. *Allergy*. 2017;72:1859-1867.
188. Bokulich NA, Chung J, Battaglia T, *et al.* Antibiotics, birth mode, and diet shape microbiome maturation during early life. *Science Translational Medicine*. 2016;8.
189. Torow N, Marsland BJ, Hornef MW, Gollwitzer ES. Neonatal mucosal immunology. *Mucosal Immunology*. 2016;10:5-17.
190. Dominguez-Bello MG, Costello EK, Contreras M, *et al.* Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;107:11971-11975.
191. Shao Y, Forster SC, Tsalki E, *et al.* Stunted microbiota and opportunistic pathogen colonization in caesarean-section birth. *Nature*. 2019;574:117-121.
192. Wang S, Ryan AC, Boyaval P, Dempsey EM, Ross PR, Stanton C. Maternal Vertical Transmission Affecting Early-life Microbiota Development. *Trends in Microbiology*. 2019;28:28-45.
193. Ferretti P, Pasolli E, Tett A, *et al.* Mother-to-Infant Microbial Transmission from Different Body Sites Shapes the Developing Infant Gut Microbiome. *Cell Host & Microbe*. 2018;24:133-14500000.
194. Dzidic M, Mira A, Artacho A, Abrahamsson TR, Jenmalm MC, Collado M. Allergy development is associated with consumption of breastmilk with a reduced microbial richness in the first month of life. *Pediatric Allergy and Immunology*. 2020;31:250-257.
195. Azad MB, Konya T, Guttman DS, *et al.* Infant gut microbiota and food sensitization: associations in the first year of life. *Clinical & Experimental Allergy*. 2015;45:632-643.
196. Sjögren YM, Jenmalm MC, Böttcher MF, Björkstén B, Sverremark-Ekström E. Altered early infant gut microbiota in children developing allergy up to 5 years of age. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2009;39:518-526.
197. Fujimura KE, Sitarik AR, Havstad S, *et al.* Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nature Medicine*. 2016;22:1187-1191.
198. Arrieta M-C, Stiemsma LT, Dimitriu PA, *et al.* Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Science Translational Medicine*. 2015;7.

199. Macpherson AJ, McCoy KD, Johansen FEE, Brandtzaeg P. The immune geography of IgA induction and function. *Mucosal immunology*. 2008;1:11-22.
200. Pabst O, Slack E. IgA and the intestinal microbiota: the importance of being specific. *Mucosal Immunology*. 2020;13:12-21.
201. Brandtzaeg P. Secretory immunity with special reference to the oral cavity. *Journal of oral microbiology*. 2013;5.
202. Kawamoto S, Maruya M, Kato LM, *et al*. Foxp3+ T Cells Regulate Immunoglobulin A Selection and Facilitate Diversification of Bacterial Species Responsible for Immune Homeostasis. *Immunity*. 2014;41:152-165.
203. Peterson DA, McNulty NP, Guruge JL, Gordon JL. IgA Response to Symbiotic Bacteria as a Mediator of Gut Homeostasis. *Cell Host & Microbe*. 2007;2:328-339.
204. Nakajima A, Vogelzang A, Maruya M, *et al*. IgA regulates the composition and metabolic function of gut microbiota by promoting symbiosis between bacteria. *Journal of Experimental Medicine*. 2018;215.
205. Mathias A, Pais B, Favre L, Benyacoub J, Corthésy B. Role of secretory IgA in the mucosal sensing of commensal bacteria. *Gut Microbes*. 2014;5:688-695.
206. Bunker JJ, Erickson SA, Flynn TM, *et al*. Natural polyreactive IgA antibodies coat the intestinal microbiota. *Science*. 2017;358.
207. Dzidic M, Abrahamsson TR, Artacho A, *et al*. Aberrant IgA responses to the gut microbiota during infancy precede asthma and allergy development. *The Journal of allergy and clinical immunology*. 2017;139:1017.
208. Holgate ST. Innate and adaptive immune responses in asthma. *Nature Medicine*. 2012;18:673-683.
209. Cerutti A. The regulation of IgA class switching. *Nature Reviews Immunology*. 2008;8:421-434.
210. Gutzeit C, Magri G, Cerutti A. Intestinal IgA production and its role in host-microbe interaction. *Immunological Reviews*. 2014;260:76-85.
211. Fagerås M, Tomićić S, Voor T, Björkstén B, Jenmalm MC. Slow salivary secretory IgA maturation may relate to low microbial pressure and allergic symptoms in sensitized children. *Pediatric research*. 2011;70:572-577.
212. Johansen FE, Kaetzel CS. Regulation of the polymeric immunoglobulin receptor and IgA transport: new advances in environmental factors that stimulate pIgR expression and its role in mucosal immunity. *Mucosal Immunology*. 2011;4:598-602.
213. Sjögren YM, Tomicic S, Lundberg A, *et al*. Influence of early gut microbiota on the maturation of childhood mucosal and systemic immune responses. *Clinical & Experimental Allergy*. 2009;39:1842-1851.
214. Sandin A, Björkstén B, Böttcher MF, Englund E, Jenmalm MC, Bråbäck L. High salivary secretory IgA antibody levels are associated with less late-onset wheezing in IgE-sensitized infants. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2011;22:477-481.
215. Johansson MA, Sjögren YM, Persson J-O, Nilsson C, Sverremark-Ekström E. Early Colonization with a Group of Lactobacilli Decreases the Risk for Allergy at Five Years of Age Despite Allergic Heredity. *PLoS ONE*. 2011;6.
216. Verschoor D, von Gunten S. Allergy and Atopic Diseases: An Update on Experimental Evidence. *International Archives of Allergy and Immunology*. 2019;180:235-243.
217. Eyerich S, Metz M, Bossios A, Eyerich K. New biological treatments for asthma and skin allergies. *Allergy*. 2020;75:546-560.
218. Miles EA, Calder PC. Can Early Omega-3 Fatty Acid Exposure Reduce Risk of Childhood Allergic Disease? *Nutrients*. 2017;9:784.
219. Alvaro-Lozano M, Akdis CA, Akdis M, *et al*. Allergen Immunotherapy in Children User's Guide. *Pediatric Allergy and Immunology*. 2020;31:1-101.
220. Stokholm J, Blaser MJ, Thorsen J, *et al*. Maturation of the gut microbiome and risk of asthma in childhood. *Nature Communications*. 2018;9:141.
221. Bisgaard H, Li N, Bonnelykke K, *et al*. Reduced diversity of the intestinal microbiota during infancy is associated with increased risk of allergic disease at school age. *Journal of Allergy and Clinical Immunology*. 2011;128:646-65200000.

222. Chiu C-Y, Chan Y-L, Tsai M-H, Wang C-J, Chiang M-H, Chiu C-C. Gut microbial dysbiosis is associated with allergen-specific IgE responses in young children with airway allergies. *The World Allergy Organization Journal*. 2019;12:100021.
223. Cuello-Garcia CA, Brozek JL, Fiocchi A, *et al*. Probiotics for the prevention of allergy: A systematic review and meta-analysis of randomized controlled trials. *J Allergy Clin Immunol*.136.
224. Organization FaAOotUNaWH. Probiotics in food - Health and nutritional properties and guidelines for evaluation. 2001. <http://www.fao.org/3/a-a0512e.pdf>.
225. Kleerebezem M, Hols P, Bernard E, *et al*. The extracellular biology of the lactobacilli. *FEMS microbiology reviews*. 2010;34:199-230.
226. Mu Q, Tavella VJ, Luo XM. Role of Lactobacillus reuteri in Human Health and Diseases. *Frontiers in Microbiology*. 2018;9:757.
227. van Baarlen P, Wells JM, Kleerebezem M. Regulation of intestinal homeostasis and immunity with probiotic lactobacilli. *Trends in Immunology*. 2013;34:208-215.
228. Smits HH, Engering A, van der Kleij D, *et al*. Selective probiotic bacteria induce IL-10–producing regulatory T cells in vitro by modulating dendritic cell function through dendritic cell–specific intercellular adhesion molecule 3–grabbing nonintegrin. *Journal of Allergy and Clinical Immunology*. 2005;115:1260-1267.
229. Forsberg A, West CE, Prescott SL, Jenmalm MC. Pre- and probiotics for allergy prevention: time to revisit recommendations? *Clin Exp Allergy*.46.
230. Boyle RJ, Ismail IH, Kivivuori S, *et al*. Lactobacillus GG treatment during pregnancy for the prevention of eczema: a randomized controlled trial. *Allergy*. 2011;66:509-516.
231. Taylor AL, Dunstan JA, Prescott SL. Probiotic supplementation for the first 6 months of life fails to reduce the risk of atopic dermatitis and increases the risk of allergen sensitization in high-risk children: A randomized controlled trial. *Journal of Allergy and Clinical Immunology*. 2007;119:184-191.
232. West CE, Hammarström ML, Hernell O. Probiotics during weaning reduce the incidence of eczema. *Pediatric Allergy and Immunology*. 2009;20:430-437.
233. Soh SE, Aw M, Gerez I, *et al*. Probiotic supplementation in the first 6 months of life in at risk Asian infants – effects on eczema and atopic sensitization at the age of 1 year. *Clinical & Experimental Allergy*. 2009;39:571-578.
234. Cabana MD, McKean M, Caughey AB, *et al*. Early Probiotic Supplementation for Eczema and Asthma Prevention: A Randomized Controlled Trial. *Pediatrics*. 2017;140.
235. Kukkonen K, Savilahti E, Haahtela T, *et al*. Probiotics and prebiotic galacto-oligosaccharides in the prevention of allergic diseases: A randomized, double-blind, placebo-controlled trial. *Journal of Allergy and Clinical Immunology*. 2007;119:192-198.
236. Wickens K, Black PN, Stanley TV, *et al*. A differential effect of 2 probiotics in the prevention of eczema and atopy: A double-blind, randomized, placebo-controlled trial. *Journal of Allergy and Clinical Immunology*. 2008;122:788-794.
237. Kalliomäki M, Salminen S, Arvilommi H, Kero P, Koskinen P, Isolauri E. Probiotics in primary prevention of atopic disease: a randomised placebo-controlled trial. *The Lancet*. 2001;357:1076-1079.
238. Dotterud CK, Storrø O, Johnsen R, Øien T. Probiotics in pregnant women to prevent allergic disease: a randomized, double-blind trial. *British Journal of Dermatology*. 2010;163:616-623.
239. Allen SJ, Jordan S, Storey M, *et al*. Probiotics in the prevention of eczema: a randomised controlled trial. *Archives of Disease in Childhood*. 2014;99:1014.
240. Kim J, Kwon J, Ahn S, *et al*. Effect of probiotic mix (Bifidobacterium bifidum, Bifidobacterium lactis, Lactobacillus acidophilus) in the primary prevention of eczema: a double-blind, randomized, placebo-controlled trial. *Pediatric Allergy and Immunology*. 2010;21.
241. Abrahamsson TR, Jakobsson T, Böttcher MF, *et al*. Probiotics in prevention of IgE-associated eczema: a double-blind, randomized, placebo-controlled trial. *J Allergy Clin Immunol*. 2007;119.
242. Kalliomäki M, Salminen S, Poussa T, Arvilommi H, Isolauri E. Probiotics and prevention of atopic disease: 4-year follow-up of a randomised placebo-controlled trial. *The Lancet*. 2003;361:1869-1871.
243. Abrahamsson TR, Jakobsson T, Björkstén B, Oldaeus G, Jenmalm MC. No effect of probiotics on respiratory allergies: a seven-year follow-up of a randomized controlled trial in infancy. *Pediatr Allergy Immunol*.24.

244. Wickens K, Barthow C, Mitchell EA, *et al.* Effects of *Lactobacillus rhamnosus* HN001 in early life on the cumulative prevalence of allergic disease to 11 years. *Pediatric Allergy and Immunology*. 2018.
245. Kalliomäki M, Salminen S, Poussa T, Isolauri E. Probiotics during the first 7 years of life: A cumulative risk reduction of eczema in a randomized, placebo-controlled trial. *Journal of Allergy and Clinical Immunology*. 2007;119:1019-1021.
246. De Benedetto A, Rafaels NM, McGirt LY, *et al.* Tight junction defects in patients with atopic dermatitis. *The Journal of allergy and clinical immunology*. 2011;127:773.
247. West CE, Dzidic M, Prescott SL, Jenmalm MC. Bugging allergy; role of pre-, pro- and synbiotics in allergy prevention. *Allergology International*. 2017;66:529-538.
248. Forsberg A, Abrahamsson TR, Björkstén B, Jenmalm MC. Pre- and postnatal administration of *Lactobacillus reuteri* decreases TLR2 responses in infants. *Clin Transl Allergy*. 2014;4.
249. Forsberg A, Abrahamsson TR, Björkstén B, Jenmalm MC. Pre- and post-natal *Lactobacillus reuteri* supplementation decreases allergen responsiveness in infancy. *Clin Exp Allergy*. 43.
250. Christie WW. <https://www.lipidmaps.org/resources/lipidweb/index.php?page=lipids/lipids.html>.
251. Han X. Lipidomics for studying metabolism. *Nature reviews Endocrinology*. 2016;12:668-679.
252. Visioli F. Lipidomics to Assess Omega 3 Bioactivity. *Journal of clinical medicine*. 2015;4:1753-1760.
253. Trzicka E, Kremmyda L-S, Stankova B, Zak A. FATTY ACIDS AS BIOCOMPOUNDS: THEIR ROLE IN HUMAN METABOLISM, HEALTH AND DISEASE - A REVIEW. PART 1: CLASSIFICATION, DIETARY SOURCES AND BIOLOGICAL FUNCTIONS. *Biomedical Papers*. 2011;155:117-130.
254. Duvall MG, Levy BD. DHA- and EPA-derived resolvins, protectins, and maresins in airway inflammation. *European journal of pharmacology*. 2016;785:144-155.
255. Barnig C, Frossard N, Levy BD. Towards targeting resolution pathways of airway inflammation in asthma. *Pharmacology & therapeutics*. 2018;186:98-113.
256. Del Moral MG, Martínez-Naves E. The Role of Lipids in Development of Allergic Responses. *Immune network*. 2017;17:133-143.
257. Lumia M, Luukkainen P, Takkinen H-M, *et al.* Cow's milk allergy and the association between fatty acids and childhood asthma risk. *Journal of Allergy and Clinical Immunology*. 2014;134:488-49000.
258. Weise C, Hilt K, Milovanovic M, Ernst D, Rühl R, Worm M. Inhibition of IgE production by docosahexaenoic acid is mediated by direct interference with STAT6 and NFκB pathway in human B cells. *The Journal of Nutritional Biochemistry*. 2011;22:269-275.
259. van den Elsen LWJ, Nüsse Y, Balvers M, *et al.* n-3 Long-chain PUFA reduce allergy-related mediator release by human mast cells in vitro via inhibition of reactive oxygen species. *British Journal of Nutrition*. 2013;109:1821-1831.
260. Hanebutt FL, Demmelmair H, Schiessl B, Larqué E, Koletzko B. Long-chain polyunsaturated fatty acid (LC-PUFA) transfer across the placenta. *Clinical Nutrition*. 2008;27:685-693.
261. Krauss-Etschmann S, Hartl D, Rzehak P, *et al.* Decreased cord blood IL-4, IL-13, and CCR4 and increased TGF-β levels after fish oil supplementation of pregnant women. *Journal of Allergy and Clinical Immunology*. 2008;121:464-470000000.
262. Bisgaard H, Stokholm J, Chawes BL, *et al.* Fish Oil-Derived Fatty Acids in Pregnancy and Wheeze and Asthma in Offspring. *The New England journal of medicine*. 2016;375:2530-2539.
263. Hansen S, Strøm M, Maslova E, *et al.* Fish oil supplementation during pregnancy and allergic respiratory disease in the adult offspring. *The Journal of allergy and clinical immunology*. 2016;139:104-1110000.
264. D'Vaz N, Meldrum SJ, Dunstan JA, *et al.* Fish oil supplementation in early infancy modulates developing infant immune responses. *Clinical & Experimental Allergy*. 2012;42:1206-1216.
265. Furuholm C, Warstedt K, Larsson J, *et al.* Fish oil supplementation in pregnancy and lactation may decrease the risk of infant allergy. *Acta Paediatrica*. 2009;98:1461-1467.
266. Furuholm C, Warstedt K, Fagerås M, *et al.* Allergic disease in infants up to 2 years of age in relation to plasma omega-3 fatty acids and maternal fish oil supplementation in pregnancy and lactation. *Pediatric Allergy and Immunology*. 2011;22:505-514.
267. Ring J, Gutermuth J. 100 years of hyposensitization: history of allergen-specific immunotherapy (ASIT). *Allergy*. 2011;66:713-724.
268. Globińska A, Boonpiyathad T, Satitsuksanoa P, *et al.* Mechanisms of allergen-specific immunotherapy: diverse mechanisms of immune tolerance to allergens. *Annals of allergy, asthma, & immunology : official publication of the American College of Allergy, Asthma, & Immunology*. 2018.

269. Hoffmann HJ, Valovirta E, Pfaar O, *et al.* Novel approaches and perspectives in allergen immunotherapy. *Allergy*. 2017;72:1022-1034.
270. Shamji MH, Durham SR. Mechanisms of allergen immunotherapy for inhaled allergens and predictive biomarkers. *The Journal of allergy and clinical immunology*. 2017;140:1485-1498.
271. van de Veen W, Wirz OF, Globinska A, Akdis M. Novel mechanisms in immune tolerance to allergens during natural allergen exposure and allergen-specific immunotherapy. *Current opinion in immunology*. 2017;48:74-81.
272. Gunawardana NC, Durham SR. New approaches to allergen immunotherapy. *Annals of Allergy, Asthma & Immunology*. 2018;121:293-305.
273. Passalacqua G, Bagnasco D, Canonica G. 30 years of sublingual immunotherapy. *Allergy*. 2020;75:1107-1120.
274. Durham SR, Penagos M. Sublingual or subcutaneous immunotherapy for allergic rhinitis? *The Journal of allergy and clinical immunology*. 2016;137:339.
275. Vogelberg C, Brüggemann B, Richter H, Jutel M. <p>Real-World Adherence and Evidence of Subcutaneous and Sublingual Immunotherapy in Grass and Tree Pollen-Induced Allergic Rhinitis and Asthma</p>. *Patient Preference and Adherence*. 2020;Volume 14:817-827.
276. Nelson H, Cartier S, Allen-Ramey F, Lawton S, Calderon MA. Network Meta-analysis Shows Commercialized Subcutaneous and Sublingual Grass Products Have Comparable Efficacy. *The Journal of Allergy and Clinical Immunology: In Practice*. 2015;3:256-266000.
277. Dhami S, Nurmatov U, Arasi S, *et al.* Allergen immunotherapy for allergic rhinoconjunctivitis: A systematic review and meta-analysis. *Allergy*. 2017;72:1597-1631.
278. Rice JL, Diette GB, Suarez-Cuervo C, *et al.* Allergen-Specific Immunotherapy in the Treatment of Pediatric Asthma: A Systematic Review. *Pediatrics*. 2018;141.
279. Poddighe D. Sublingual immunotherapy for pediatric allergic rhinitis: The clinical evidence. *World Journal of Clinical Pediatrics*. 2016;5:47.
280. Möller C, Dreborg S, Ferdousi HA, *et al.* Pollen immunotherapy reduces the development of asthma in children with seasonal rhinoconjunctivitis (the PAT-study). *The Journal of allergy and clinical immunology*. 2002;109:251-256.
281. Valovirta E, Berstad AK, de Blic J, *et al.* Design and recruitment for the GAP trial, investigating the preventive effect on asthma development of an SQ-standardized grass allergy immunotherapy tablet in children with grass pollen-induced allergic rhinoconjunctivitis. *Clinical therapeutics*. 2011;33:1537-1546.
282. Niggemann B, Jacobsen L, Dreborg S, *et al.* Five-year follow-up on the PAT study: specific immunotherapy and long-term prevention of asthma in children. *Allergy*. 2006;61:855-859.
283. Jacobsen L, Niggemann B, Dreborg S, *et al.* Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy*. 2007;62:943-948.
284. Valovirta E, Petersen TH, Piotrowska T, *et al.* Results from the 5-year SQ grass sublingual immunotherapy tablet asthma prevention (GAP) trial in children with grass pollen allergy. *The Journal of allergy and clinical immunology*. 2018;141:529.
285. Akdis M, Akdis CA. Mechanisms of allergen-specific immunotherapy: Multiple suppressor factors at work in immune tolerance to allergens. *Journal of Allergy and Clinical Immunology*. 2014;133:621-631.
286. Fan D-CC, Wang X-DD, Wang C-SS, Wang Y, Cao F-FF, Zhang L. Suppression of Immunotherapy on Group 2 Innate Lymphoid Cells in Allergic Rhinitis. *Chinese medical journal*. 2016;129:2824-2828.
287. Lao-Araya M, Steveling E, Scadding GW, Durham SR, Shamji MH. Seasonal increases in peripheral innate lymphoid type 2 cells are inhibited by subcutaneous grass pollen immunotherapy. *Journal of Allergy and Clinical Immunology*. 2014;134:1193-11950000.
288. Xian M, Feng M, Dong Y, Wei N, Su Q, Li J. Changes in CD4+CD25+FoxP3+ Regulatory T Cells and Serum Cytokines in Sublingual and Subcutaneous Immunotherapy in Allergic Rhinitis with or without Asthma. *International Archives of Allergy and Immunology*. 2020;181:71-80.
289. Custovic A, Simpson BM, Murray CS, Lowe L, Woodcock A, and Group NAC. The National Asthma Campaign Manchester Asthma and Allergy Study. *Pediatric Allergy and Immunology*. 2002;13:32-37.
290. Simpson A, Tan VYF, Winn J, *et al.* Beyond Atopy. *American Journal of Respiratory and Critical Care Medicine*. 2010;181:1200-1206.

291. Belgrave DC, Granell R, Simpson A, *et al.* Developmental profiles of eczema, wheeze, and rhinitis: two population-based birth cohort studies. *PLoS medicine*. 2014;11.
292. Fontanella S, Frainay C, Murray CS, Simpson A, Custovic A. Machine learning to identify pairwise interactions between specific IgE antibodies and their association with asthma: A cross-sectional analysis within a population-based birth cohort. *PLOS Medicine*. 2018;15.
293. Deliu M, Fontanella S, Haider S, *et al.* Longitudinal trajectories of severe wheeze exacerbations from infancy to school age and their association with early-life risk factors and late asthma outcomes. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2020;50:315-324.
294. Janson PC, Linton LB, Bergman EA, *et al.* Profiling of CD4+ T cells with epigenetic immune lineage analysis. *J Immunol*. 2010;186.
295. Bibikova M, Barnes B, Tsan C, *et al.* High density DNA methylation array with single CpG site resolution. *Genomics*. 2011;98:288-295.
296. Moran S, Arribas C, Esteller M. Validation of a DNA methylation microarray for 850,000 CpG sites of the human genome enriched in enhancer sequences. *Epigenomics*. 2015;8:389-399.
297. Morin AM, Gatev E, McEwen LM, *et al.* Maternal blood contamination of collected cord blood can be identified using DNA methylation at three CpGs. *Clinical Epigenetics*. 2017;9:75.
298. Opstelten R, Slot MC, Lardy NM, *et al.* Determining the extent of maternal-foetal chimerism in cord blood. *Scientific Reports*. 2019;9:5247.
299. Jarde A, Lewis-Mikhael A-M, Moayyedi P, *et al.* Pregnancy outcomes in women taking probiotics or prebiotics: a systematic review and meta-analysis. *BMC Pregnancy and Childbirth*. 2018;18:14.
300. Saccone G, Saccone I, Berghella V. Omega-3 long-chain polyunsaturated fatty acids and fish oil supplementation during pregnancy: which evidence? *The Journal of Maternal-Fetal & Neonatal Medicine*. 2015;29:1-9.
301. Chadwick R, Oconnor A. Epigenetics and personalized medicine: prospects and ethical issues. *Personalized Medicine*. 2013;10:463-471.
302. de Jager W, Rijkers GT. Solid-phase and bead-based cytokine immunoassay: A comparison. *Methods*. 2006;38:294-303.
303. Keustermans G, Hoeks S, Meerding JM, Prakken BJ, de Jager W. Cytokine assays: An assessment of the preparation and treatment of blood and tissue samples. *Methods*. 2013;61:10-17.
304. McKinnon KM. Flow Cytometry: An Overview. *Current Protocols in Immunology*. 2018;120.
305. Sandoval J, Heyn H, Moran S, *et al.* Validation of a DNA methylation microarray for 450,000 CpG sites in the human genome. *Epigenetics*. 2011;6:692-702.
306. Solomon O, MacIsaac J, Quach H, *et al.* Comparison of DNA methylation measured by Illumina 450K and EPIC BeadChips in blood of newborns and 14-year-old children. *Epigenetics*. 2018:1-10.
307. Uam JW, Holt BJ, Ron-Hay MJ, *et al.* Inhalant allergen-specific T-cell reactivity is detectable in close to 100% of atopic and normal individuals: covert responses are unmasked by serum-free medium. *Clinical & Experimental Allergy*. 1995;25:634-642.
308. Kaldjian EP, Chen G-H, Cease KB. Enhancement of lymphocyte proliferation assays by use of serum-free medium. *Journal of Immunological Methods*. 1992;147:189-195.
309. Chowdhury F, Williams A, Johnson P. Validation and comparison of two multiplex technologies, Luminex® and Mesoscale Discovery, for human cytokine profiling. *Journal of Immunological Methods*. 2009;340:55-64.
310. McKay HS, Margolick JB, Martinez-Maza O, *et al.* Multiplex assay reliability and long-term intra-individual variation of serologic inflammatory biomarkers. *Cytokine*. 2017;90:185-192.
311. Assarsson E, Lundberg M, Holmquist G, *et al.* Homogenous 96-Plex PEA Immunoassay Exhibiting High Sensitivity, Specificity, and Excellent Scalability. *PLoS ONE*. 2014;9.
312. Houseman EA, Accomando WP, Koestler DC, *et al.* DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC bioinformatics*. 2012;13:86.
313. Ritchie ME, Phipson B, Wu D, *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic acids research*. 2015;43.
314. van Iterson M, van Zwet EW, Consortium B, Heijmans BT. Controlling bias and inflation in epigenome- and transcriptome-wide association studies using the empirical null distribution. *Genome Biology*. 2017;18:19.

315. de Weerd HA, Badam TVS, Martínez-Enguita D, *et al.* MODifieR: an ensemble R package for inference of disease modules from transcriptomics networks. *Bioinformatics (Oxford, England)*. 2020;36:3918-3919.
316. Wickman M, Lupinek C, Andersson N, *et al.* Detection of IgE Reactivity to a Handful of Allergen Molecules in Early Childhood Predicts Respiratory Allergy in Adolescence. *EBioMedicine*. 2017;26:91-99.
317. Bush A. Cytokines and Chemokines as Biomarkers of Future Asthma. *Frontiers in pediatrics*. 2019;7:72.
318. Pavord ID, Beasley R, Agusti A, *et al.* After asthma: redefining airways diseases. *Lancet (London, England)*. 2018;391:350-400.
319. Booth V, Clark-Lewis I, Sykes BD. NMR structure of CXCR3 binding chemokine CXCL11 (ITAC). *Protein Science*. 2004;13:2022-2028.
320. Lejeune S, Pichavant M, Engelmann I, *et al.* Severe preschool asthmatics have altered cytokine and anti-viral responses during exacerbation. *Pediatric Allergy and Immunology*. 2020;31:651-661.
321. Lockett GA, Huoman J, Holloway JW. Does allergy begin in utero? *Pediatr Allergy Immunol*.26.
322. White GP, Watt PM, Holt BJ, Holt PG. Differential patterns of methylation of the IFN-gamma promoter at CpG and non-CpG sites underlie differences in IFN-gamma gene expression between human neonatal and adult CD45RO- T cells. *J Immunol*.168.
323. White GP, Hollams EM, Yerkovich ST, *et al.* CpG methylation patterns in the IFNgamma promoter in naive T cells: variations during Th1 and Th2 differentiation and between atopics and non-atopics. *Pediatr Allergy Immunol*.17.
324. Jacoby M, Gohrbandt S, Clausse V, Brons NH, Muller CP. Interindividual variability and co-regulation of DNA methylation differ among blood cell populations. *Epigenetics*.7.
325. Murphy A, Tantisira KG, Soto-Quirós ME, *et al.* PRKCA: a positional candidate gene for body mass index and asthma. *American journal of human genetics*. 2009;85:87-96.
326. Melén E, Himes BE, Brehm JM, *et al.* Analyses of shared genetic factors between asthma and obesity in children. *The Journal of allergy and clinical immunology*. 2010;126:631.
327. Prescott SL, Irvine J, Dunstan JA, Hii C, Ferrante A. Protein kinase Czeta: a novel protective neonatal T-cell marker that can be upregulated by allergy prevention strategies. *The Journal of allergy and clinical immunology*. 2007;120:200-206.
328. Wang C, Li H, Cao L, Wang G. Identification of differentially expressed genes associated with asthma in children based on the bioanalysis of the regulatory network. *Molecular medicine reports*. 2018;18:2153-2163.
329. Gorfu G, Rivera-Nieves J, Ley K. Role of beta7 integrins in intestinal lymphocyte homing and retention. *Current molecular medicine*. 2009;9:836-850.
330. Eigenmann PA, Tropia L, Hauser C. The mucosal adhesion receptor alpha4beta7 integrin is selectively increased in lymphocytes stimulated with beta-lactoglobulin in children allergic to cow's milk. *The Journal of allergy and clinical immunology*. 1999;103:931-936.
331. Xiong N, Fu Y, Hu S, Xia M, Yang J. CCR10 and its ligands in regulation of epithelial immunity and diseases. *Protein & cell*. 2012;3:571-580.
332. Hijnen D, Nijhuis E, Bruin-Weller M, *et al.* Differential expression of genes involved in skin homing, proliferation, and apoptosis in CD4+ T cells of patients with atopic dermatitis. *The Journal of investigative dermatology*. 2005;125:1149-1155.
333. Kanaoka Y, Maekawa A, Austen KF. Identification of GPR99 protein as a potential third cysteinyl leukotriene receptor with a preference for leukotriene E4 ligand. *The Journal of biological chemistry*. 2013;288:10967-10972.
334. Liu Z, Yelverton RW, Kraft B, Tanner SB, Olsen NJ, Aune TM. Highly conserved gene expression profiles in humans with allergic rhinitis altered by immunotherapy. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2005;35:1581-1590.
335. Martino D, Maksimovic J, Joo JH, Prescott SL, Saffery R. Genome-scale profiling reveals a subset of genes regulated by DNA methylation that program somatic T-cell phenotypes in humans. *Genes and immunity*.13.
336. Kaur S, Bansal Y, Kumar R, Bansal G. A panoramic review of IL-6: Structure, pathophysiological roles and inhibitors. *Bioorganic & Medicinal Chemistry*. 2020;28:115327.
337. Baccarelli A, Rusconi F, Bollati V, *et al.* Nasal cell DNA methylation, inflammation, lung function and wheezing in children with asthma. *Epigenomics*. 2012;4:91-100.

338. Vähämiko S, Laiho A, Lund R, Isolauri E, Salminen S, Laitinen K. The impact of probiotic supplementation during pregnancy on DNA methylation of obesity-related genes in mothers and their children. *European Journal of Nutrition*. 2018;1-11.
339. Salmond RJ, Filby A, Qureshi I, Caserta S, Zamoyska R. T-cell receptor proximal signaling via the Src-family kinases, Lck and Fyn, influences T-cell activation, differentiation, and tolerance. *Immunological Reviews*. 2009;228:9-22.
340. van Dijk SJ, Zhou J, Peters TJ, *et al*. Effect of prenatal DHA supplementation on the infant epigenome: results from a randomized controlled trial. *Clinical Epigenetics*. 2016;8:114.
341. Amarasekera M, Noakes P, Strickland D, Saffery R, Martino DJ, Prescott SL. Epigenome-wide analysis of neonatal CD4(+) T-cell DNA methylation sites potentially affected by maternal fish oil supplementation. *Epigenetics*. 9.
342. Hatzler L, Panetta V, Lau S, *et al*. Molecular spreading and predictive value of preclinical IgE response to *Phleum pratense* in children with hay fever. *The Journal of allergy and clinical immunology*. 2012;130:894-90100000.
343. Scaparrotta A, Verini M, Consilvio NP, *et al*. Sensitization to timothy grass pollen allergenic molecules in children. *Multidisciplinary respiratory medicine*. 2013;8:17.
344. Cipriani F, Mastroianni C, Tripodi S, *et al*. Diagnostic relevance of IgE sensitization profiles to eight recombinant *Phleum pratense* molecules. *Allergy*. 2018;73:673-682.
345. Scadding GW, Shamji MH, Jacobson MR, *et al*. Sublingual grass pollen immunotherapy is associated with increases in sublingual Foxp3-expressing cells and elevated allergen-specific immunoglobulin G4, immunoglobulin A and serum inhibitory activity for immunoglobulin E-facilitated allergen binding to B cells. *Clinical & Experimental Allergy*. 2010;40:598-606.
346. Varona R, Ramos T, Escribese MM, *et al*. Persistent regulatory T cell response 2 years after 3 years of grass tablet SLIT: links to reduced eosinophil counts, sIgE levels and clinical benefit. *Allergy*. 2018.
347. Durham SR, Emminger W, Kapp A, *et al*. SQ-standardized sublingual grass immunotherapy: Confirmation of disease modification 2 years after 3 years of treatment in a randomized trial. *Journal of Allergy and Clinical Immunology*. 2012;129:717-72500000.
348. Liu Y, Xing Z, Wang J, Geng C. Salivary Immunoglobulin A, E, and G4 Levels Specific to *Dermatophagoides pteronyssinus* in Allergic Rhinitis Patients Treated With Subcutaneous Immunotherapy. *American Journal of Rhinology & Allergy*. 2018;1453566771.
349. van de Veen W, Akdis M. Role of IgG4 in IgE-mediated allergic responses. *The Journal of allergy and clinical immunology*. 2016;138:1434-1435.
350. Boonpiyathad T, Satitsuksanoa P, Akdis M, Akdis CA. IL-10 producing T and B cells in allergy. *Seminars in Immunology*. 2019;44:101326.
351. Fujieda S, Zhang K, Saxon A. IL-4 plus CD40 monoclonal antibody induces human B cells gamma subclass-specific isotype switch: switching to gamma 1, gamma 3, and gamma 4, but not gamma 2. *Journal of immunology (Baltimore, Md : 1950)*. 1995;155:2318-2328.
352. Nies JH, Bär C, Schlenvoigt G, Fahlbusch B, Zwacka G, Markert UR. IL-4 supplemented B-cell cultures of allergic children show reduced IgA and IgG production in response to additional stimulation with IL-10. *Journal of investigational allergology & clinical immunology*. 2002;12:99-106.
353. Haruna T, Kariya S, Fujiwara T, *et al*. Role of whole saliva in the efficacy of sublingual immunotherapy in seasonal allergic rhinitis. *Allergology International*. 2018;68:82-89.
354. Ellis AK, Tenn MW, Steacy LM, *et al*. Lack of effect of Grastek® on birch pollen-induced allergic rhinoconjunctivitis in the Environmental Exposure Unit. *Annals of Allergy, Asthma & Immunology*. 2018.
355. Harb H, Irvine J, Amarasekera M, *et al*. The role of PKCζ in cord blood T-cell maturation towards Th1 cytokine profile and its epigenetic regulation by fish oil. *Bioscience Reports*. 2017;37.
356. James LK, Shamji MH, Walker SM, *et al*. Long-term tolerance after allergen immunotherapy is accompanied by selective persistence of blocking antibodies. *Journal of Allergy and Clinical Immunology*. 2011;127:509-51600000.
357. Shamji MH, Ljørring C, Francis JN, *et al*. Functional rather than immunoreactive levels of IgG4 correlate closely with clinical response to grass pollen immunotherapy. *Allergy*. 2012;67:217-226.
358. Custovic A, Henderson J, Simpson A. Does understanding endotypes translate to better asthma management options for all? *Journal of Allergy and Clinical Immunology*. 2019;144:25-33.

Papers

The papers associated with this thesis have been removed for copyright reasons. For more details about these see:

<http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-170575>

FACULTY OF MEDICINE AND HEALTH SCIENCES

Linköping University Medical Dissertations No. 1755, 2020
Division of Inflammation and Infection
Department of Biomedical and Clinical Sciences
Linköping University
SE-581 83 Linköping, Sweden

www.liu.se