Linköping University Medical Dissertations No 1090

Weight gain in children

-possible relation to the development of diabetes

Karina Huus

Division of Pediatrics and Diabetes Research Centre Department of Clinical and Experimental Medicine Faculty of Health Science, Linköping University SE-581 85 Linköping, Sweden



Linköping University

Linköping 2009

© Karina Huus, 2009 Cover illustration: Linus Huus ISBN 978-91-7393-729-0 ISSN 0345-0082

Papers I, III and IV have been reprinted with permission from Blackwell Publishing Ltd Paper II has been reprinted with permission from BioMed Central London Ltd

Printed in Sweden by

Distributed by

Division of Paediatrics

Department of Clinical and Experimental Medicine

Faculty of Health Sciences, Linköping University

SE-551 85 Linköping, Sweden

Phone:+46-13-22 20 00; Fax: +46-13-14 81 56

To my family

ABSTRACT

Background: The prevalence of overweight and obesity among children has increased the last decades and is now defined as a global epidemic disease by the World Health Organization. Also the incidence of type 1 diabetes has increased and there are some hypothesises that argue there is a connection between overweight/obesity and type 1 diabetes.

Aim: The general aim of this thesis was to study factors contributing to the development of overweight and obesity among children and to study possible relations to the development of diabetes.

Method: All Babies in Southeast Sweden, ABIS, is a prospective cohort study. The study includes all babies who were born in southeast Sweden between Oct 1^{st} 1997 until Oct 1^{st} 1999 and the design was to follow them up to school age in ABIS I and to follow them until 14 years in ABIS II, of the eligible 74 % entered the study. The families have answered questionnaires and biological samples were taken mainly from the children at the different time points: birth, 1 year, 2.5 years, 5 years and 8-9 years. In this thesis studies have been made including the whole cohort, but some studies have also been made involving only a part of the children.

Results: The prevalence of overweight and obesity among children in the ABIS study was 12.9% overweight and 2.5% obese at 5 years of age. One risk factor which appeared to have a great impact on the development of overweight and obesity at 5 years of age was the child's own BMI at an early age and also the heredity for overweight/ obesity and the heredity for type 2 diabetes. If the father had a university degree, the child was less likely to be obese at 5 years of age. Other factors, such as the parents' age, if the child had any siblings, and if the child lived with a single parent, did not show any significant correlation to the child's BMI at 5 years of age.

Early nutrition has been studied and no correlation could be found between breastfeeding less than 4 months and the development of overweight/obesity at 5 years of age. The parents answered questions about how frequent the child ate different food at 2.5 years and at 5 years. Intake of sweet lemonade was the only single food which was correlated to a higher BMI in 5 years old children. Porridge seemed to be protective against overweight/ obesity. In one of the studies the physical activity was measured by a step counter. The fewer steps the children were taking, the higher BMI and waist circumference they had. Low physical activity was also associated with a higher C-peptide value and decreased insulin sensitivity. Children who spent more time in front of TV/video had a higher fasting blood glucose value.

Conclusions: A strong factor for the development of overweight and obesity among children is the child's own BMI at an early age and also its heredity for overweight/ obesity and the heredity for type 2 diabetes. Early nutrition did not show any obvious correlations with overweight and obesity at 5 year old children. Low physical activity was associated with higher fasting C-peptide value and decreased insulin sensitivity. Low physical activity may cause β -cell stress which might contribute to an autoimmune process in individuals genetically predisposed to autoimmunity and, thereby, to the increasing incidence of Type 1 diabetes in children.

SAMMANFATTNING

Bakgrund: Förekomsten av övervikt och fetma bland barn har ökat under de senaste decennierna och klassas av Världshälsoorganisationen (WHO) som en global epidemi. Antalet barn som insjuknar i typ 1 diabetes har också ökat och det finns en del hypoteser som argumenterar för att det finns en koppling mellan övervikt/fetma och typ 1 diabetes.

Syfte: Den här avhandlingens syfte var att studera faktorer som bidrar till utvecklingen av övervikt och fetma hos barn och att studera om det möjligen finns en relation till utvecklingen av typ 1 diabetes.

Metod: Alla Barn I Sydöstra Sverige, ABIS, är en prospektiv kohort studie. Alla barn, som föddes mellan 1:a oktober 1997 till 1:a oktober 1999 i sydöstra Sverige, erbjöds delta. Barnen följdes sedan upp till skolåldern i ABIS I och till 14 års ålder i ABIS II. Från starten valde 74% av de tillfrågade familjerna att gå med i studien. Familjerna har besvarat frågeformulär, och biologiska prover är tagna huvudsakligen från barnen vid de olika åldrarna: födseln, 1 år, 2.5 år, 5 år och 8-9år. I avhandlingen ingår dels studier med hela ABIS kohorten, men i två av studierna deltar endast en del av barnen.

Resultat: Hos de barn som igår i ABIS studien var 12,9% överviktiga vid 5 års ålder och 2,5% var feta. En faktor som visade sig ha betydelse för utvecklingen av övervikt och fetma hos 5 år gamla barn var barnets eget BMI vid tidig ålder samt hereditet för övervikt/fetma och hereditet för typ 2 diabetes. Om föräldrarna, framförallt pappan, läst på högskola eller universitet var barnen mindre ofta överviktiga/feta. Andra faktorer som föräldrarnas ålder, om barnet hade några syskon och om barnet levde med en ensamstående förälder visade sig inte ha betydelse för utvecklingen av övervikt och fetma hos barnen. Tidig uppfödning har också studerats. Vi fann ingen korrelation mellan kort amning, dvs. mindre än 4 månader, och utvecklingen av övervikt/ fetma hos 5 år gamla barn. Föräldrarna har också fått svara på hur frekvent barnet åt olika livsmedel vid 2.5 år och vid 5 år. Saft var det enda livsmedel som enskilt hade etf samband med utveckling av övervikt och fetma. Låg fysisk aktivitet var också associerad till ett högre C-peptidvärde och minskad insulinkänslighet. Barn som tittar mycket på TV/video hade ett högre fasteblodsocker.

Konklusion: Av betydelse för utveckling av övervikt och fetma hos barn är barnets eget BMI i tidig ålder och dess hereditet för övervikt och fetma samt hereditet för typ 2 diabetes. Tidig nutrition verkar inte ha några uppenbara samband med övervikt och fetma hos 5 år gamla barn. Låg fysisk aktivitet var associerad till högt faste C-peptid och ökad insulinresistens, vilket skulle kunna stressa β -cellerna och därmed, i enlighet med β - cell stress hypotesen, kunna bidra till en ökad förekomst av typ 1 diabetes hos barn.

ORIGINAL PAPERS

This thesis is based on the following papers, conducted within the ABIS-project, which are referred to in the text by Roman numerals given below:

- I. Huus K, Ludvigsson J F, Enskär K, Ludvigsson J. Risk factors in childhood obesity findings from the All Babies In Southeast Sweden (ABIS) cohort. *Acta Paediatrica 2007; 96(9):1321-5.*
- II. Huus K, Ludvigsson J F, Enskär K, Ludvigsson J. Exclusive breastfeeding of Swedish children and its possible influence on the development of obesity: a prospective cohort study. BMC Pediatrics 2008; 9;8(1):42.
- III. Huus K, Brekke H K, Ludvigsson J F, Ludvigsson J. Relationship of food frequencies as reported by parents to overweight and obesity at 5 years. *Acta Paediatrica 2008 sep 24 (Epub ahead of print)*
- IV. Ludvigsson J, Huus K, Eklöv K, Klintström R, Lahdenperä A. Fasting plasma glucose levels in healthy preschool children: effects of weight and life style. *Acta Paediatrica 2007; 96(5):706-9.*
- V. Huus K, Raustorp A, Ludvigsson J. Physical activity, blood glucose and C-peptide in healthy school- children. Submitted

CONTENTS

INTRODUCTION
Diabetes1
Definition and diagnosis of diabetes1
Classification of diabetes mellitus2
Aetiology4
Epidemiology7
Living with T1D7
The link between child overweight/obesity and the development of T1D10
Overweight/ obesity
Definition of overweight/ obesity11
Aetiology12
Epidemiology13
Living with overweight/ obesity13
Modern Life style15
AIM OF THE THESIS
METHOD
Participants
Samples21
Procedure
Data collection22
Analyses and statistics
Ethical considerations27
RESULTS
DISCUSSION
Main methodological issues
Main results
CONCLUSIONS
Clinical implications
ACKNOWLEDEMENTS
REFERENCES

ABBREVIATIONS

ABIS	All Babies in Southeast Sweden
AOR	Adjusted Odds Ratio
BMI	Body Mass Index
CHS	Child Health Services
CI	Confidence Interval
DKA	Diabetic Keto Acidosis
FFQ	Food Frequency Questionnaire
GADA	Glutamic Acid Decarboxylase Antibodies
HOMA	Homeostasis Model Assessment
HRQOL	Health-Related Quality Of Life
IAA	Auto-Antibodies to Insulin
ICA	Islet Cell Auto-antibodies
ICC	Intraclass Correlation Coefficient
IOTF	International Obesity Task Force
IR	Insulin Resistance
OGTT	Oral Glucose Tolerance Test
OR	Odds Ratio
T1D	Type 1 Diabetes
T2D	Type 2 Diabetes
WC	Waist Circumference
WHO	World Health Organization

INTRODUCTION

The prevalence of overweight and obesity among children has increased and the World Health Organization (WHO) describes obesity as a global epidemic disease ¹. Worldwide, more people are overweight than underweight ². Also the incidence of type 1 diabetes (T1D) is increasing among children in most regions of the world ³. There are some hypotheses that argue there is a connection between overweight and T1D ^{4, 5}.

Diabetes

The disease diabetes mellitus has been known for more than 3 500 years. The word "diabetes" comes from a Greek doctor, Aretaios from Kappadokien (year 120-180), and it means that a person with diabetes drinks a lot and passes large urine quantities. Mellitus means honey-sweet and refers to the smell of the urine. First in 1921 Banting and Best discovered insulin and were able to state that lack of insulin is the cause of diabetes mellitus 6 .

Definition and diagnosis of diabetes

Diabetes mellitus comprises a group of metabolic diseases characterized by hyperglycemia resulting from defects in insulin secretion, insulin action, or both. Diabetes with chronic hyperglycemia is associated with long-term damage, dysfunction and failure of various organs, especially kidneys, eyes, heart, nerves and blood vessels ⁷.

Symptoms which are associated with diabetes are caused by hyperglycemia and include polyuria, polydipsia, weight loss and blurring of vision. Children with diabetes often present severe symptoms at onset with very high glucose levels, glucosuria and ketonuria⁷.

The diagnosis is based on measurements of plasma/blood glucose in combination with clinical symptoms as mentioned above. The diagnostic criteria are the same for both adults and children, but for children the diagnosis is made by plasma/blood glucose test without an oral glucose tolerance test (OGTT), since the symptoms usually are clear and blood glucose high. If a person does not show any symptoms, the diagnosis should be made only after

repeated plasma/blood glucose tests where the value should be in the diabetic range, or if the OGTT test meets the diagnostic criteria for diabetes⁸.

The diagnose of diabetes mellitus can be confirmed by symptoms of diabetes plus casual plasma concentration ≥ 11.1 mmol/L (casual is defined as any time of the day without regard to the time when the last meal was taken). Classic symptoms of diabetes include polyuria, polydipsia and unexplained weight loss. Fasting plasma glucose ≥ 7.0 mmol/L (fasting is defined as no calorie intake for at least 8 hours), or the 2-h post load glucose ≥ 11.1 mmol/L in capillary blood during an OGTT also gives the diagnosis. OGTT should be performed as described by WHO, using a glucose load containing equivalent of 75g anhydrous glucose dissolved in water ^{7,9}.

Classification of diabetes mellitus

The WHO has divided diabetes into four different groups: T1D, Type 2 diabetes (T2D), gestational diabetes and other specific types, depending on aetiology. Patients with any form of diabetes may need insulin at some stage of their disease but that does not in itself define the aetiological class 8 .

T1D

This form of diabetes is a result of an autoimmune mediated destruction of the β -cells in the pancreas which usually leads to absolute insulin deficiency. In some individuals this destruction of β -cells is rapid and in some individuals it is slow. Children often have the rapid form, but also in adulthood the rapid form is seen. The slow form often occurs in adults and is sometimes referred to as latent autoimmune diabetes in adults (LADA). The first manifestation of the disease can, especially among children and adolescents, be presented with ketoacidosis. Others have a modest fasting hyperglycaemia and they can, if they are exposed to infection or other stress, rapidly change to severe hyperglycaemia and/or ketoacidosis. Some individuals, especially among adults, can have a residual β -cell function for some years. However, after some eyars of the disease there is usually little or no insulin secretion as manifested by low levels of C-peptide in the plasma. In 85-90% of all individuals with T1D, where fasting diabetic hyperglycaemia is initially detected, there are markers of immune destruction, including auto-antibodies to e.g. insulin and/or to glutamic acid decarboxylase (GAD)⁸.

T2D

These patients have relative (rather than absolute) insulin deficiency. The persons are resistant to the action of insulin and at least in the beginning the patients do not need insulin treatment. This type of diabetes can be undiagnosed for many years because it does not give enough noticeable symptoms of hyperglycemia. However, these patients have an increased risk of developing macro vascular and micro vascular complications. The aetiology of this form of diabetes is at least to some extent genetic in combination with increased weight, low physical activity and perhaps stress. The insulin secretion is defective and insufficient to compensate for the insulin resistance. Weight reduction, increased physical activity and/or pharmacological treatment can increase the insulin sensivity or otherwise the own insulin secretion can be stimultated by drugs or insulin can be given ⁸.

Gestational diabetes

Gestational diabetes is when pregnant women without previously diagnosed diabetes exhibit a high blood glucose value. The women often have very few symptoms and often gestational diabetes is diagnosed through screening. Approximately 7% of all pregnancies are complicated by gestational diabetes. The main problem with gestational diabetes is its negative effects on the foetus and newborn child. Mostly this type of diabetes disappears again after pregnancy, but can sometimes reappear, usually as T2D ¹⁰.

Other specific types

This includes less common types of diabetes mellitus where the cause can be identified in a relatively specific manner. One group which includes this form is for example genetic defects on the β -cell function or insulin action. These forms of diabetes are frequently characterized by onset of hyperglycemia at a rather early age (generally before 25 years of age). They are referred to as maturity onset diabetes of the young (MODY) and are characterized by low insulin secretion with minimal or no defects in insulin action. In this type also different rare cases of diabetes are included caused by diseases of exocrine pancreas, drugs or chemicals induced or caused by infections ⁷.

Aetiology

The aetiology of T1D is in many ways unknown but the development of the disease is believed to be caused by a combination of genetic susceptibility genes, immune dysregulation and environmental factors ¹¹. The autoimmune process may start years before the clinical onset of T1D and the autoimmune process leading to diminishing insulin production ¹². Different genes for T1D provide susceptibility towards the disease and a number of environmental factors also contribute to the development of T1D (figure 1) ¹¹.



Figure 1. Scheme of T1D development. The interactions between genes and the immune system and with the environmental factors may trigger an autoimmune response leading to β -cell loss and to T1D (adapted from Harrison LC)¹³.

Genetic risk of T1D

Several different genes are involved in the pathogenesis for T1D. These genes are not necessary for developing T1D and they are not by themselves sufficient for developing T1D. There are at least four loci with strong evidence of association with T1D. The first is a human

leukocyte antigen (HLA), the most prominent of the loci to increase the risk for T1D ¹⁴. This region is located on chromosome 6p21, including the HLA-DRB1, -DQA1 and –DRQ genes. Other less important loci are: a region 5' to the insulin gene (INS) on chromosome 11p15, the CTLA-4 gene region on chromosome 2q31 and the protein tyrosine phosphatase-22 (PTPN22) ¹⁵.

Auto-antibodies against β-cell antigens

Years before clinical manifestation of T1D, immune and metabolic changes can be detected. The immune changes involve both humoral and cellular responses which persist over a prolonged period until the diagnose of T1D 16 .

Islet cell auto-antibodies

Islet cell auto-antibodies (ICA) are not specific for the β -cells because they react also with other cells in the pancreas, but they give valuable information because they could be found in 80-90% of the children who develop T1D (table 1). These auto-antibodies can be found many years before the clinical onset of T1D¹⁷. However, the ICA are not a truly independent marker, because the ICA signal is proven to be attributable to reactivity against GADA, IA-2 and IAA¹⁸.

Auto-antibodies to insulin

Insulin is the only β -cell specific auto-antigen. Insulin auto-antibodies (IAA) are usually the first auto- antibodies to appear. IAA correlates inversely with age. The older the child is, the lower level of IAA (Table 1)¹⁹.

Glutamic acid decarboxylase

GADA (Glutamic Acid Decarboxylase Antibodies) is an auto-antibody against a peptide, an enzyme, released in the islet of Langerhans in pancreas. The antigen was for the first time detected in plasma from diabetic children in Linköping ²⁰. GADA is widely used to define if healthy individuals have an increased risk to get T1D, as in ABIS, and is also used to define if diabetes is of an autoimmune type (Table 1) ²¹. In a controlled randomized trial in T1D children and adolescents with recent onset T1D it has been shown that GAD-alum treatment preserves residual insulin secretion ²². Further studies are now ongoing.

Protein tyrosine phosphatase

IA-2A, is an auto-antibody directed against the intracellular part of the IA-2 protein (tyrosine phophotase). This autoantibody is also used for screening of high risk individuals, eg in ABIS (Table 1) $^{16, 23}$

Table 1. Occurrence of auto-antibodies against pancreas in healthy individuals and in individuals with T1D.

Auto-antibodies	Shortening	Healthy	Individuals with	References
		individuals	type 1 diabetes	
Islet cell auto-antibodies	ICA	1-3 %	85 %	18, 21, 24-26
Auto-antibodies to insulin	IAA	0.7-3%	40-70 %	16, 18, 21, 24, 26-28
Glutamic acid decarboxylase antibodies	GADA	0-3 %	50-80 %	16, 18, 21, 24- 27, 29
Protein tyrosine phosphatase	IA-2A	0-2.5 %	55-80 %	24, 26, 29

To be able to predict T1D, multiple defined auto-antibodies have been studied. In a cohort of 4 505 school children, 97.5 % had no auto-antibodies, 2.3% had a single auto-antibody and 0.3% of the children had multiple auto-antibodies. Of the children who developed T1D within 8 years, all of them had multiple auto-antibodies 18 .

Clinical characteristics at diagnosis of T1D

The most classic symptoms at onset of T1D are **p**olydipsia, polyuria and weight loss. For children developing T1D the symptoms develop quite rapidly, within weeks or months. The mean duration of symptoms before diagnosis is longer the older the child is $^{30, 31}$.

When a child develops T1D with clinical symptoms, the destruction of islet β -cells causing insulin deficiency and the glucose concentration in the blood are raised beyond the renal threshold and the reabsorption of glucose in the renal is incomplete. Some glucose remains in the urine, glucosuria. This also results in an increased urine production due to the increased osmotic pressure of the urine and inhibits reabsorption of water. This leads to increased thirst,

lost blood volume will be replaced osmotically by fluid in the body cells which causes dehydration ³².

Some children can also experience blurred vision. It is the prolonged high blood glucose which causes glucose absorption, and this can lead to changes in the shape of the lenses in the eyes. This symptom often disappears when the blood glucose value is normalized ³³. Children developing T1D may initially present diabetic ketoacidosis (DKA). About 29.4% of the children who develop TID gets DKA at onset. This is an extreme state of metabolic dysregulation. The children smell of acetone and their breath is rapid and deep (kussmaul breathing). The children also have polyuria, nausea, are vomiting and have abdominal pain. DKA can lead to coma and death ^{32, 34}.

Epidemiology

The incidence of T1D is increasing among children in most regions of the world from 0.1/100 000 children below the age of 15 in low incidence countries as China and Venezuela to 32-40/100 000 in high incidence countries as Sweden, Finland, and Sardinia in Italy ^{35, 36}. The incidence rate has increased dramatically over the last decades in the western countries and in Sweden the incidence has almost doubled during the last 20 years ^{37, 38}. Boys and girls are equally affected in most populations and the increase seems to be highest in the youngest age group ³. However, according to a study of Pundziute-Lycka et al (2002) the increasing incidence of T1D in Swedish children may to some extent be a shift so the children are younger when they get the diagnosis ³⁹.

Living with T1D

The families' lived experience, when a child develops T1D, is described like an ongoing process including learning about the inevitable and about the extent. The families have to learn new things depending on what has happened and they therefore need individualized treatment in the beginning ⁴⁰. After one year the families describe that they have had an ordinary, yet different, year, but it is important that health professionals make use of the families' experience to be able to support them ⁴¹. Studies asking the child or its family about living with T1D show that the children experience the restrictions in their social life, the need to live a regular life and a feeling of being different as aspects they find difficult living with ⁴²⁻⁴⁴.

Factors related to development of T1D

Both genetic and environmental risk factors contribute to the risk of T1D in children. Different environmental factors have been suggested as trigger mechanisms of this autoimmune β -cell destruction.

Infections, such as viruses, are one of the most probable risk factors, but this has been very difficult to prove because of the time between their possible trigger effect and the manifestation of T1D. New evidence strongly suggests that enterovirus infections can be one major trigger of T1D ⁴⁵. Enteroviruses, and in particular the Coxsackie virus, have been proposed as triggers of β -cell autoimmunity and T1D ⁴⁶. An additional hint was also that it was observed that T1D was usually diagnosed in late summer and fall, this matching the seasonal pattern of enterovirus in northern Europe ⁴⁷. Another study has shown that countries with high incidences of T1D (Finland and Sweden) also have a lower frequency of enterovirus infections compared to countries where T1D is less common. This finding is congruent with the polio hypothesis, which is based on experience from another enterovirus disease, poliomyelitis, where the risk virus-induced motor neuron damage increases when the frequency of polivirus infections in the population decreases ^{48, 49}.

Another possibility is that the enterovirus infection occurs in utero or neonatally in countries with low frequency of enterovirus infections. As maternal enterovirus antibody levels are low in these countries, the child gets no protection 50 .

Even if the association between enterovirus infections and T1D has been studied in many epidemiological studies, there are still conflicting findings and the causality has not been proved. It is still possible that this association is caused by some unidentified confounding factor $^{46, 50}$.

Hygiene, Kolb and Elliot (1994) described the hygiene hypothesis. It was from the beginning introduced in the field of allergy research but could also be linked to T1D. The hygiene hypothesis suggests that due to better hygiene resulting in reduced exposure to microbial antigens early in life and by that a reduced need for a strong immune defense ⁵¹ which may lead to imbalance of the immune system with for instance autoimmune diseases or allergies as a consequence. Infections, especially in childhood, may in fact prevent or delay diseases ⁵².

Early infant feeding has also been suggested as a trigger for β -cell destruction. Short duration of breastfeeding (\leq 2months), an early introduction of cow's milk, and late introduction of gluten as well as a high consumption of milk (at the age of 1 year) have been found to be risk factors for the introduction of β -cell auto-antibodies in 2.5-years-old children ⁵³. Breast milk protects the child against infections during the first months by maternally transmitted immunity and this could also increase the child's resistance to other possible triggers of diabetes associated autoimmunity ⁵⁴. It is the cow's milk protein which has been proposed as a trigger of the activation of the immune system in the destructive process leading to T1D ⁵⁵.

Another study on pre-school children confirmed the results that long duration of breastfeeding, late introduction of bottle feeding and current consumption of cow's milk seem to reduce the risk of developing T1D 56 .

In a review of Virtanen and Knip 2003 they discuss the role of nutritional factors as potential risk factors in the development of T1D. Many different factors, such as infant feeding, cow's milk, different vitamins (C, D and E), have been reported as possibly protective against T1D. But the conclusions are that more longitudinal studies are needed from pregnancy until the children develop T1D 57 .

Maternal food consumption during pregnancy has been suggested as a trigger of β -cell destruction. Different food groups, such as potatoes, other root-vegetables, gluten-containing foods, non-gluten cereal grains, cow's milk products, fruits, vegetables, meat and fish, have been studied. It was found that an increased consumption of potatoes during the last three months of pregnancy was associated with a delayed time in the development of islet autoimmunity (IA) ⁵⁸. It is not the potato in itself but the toxic contamination of a class of streptomyces toxins that can accelerate the onset of autoimmune diabetes ⁵⁹. Also the intake of vitamin D has been studied. Mothers who continuously used cod liver oil during pregnancy, had lower risk of T1D in the offspring ⁶⁰. Measures of the intake of vitamin D in the food during pregnancy shows that it may have a protective effect on the appearance of IA in the offspring ⁶¹. However, if the mother used multivitamin supplements during pregnancy, this did not give the same results ^{60, 61}.

Psychological stress has also been suggested as a risk factor. It is known that psychological stress decreases the insulin sensitivity and increases the insulin resistance and may hence be important in the development of T1D $^{62, 63}$. Also, the mothers' experience of serious life events is associated with diabetes-related autoimmunity in children 64 .

The link between child overweight/obesity and the development of T1D

Wilkin TJ (2001) proposed the accelerator hypothesis which could be the link between child overweight/obesity and the development of T1D. This hypothesis suggests that T1D and T2D have the same origin. The increased insulin resistance is associated with the epidemic of (childhood) overweight and obesity and creates greater insulin secretory demand on the islets, leading to acceleration of β -cell destruction ⁴. The different genes affect the tempo of the β -cell loss and thus determine the age of the child when getting diabetes ⁶⁵.

An extension of the acceleration hypothesis is the β -cell stress hypothesis suggesting that any phenomenon that induces insulin resistance and/or increased insulin demand and thereby adds extra pressure on the β -cells should be regarded as a risk factor for T1D [5, 66]. Psychological stress, puberty and rapid growth could be such phenomena [64, 66].

Overweight/ obesity

Nowadays humans are among the fattest of all mammals. For humans, fat is an energy reserve. For other mammals fat mostly appears to serve as insulation from the cold. Obesity has never been a common health problem in the human history, nor has it been realistically possible for most of the people to be obese because of frequent food shortages ⁶⁶. However, in our days the prevalence of overweight and obesity among children has increased rapidly during the last 20 years. The WHO describes obesity as a global epidemic disease ¹. Also in Sweden the prevalence is increasing ⁶⁷. For the first time in 200 years life expectancy for young people may be decreased because of the prevalence of obesity ⁶⁸.

Definition of overweight/ obesity

Obesity is the result of excess body fat. Body Mass Index (BMI) is used to classify overweight and obesity in adults. BMI is the weight in kilograms divided by the square of the height in meters $(kg/m^2)^{-1}$. It is the same for both genders and for all ages of adults. WHO defines overweight in adults as BMI equal to or more than 25 and obesity as BMI equal to or more than 30. Overweight and obesity among children are defined according to International Obesity Task Force (IOTF) based on a study of Cole et al. This cut-off is equivalent to BMI in adults (overweight BMI >25 and obese BMI >30) (Figure 2). The definition is based on an international survey of six large nationally representative cross sectional growth studies ⁶⁹. To assist international comparisons in epidemiological studies, the Cole et al reference values have several strengths, they are based on large data set from different countries, the BMI cut-offs are linked to adult cut-offs , and they are simple to use. They are also consistent for children and adolescents. But there are also some concerns regarding the definition. Almost all data derive from western populations. In the population, around which the cut-offs are built, there is a variation in the prevalence of overweight/obesity, and the characteristics of the population, like the anthropometry, are unknown ⁷⁰.



Figure 2. The lines represent the BMI cut-offs within the different ages (Source: Nowicka P, Flodmark CE. Barnövervikt i praktiken- evidensbaserad familjeviktskola. P:16. Printed with permission from Studentlitteratur, Sweden)

Another way to measure overweight and obesity is Waist Circumference (WC). In adults it is well known that a more central fat distribution is associated with an increased risk of ill health ⁶⁶. However, also the prediction of health risks associated to overweight/obesity in children is improved by additional inclusion of WC ⁷¹.

Aetiology

Overweight and obesity have been seen as something which could disappear if a person changed life style ⁷². However, research has shown that it is not that simple. There are many different factors which interfere with each other in the development of overweight and obesity, such as genes, social factors, behavior, and cultural factors ⁷²⁻⁷⁴.

Childhood obesity leads to health consequences both in childhood and in adulthood. Obesity in childhood might lead to psychological consequences. Obese children are more likely to have low self-esteem and behavioral problems than non–obese children, and that risk seems to be even greater in girls than in boys. Obesity in childhood also promotes cardiovascular risks, such as dyslipidaemia and hypertension. Another clinical consequence is an increased risk of developing asthma symptoms. Also a higher risk of developing T1D has been reported ^{75, 76}.

Long term consequences of obesity in childhood have an adverse effect on the social and economic outcomes. There is some evidence that this effect might be more marked in women. Cardiovascular risk factors are higher among persons who were obese as children ⁷⁶. Obesity in childhood leads to adult obesity. Four out of five obese teenagers remain obese in adulthood ⁷⁷.

Epidemiology

During the past two decades, the prevalence of obesity has increased worldwide ¹. In a study from the US it was found that between 1999-2004 there was an increase in overweight among children, adolescents and men, while it might have stabilized among women ⁷⁸. In Europe, especially in the southern and western parts, the prevalence of overweight is higher among children ⁷⁹. Compared to other countries, Sweden has a low incidence of overweight and obesity among children, but the development during the last decades is alarming. In a study from the county of Östergötland it has been found that the prevalence of overweight among 10- year old children was 22% in both boys and girls, of which 4% and 5% respectively were obese ⁸⁰. In another Swedish study published in 2007 it was found that the obesity epidemic in children 10-11 years of age may be decreasing in urban Sweden and among girls it may possibly be reversed. From 2000/2001 until 2004/2005 the prevalence among overweight girls decreased from 19.6% to 15.9% and among obese girls from 3.0% to 2.5% ⁸¹.

Obese children, both girls and boys, are at risk of becoming obese adults. A study following children from 3 to 35 years of age showed that the higher the children's or adolescents' BMI is, the greater the risk becomes of being overweight at 35 years ⁸².

Living with overweight/ obesity

Children with overweight/obesity have a lower health-related quality of life (HRQOL) compared to normal weight children ⁸³. Especially obese children have a lower health quality of life in physical, social and school domains compared with normal weight children. The moderately obese children had similar emotional and school HRQOL, but they had lower HRQOL in the physical domain. Parents to obese children perceived their children's quality of life events lower than the children did themselves ⁸⁴.

Factors related to development of overweight/ obesity

By some estimates **Genetic variation** stands for 40-70% of the within-population variation in obesity. It is very complex to find out which genes that are involved in causing obesity in humans. Mostly there is an interaction of multiple genes, environmental factors and behaviour. There are at least 21 different genes or markers which can be linked to obesity ⁷³.

Breastfeeding. Some studies report that breastfeeding protects against obesity ⁸⁵⁻⁸⁸, while other studies show that breastfeeding does not protect against obesity ^{89, 90}. Several possible biological mechanisms for a protective effect against obesity have been discussed. Rolland-Cachera et al (1995) observed that a higher intake of protein early in life, regardless of type of feeding, was related to an increased risk of obesity ⁹¹. Infants who are breastfed have a lower intake of protein and reduced energy metabolism ⁹². Another possible biological mechanism could be that breastfed and formula-fed infants have different hormonal response to feeding. Formula feeding leads to a greater insulin response resulting in fat deposition and that the adipocytes increase in number ⁹³. Other studies suggest that infants who have been breastfed more readily adapt to new foods, such as vegetables, thus reducing the caloric density of their subsequent diets ⁹⁴.

Decreased **physical activity** has been proposed to be one of the greatest reasons for obesity ⁹⁵. TV-viewing, using computers and other sedentary occupations are also proposed as causes of obesity. However, the results are pointing in different directions ⁹⁶. Reduced television viewing and computer use may lead to a decrease in BMI, but this result might be due to the change in **energy intake** more than to the change in physical activity ⁹⁷.

There are studies which show an association between **dietary factors** and BMI ⁹⁸⁻¹⁰⁵. Especially dietary fat intake ^{98-102, 104, 105} but also the protein intake ^{105, 106} have been related to BMI. However, other studies have not confirmed these results ^{103, 107-109}.

Modern Life style

The life style for children has changed during the last decades. Their life style has changed to include less physical activity and a change of the food consumption. Food has become more affordable to a larger number of people over the last decades and also the price of food has decreased substantially in relation to the income. The means of nourishment have also changed to be a marker of life style and a source of pleasure ¹¹⁰. Inactivity, such as television viewing and computer games, is associated with increased prevalence of obesity ¹¹¹. In the last decades inactivity among children has increased and it has been found that parents prefer letting their children watch television at home rather than playing outside because then the parents can keep up with what they are doing and still be able to have an eye on the children. Children are also more often driven by their parents to school and they participate less frequently in sports and physical education ^{110, 112}.

AIM OF THE THESIS

The general aim of this thesis was to study factors contributing to the development of overweight and obesity among children and to study possible relations to the development of diabetes

Specific aims:

- To identify risk factors for overweight/obesity among children.
- To investigate if breastfeeding and early nutrition are related to overweight/obesity in 5 year -old children.
- To examine whether a modern life style with high-energy intake and low levels of physical activity influences the blood glucose value and insulin sensitivity in healthy children.

METHOD

All papers in the current thesis are based on data from the All Babies In Southeast Sweden (ABIS) project or from parts of the ABIS project. ABIS is a prospective, longitudinal cohort study aiming at studies of environmental factors affecting development of immune-mediated diseases in children. The study was initiated by Professor Johnny Ludvigsson at the Department of Molecular and Clinical Medicine. Division of Pediatrics, Faculty of Health Sciences, Linköping, Sweden.

The ABIS study is divided into two parts, ABIS I and ABIS II.

The study was designed to include all babies who were born in southeast Sweden between Oct 1st 1997 until Oct 1st 1999 and to follow them up to school age in ABIS I and to follow them until 14 years in ABIS II. In ABIS I data were collected at 4 different time points, at birth, at 1 year, at 2. 5 years and at 5 years. In ABIS II data were collected at 8-9 years and are planned to be collected at 11 years and 14 years (Figure 3).

ABIST	ABIS II
1997 1998 1999 2000 2001 2002 2003 2004 2005	2006 2007 2008 2009 2010 2011 2012 2013 2014
At-birth 0ct 97-0ct 99	8 <u>-9-yearfollo</u> w-up
1-year follow-up Oct 98- Oct 00	11-year follow-up
2.5-yearfollow-up (2-3 yrs) Oct 99-Oct 02	14 <u>-year follow-</u> up
5.5-year follow-up (5-6 yrs) Oct 02-Oct 05	

Figure 3: The time table of the ABIS projects

At each time point questionnaire data and biological samples, mainly from the children, were collected. The data in ABIS I were collected by nurses at the Child Health Services (CHS).

Data were collected in connection with the four childhood check-ups at the CHS, i.e. at 1 week, 1 year, 2-3 years and 5-6 years of age. In the 8-9 year follow-up of ABIS II the questionnaire and equipment for biological samples were sent home to the families. The parents and their children went to the nearest welfare center to take blood samples and they sent in the other biological samples by themselves together with the questionnaires.

Each questionnaire contained approx 150 questions. The questions asked in the different studies concerned BMI of the children and their parents and other questions regarded siblings and heredity for T1D and T2D among relatives. Several questions regarding food frequency, including breastfeeding, activity and inactivity. Regarded the parents, questions concerning their age, education, if they were a single parent, if they were born abroad and their smoking habits were analyzed.

Participants

All 21 700 parents-to-be in Southeast Sweden during Oct 1st 1997 and Oct 1st 1999 were invited to participate in the ABIS study. The questionnaire response rate at infant birth was 74% yielding a sample of 16 058 subjects (51.8% boys and 48.2% girls). The 1-year questionnaire was completed by 10 836 families, and 7 356 completed the 5-year questionnaire. In ABIS II the questionnaire was sent by regular post to all who participated from birth and answered that questionnaire. Parents of 3 952 children and 3 837 children themselves answered the 8-9 year questionnaire. No reminders have been sent out (September 2008, the collection is ongoing) (Figure 2).



Figure 2. Number of participating families in the different years when the data were collected (collection of data of the 8 year follow up is still ongoing, September 2008).

The representation of participants in the ABIS study is shown in table 2. Characteristics for children and parents at baseline (child birth) for all participants and for those who completed the 5 year questionnaire are presented. In addition, the table shows data from answers of the 5 year questionnaires. The numbers are based on those from whom we have data on the specific parameters.

	Baseline (n \geq 14 244)	Baseline ($n \ge 5$ 999)	5 years ($n \ge 5$ 999)
Boys/girls, %	48.2/51.8	48.1/51.9	48.1/51.9
BMI girls*	-	-	16.01 (1.77)
BMI boys*	-	-	16.09 (1.59)
BMI boys and girls*	-	-	16.05 (1.68)
Maternal age, years	29.6	29.9	34.9
Paternal age, years	32.1	32.3	37.3
BMI mother*	23.8 (3.9)#	23.7 (3.9)#	24.1 (4.0)
BMI father*	25.0 (3.0)#	25.0 (2.9)#	25.6 (3.1)
Mother born abroad, %	6.6	5.3	5.3
Father born abroad, %	7.2	5.2	5.2
University degree, mother %	31.7	34.5	39.4
University degree, father %	24.6	25.3	26.6
Single parent, %	2.1	1.2	6.6
Siblings*	1.4(0.5)	1.4(0.5)	-
Smoking, mother %	11.1	7.8	10.7
Smoking, father %	-	-	9.3

Table 2. Characteristics of the children and the parents at baseline (in all participants) and at baseline and 5 years in those who continued to participate until 5 years of age.

BMI = Body Mass Index

* Data are presented as mean values (standard deviation)

Information collected at 1-year-examination.

The baseline values for parental BMI, parental age, level of education, if the parents were born abroad, maternal smoking habits and parental civil status differ very little between the initial cohort and between those who still participated at 5 years of age. Parents in the families who dropped out of the study, tended to be slightly younger and to have a lower level of education. These parents were often smokers, were often born in a foreign country and were often a single parent at the time of birth.

Samples

Three of the papers are based on data collected from the questionnaire sent out in the ABIS study (papers I. II. III) and two of the papers (papers IV, V) are based on subsamples of the ABIS study.

Papers I, II and *III* were based on the whole ABIS cohort and questions from the birth, 1 year, 2.5 year and 5 year questionnaires. In the different studies all available data are used even if it meant that the number of participants differs in the analyses.

Paper IV was based on a subsample of the ABIS study, 127 parents and their children from 6 different pre-school units in the Linköping area participated. The children were between 5 and 7.5 years old and the mean age was 6.6 years. There were 56% girls and 44% boys. Fasting plasma glucose samples were provided from 106 children.

Paper V was based on a subsample of the ABIS study. Totally 199 children participated and questions in the 8 year questionnaire have been answered by 154 parents. Number of pedometer steps was collected from 192 children, 53.2% girls and 51.8% boys.

Procedure

In the ABIS study the at-birth questionnaire was given to the mothers when they left the hospital after the delivery. Biological samples, such as blood, urine, stool, hair from the mother and breast milk, were taken. The questionnaire was filled in and returned either immediately or to the Child Health Services at the first check up.

The questionnaires at 1 year, 2,5 years and 5 years were handed out to the parents at the Child Health Service. They were returned immediately or mailed back when completed. Biological samples, such as blood, urine and stool, were taken. Some Child Health Services reminded (by phone) those parents who had not returned the questionnaires. Also at 1 year the parents were contacted if they had not completed the full questionnaire. This was not done in the 2.5 year and 5 year questionnaires.

In study IV, information letters and questionnaires were sent out to the children and their parents by the pre-school teacher who also collected them after they had been filled in. The

fasting plasma glucose measurement was made one morning before breakfast. Also WC, weight and height were measured.

The 8 years questionnaire was sent home to the parents and their children by ordinary mail. The participants were also given the opportunity to answer the questionnaire on the home page on internet. In paper V, which is a subsample of the whole cohort, the school nurse measured height and weight. Also fasting blood samples were gathered.

Physical activity was measured by a step counter, cable tie sealed Yamax pedometers (SW-200 Tokyo, Japan) during four consecutive weekdays. The number of daily mean steps was measured over four consecutive days as recommended in order to assure reliable results and avoid reactivity ^{113, 114}. The school nurses and physical educators carried out the data collections. The pedometers were collected each day (every 24 hours) by a research assistant, who unsealed them, documented the number of daily mean steps, reset and resealed the devices and returned them to the children. The children were also asked to complete a brief survey to verify that the pedometers were worn according to the instructions during the entire time on the previous day.

Data collection

The ABIS questionnaires cover a large number of different subjects, such as medical issues, environmental factors, nutrition, psychological variables (117-196 questions). Relevant to the current thesis are questions about weight, height, WC, nutrition including breastfeeding, physical activity and some socioeconomic factors.

Weight and height

All the papers include variables with weight and height. Overweight and obesity were defined according to Cole et al ⁶⁹, age and gender adjusted. Weight and height data were received from the parents in papers I, II and III. In papers IV and V, weight and height were measured at the same time as other data were collected.

Paper I

BMI was related to different variables which could be associated with overweight/obesity, family situation (single vs. cohabitant), maternal age (\leq 34 years at child birth vs. \geq 35 years), paternal age (\leq 36 years at child birth vs. \geq 37 years), number of siblings (0 vs. \geq 1), infant birth weight (\leq 94th percentile vs. \geq 95th percentile), maternal BMI (\leq 24.99 vs \geq 25), paternal BMI (\leq 24.99 vs \geq 25), maternal and paternal education (university degree: no vs. yes). Heredity for obesity or T1D was also used.

Paper II

The questionnaire at 1 year contained questions about exclusive breastfeeding, introduction of infant formula or gruel and the introduction of cow's milk and the response alternatives were: < 1 month to >8 months. Regarding any breastfeeding there was a question in the 2.5 year questionnaire where the parents stated how many months after birth the mother stopped breastfeeding the child. Short-term exclusive breastfeeding was defined as < 4 months of exclusive breastfeeding.

Breastfeeding and the children's BMI were related to different variables such as family situation (single vs. cohabitant), maternal age (\leq 34 years at child birth vs. \geq 35 years), paternal age (\leq 36 years at child birth vs. \geq 37 years), number of siblings (0 vs. \geq 1), maternal BMI (\leq 24.99 vs \geq 25), paternal BMI (\leq 24.99 vs \geq 25), maternal and paternal education (university degree: no vs. yes).

Paper III

The questionnaire at 2.5 years contained a 30 items Food Frequency Questionnaire (FFQ) and the questionnaire at 5 years contained a 34 items FFQ regarding the children's diet. The frequency categories used were: A. daily, B. 3-5 times/week, C. 1-2 times/week, D. less than once a week and in the 5 year questionnaire, E. never. Fat content of dairy foods was not specified in the questionnaire. Foods which were expected to be associated with overweight or obesity due to high energy content (foods high in fat and/or sugar like snack foods/ candy, cheese, sausage) and foods with low energy content (vegetables, fruits, porridge) as well as foods associated to BMI in previous studies (milk, sugar containing beverages) were included in the analyses.

Paper IV

The questionnaire which was used in this paper was answered by the parents and a part of the questions was a FFQ containing 16 items regarding the children's diet. The frequency categories which were used were: A. more than 1/day, B. every day, C. 3-6 times/week, D.1-2 times/week, E. 1-3 times/month, F. more seldom. There were also questions about physical activity/inactivity, heredity for T1D and heredity for overweight/obesity. Weight, length, WC were measured and fasting plasma glucose levels were registered.

Paper V

In this paper the 8-years questionnaire from ABIS was used. Five different questions about physical activity were analysed: Number of hours spent in front of TV/video, hours spent in front of computer, hours spent in physical activity (playing, running around), hours spent reading /homework plus one further question concerning hours spent in car transportation. The scales used for these questions were 1=0.15 minutes, 2=1/2 hour, 3=1 hour, 4=2 hours, 5=3 hours, 6=4 hours, 7=5 hours, 8=6 hours, 9=7 hours or more. The questions were asked with regard to schooldays and weekend days respectively. Physical activity was measured by a step counter, height, weight and WC were also measured and blood samples, such as fasting blood glucose, HbA1c and C-peptide, were taken.

Analyses and statistics

SPSS version 11.5-14 for windows was used ¹¹⁵.

Descriptive statistics with means or medians were calculated in all the papers. In paper IV scatter plots were used to illustrate the results.

All participating children with data on each question were used in the different analyses. This explains why the number of participants differs between the analyses.

Correlations were assessed using Pearson's correlation for parametric variables and Spearman's correlation for non-parametric variables (Papers IV and V).

In paper I we used the Chi-square test. This test compares the observed numbers in each of the four categories in the contingency table with the numbers to be expected ¹¹⁶. The Chi-square test was used for analyzing heredity for overweight/obesity in paper I.

Student's t-test (independent) was used to compare mean values in the different groups. The t-test was used in paper V when we compared boys and girls and when comparing normal weight children with overweight/obese children.

Simple and/or multiple logistic regressions were used in the papers I, II and III. Multiple logistic regressions were used to investigate how a dichotomous outcome variable (dependent variable) is related to more than one exposure variable (independent variable). The logistic regression analysis was used to explore predictors and to calculate Odds Ratio (OR) with 95% confidence intervals (CI). In paper I logistic regressions were used to investigate the relationship between a number of risk factors and the risk of overweight/obese. In paper III different foods were related to normal weight and overweight/ obese children. In paper II different risk factors were related to both short and long breastfeeding and to normal weight or overweight/obese children.

Intraclass correlation coefficient (ICC) was used to be able to demonstrate the similarity between two different test methods that measure a continuous variable. In the validation of the children's height and weight intraclass correlation is used. The reason for using ICC instead of another correlation like the Pearson's was that Pearson's correlation is between two different groups, for instance between men and women. However, in the intraclass correlation the trait's mean and variance are derived from pooled estimates across all members of all groups. The intraclass correlation gives the between group variance divided by the total variance ¹¹⁶.

P-values (significance level) are used to assess the strength of evidence against the null hypothesis. This means that there is no true association in the population from which the sample was drawn. The smaller the P-value is, the stronger the evidence is against the null hypothesis ¹¹⁷. In papers I, II, IV and V the P-value < 0.05 was considered statistically significant. In paper III, due to multiple comparisons and to the risk of false-positive statistical significance, only the P-value < 0.01 was regarded as statistically significant.

Homeostatic model assessment (HOMA) are used to assessing β -cell function and insulin resistance (IR) from fasting glucose and insulin or C-peptide concentrations ¹¹⁸. In paper V, HOMA IR and HOMA β -cell were calculated.

Validity

The validity term means that the measurement shows what it claims to measure ¹¹⁹. All data in the ABIS project are given by the parents. In all papers data on weight and height are used. Therefore a validation of these data has been done. Eight different schools in the ABIS area where they had the children's records from the CHS were included in this validation. Data on 145 children at 1 year and 5 years of age were registered from the child health records and compared with the data from the parents. This validation showed a high correlation between weight and height having an intraclass correlation coefficient of P<0.001. A validation of height and weight has also been done at 8 years of age (intraclass correlation coefficient of P<0.001).

A validation has also been done in paper V where the parents answered questions about the children's physical activity and inactivity. Physical activity was measured also by a step counter/ pedometer.

Reliability

This is the precision of the measuring. Reliability measures the degree at which it is consistent and stable. A test which is stable and repeatable has reliability ¹¹⁹. P-values (significance level) are used to assess the strength of evidence against the null hypothesis. This means that there is no true association in the population from which the sample was drawn. The smaller the P-value is, the stronger the evidence is against the null hypothesis ¹¹⁷. In papers I, II, IV and V the P-value < 0.05 was considered statistically significant. In paper III, due to multiple comparisons and to the risk of false-positive statistical significance, only the P-value < 0.01 was regarded as statistically significant.

Ethical considerations

When conducting research involving children, the research should be performed according to accepted ethical guidelines and rules. The ABIS study was formally approved by the Research Ethics Committee of the Faculty of Health Science at the University of Linköping, Sweden. The ABIS study was also approved by the Research Ethic Committee of the Medical Faculty of Lund University.

Performing research including children and their families in a large scale study as the ABIS can cause ethical conflicts. Ethical questions and principles were considered in the following way with respect to the four ethical principles, respect for autonomy, beneficence, nonmaleficence and justice ¹²⁰.

Respect for autonomy

Autonomy is the capacity for self-determination. To respect an autonomous individual is to acknowledge that person's right to make choices out of his or her own values and beliefs¹²¹. In the ABIS study the parents received oral information and they were also invited to see a video film about the ABIS project before their child was born. On the first page of the questionnaire information was given that participation is voluntary. Return of the completed at-birth questionnaire and/or biological samples were considered as an informed consent to participate. In study IV blood-samples and some other measurements were taken from preschool children and they were informed that the participation was voluntary. Also in the last study V blood-samples were taken from 8 years old children. The children carried a pedometer for 4 days and some other measurements were taken. They were also informed about the voluntariness and some chose to participate only in some parts.

Beneficence

The principle is that we should do good to others ¹²¹. There are no sharp breaks between nonmaleficence and beneficence, but the beneficence principle potentially demands more than nonmaleficence because it requires positive steps to help others ¹²⁰. The main aim of the ABIS study is to study the aetiology of diabetes and other immune-mediated diseases, and, if answers can be found and thus also a cure or a way of making life easier for children with diabetes or other immune mediated diseases, we have done something good. However, children who have participated in the ABIS study can, if the research finds a way to prevent

and treat T1D and if the research shows how to identify risk individuals depending on how specific the prevention must be, get direct benefits from the study.

Nonmaleficence

This principle implies not to harm intentionally or willingly ¹²¹. In the ABIS study the parents and children answered questionnaires, and biological samples, such as blood-samples, were also taken. It could always be discussed if the child is harmed or not. When the child are small, the parents make the decision, but the older the children are, the more they can decide for themselves. The parents and the children have got information about that their participation is voluntary, and sometimes, when a child has not consented to take some biological samples, the parents have nevertheless answered the questionnaire.

Justice

In general, justice means fairness, for example ensuring equality of opportunity for all individuals and equality of the outcome for different groups in society ¹²¹. In ABIS the opportunity to participate in the study was the same for all mothers in the southeast of Sweden who visited the Maternal Health Services. All participating families have answered the same questionnaires and the same biological samples have been asked for. However, papers IV and V are based on subsamples of the ABIS where the opportunity to participate has been given to a small group in a pre-school class or in an ordinary school class.

The ABIS study is a large prospective cohort study and in studies like this there are often ethical concerns about the risk of creating distress and anxiety. Therefore, in the ABIS study there have been several papers on ethical concerns. The first study about ethical concerns was made in 2002 by Gustafsson-Stolt et al. and thereafter a number of studies have been made. All these studies, either based on interviews or questionnaires, show that a majority believed that a study like the ABIS was good and they wanted to participate because they felt they could do something for the research. The mothers who did not want to participate were concerned about putting the child through frequent blood testing and some felt that the project involved some kind of "experimentation" ¹²². Gustafsson-Stolt et al (2003) found that the parents ´ attitudes towards the project were prominently positive. The parents were positive to research screening and felt that this kind of research is important ¹²³.

RESULTS

All papers include BMI of the children and in table 3 the BMI of the whole cohort is presented. In papers IV and V, subsamples of the whole cohort are used.

Table 3. BMI distribution in the ABIS study (the whole cohort)

	Overweight	Obesity
Children at 2. 5 years	12.9%	2.5%
Children at 5 years	12.9%	4.3%
Children at 8-9 years	12.0%	2.3%
*Mothers	23%	7%
*Fathers	41%	6%
■ Mothers	23%	8.5%
■ Fathers	46%	7.5%

*1 year after the child's birth

■ 5 years after the child's birth

Paper I

In paper I a multiple logistic regression with the children's BMI > 25 and > 30, age and gender adjusted according to Cole et al ⁶⁹ as our dependent variable and different risk factors as independent variables was made. The independent variables include the parents' age, parents' education, if the child had any siblings, and if the child lived with a single parent. The analysis shows that children whose fathers had a university degree were less likely to be obese at 5 years of age (adjusted odds ratio (AOR) = 0.74; 95% CI= 0.60-0.91; p= 0.005) and the result was similar when BMI >30 (age and gender adjusted) was used as an dependent variable. A borderline association was also found for maternal education and children's BMI>25 (AOR= 0.84; 95% CI= 0.70-1.00; p=0.054). The other variables in the analysis were not significantly associated.

In another multiple regression analysis of the children's BMI as a dependent variable the children's birth BMI > 95th percentile and the parents' BMI >25 were used as independent variables. A high birth BMI was positively associated with BMI at 5 years of age (AOR= 2.25; 95% CI= 1.68-3.02; p = < 0.001) and also the parental BMI was positively associated with the children's BMI at 5 years of age (high BMI in mothers: AOR = 1.97; 95% CI = 1.67-2.32; p < 0.001; and high BMI in fathers: AOR= 1.96; 95% CI= 1.64-2.33; p < 0.001). Similar results were found when the parents BMI < 30 was used as an independent variable.

Heredity: It was found that children, who had heredity for T2D, had a higher BMI (p<0.001; $N= 6\ 213$) at 5 years of age and children, who had heredity for overweight, also had a higher BMI (p<0.001; $N= 6\ 213$) at 5 years of age.

Paper II

In paper II it was found to be a weak association between short-term exclusive breastfeeding and obesity in 5 year old children (simple logistic regression: OR=1.44; 95% CI=1.00-2.07; P=0.050). However, when adjusting for other variables, such as parental age, single parent, parent born abroad, parents' smoking habits and the parents' education, the association between short-term exclusive breastfeeding and obesity in 5 year old children did not exist any longer. The only association found in the multivariate analysis was maternal smoking and obesity at 5 years of age (AOR= 1.72; 95% CI= 1.00-2.93; P=0.048).

At the age of three months 78.4% of the children were exclusively breastfed and the median age for introduction of infant formula or gruel was 5 months. The median duration of any breastfeeding was 8 months. Any breastfeeding was also associated with the parents' age. Older mothers breastfed for a longer time than younger mothers (R2=0.028, P<0.001).

Short-tem exclusive breastfeeding (less than 4 months) was in a multiple logistic regression associated with maternal BMI> 30 (AOR=1.07; 95% CI=1.05-1.09; P<0.001) and maternal smoking (AOR=1.43; 95% CI= 1.05-1.95; P= 0.023). It was also more common with short exclusive breastfeeding among single mothers (AOR= 2.10; 95% CI= 1.43-3.09; P<0.001). It was less common with short exclusive breastfeeding among parents with a university degree (mother: AOR=0.74; 95% CI= 0.61-0.90; P=0.003, father: AOR=0.73; 95% CI= 0.58-0.92; P=0.008).

Paper III

In an analysis where overweight/obese children at 5 years of age and food frequencies at 2.5 years were studied, the only food that was positively associated with (contributed to) overweight/obesity was the consumption of cheese (AOR=0.070; 95% CI= 0.55-0.90; P= 0.005). Negatively associated with (protect against) overweight/obesity was porridge (AOR= 0.57; 95% CI= 0.39-0.84; P= 0.004), fried potatoes /pommes frites (AOR=0.77; 95% CI= 0.54-0.92; P= 0.005) and cream /crème fraiche (AOR=1.19; 95% CI= 1.00-1.41; P= 0.046).

Based on previous results from this population 124 , analyses were performed adjusted for parental BMI, parental education and heredity for T1D and T2D. When adjusting for these risk factors, the frequency in intake of fried potatoes/pommes frites (AOR=0.75; 95% CI=0.62-0.92; P= 0.006) and porridge (AOR=0.55; 95% CI= 0.36-0.85; P=0.007) remained significantly negatively associated with overweight/obesity.

In another analysis where overweight/obesity at 5 years of age and food frequencies at 5 years were studied. The intake of chocolate (AOR=0.76; 95% CI=0.65-0.90; P= 0.001) and lemonade (AOR=1.38; 95% CI=1.12-1.70; P= 0.003) were positively associated, whereas cream/crème fraiche (AOR=1.23; 95% CI=1.06-1.43; P= 0.007), pastries (AOR=0.43; 95% CI=0.24-0.79; P=0.007), chocolate (AOR=0.76; 95% CI=0.65-0.90; P= 0.001) and candy (non-chocolate) (AOR=1.46; 95% CI=1.14-1.87; P= 0.003) were negatively associated with overweight/obesity at 5 years of age. When adjusting for known risk factors only the frequency in intake of candy (non-chocolate) AOR=1.60; 95% CI=1.22-2.12; P=0.001) remained negatively associated with overweight/obesity.

Paper IV

In paper IV, playing outdoors was associated with a higher degree of physical exercise (p<0.05, $R^2=0.082$) and it was also correlated to a higher BMI (p<0.001, $R^2=0.228$). Children who watched TV often had a higher BMI (p<0.01, $R^2=0.228$) and the WC was correlated with how many hours per week the children played outdoors (p<0.01, $R^2=0.151$). BMI was also correlated to the consumption of sweet lemonade (p<0.05, $R^2=0.041$). In paper IV no correlations were found between fasting plasma glucose levels and physical activity, eating habits or the children's BMI.

Paper V

When the children carried a pedometer for 4 days it was found that boys took more steps than girls, even though boys were said to spend more time in front of the computer (paper IV). The fewer steps the children were taking, the higher BMI (p=0.019) and WC (p= 0.018) they had. C-peptide (P=0.044), HOMA IR (P= 0.046) and HOMA β -cell (P=0.041) were also significantly correlated to how many daily mean steps the children were taking. A higher HOMA IR was also correlated to a larger WC (P=0.041).

How many hours per day the child spent in front of TV / video were correlated to larger WC (school days: P= 0.033 weekend days: P= 0.016). Children who spent more hours per day in front of TV / video were also correlated to higher fasting blood glucose values (school days P=0.016). Children who spent long time sitting in a car had higher fasting blood glucose values (P= 0.037). The more time the children spent in front of TV/DVD on schooldays, the less time they spent on physical activity in the weekends (P= 0.045), Table 4.

	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.	13.	14.	15.	16.
MI																
'aist circum ference	.797**															
asting blood glucose	<i>019</i> .	.079														
asting C-peptide	.149	.175	.028													
bA1C	062	126	028	.016												
OMA IR	.172	.221*	.082	*166.	600.											
OMAβ-cell	.040	.072	.468**	.831**	.030	.805**										
aily mean steps	185*	.212*	.095	205*	062	210*	.239*	,								
iours spent in front of TV deo/DVD (S)	.136	.198*	233*	.185	.072	,174	077	.051								
iours spent in front of TV deo/DVD (W)	.133	.206*	.063	.064	.084	.041	075	-070	.623**							
ours spent in front of computer (S)	027	.055	077	.073	153	.037	.087	.117	.163*	.232*	ı					
ours spent in front of computer (W)	020	.032	002	.016	167	006	.019	.109	.109	.193*	.749**					
iours spent in physical activity laying, running around) (S)	059	135	.015	004	600.	017	008	.085	033	004	-097	083				
lours spent in physical activity laying, running around) (W)	124	102	059	.029	.610	600.	.030	030	164*	076	127	128	.705*			
ours spent reading/ homework (S)	-000	.119	-009	.042	218*	.046	.035	046	.091	.071	.154	.072	018	600.		
ours spent reading/ homework (W)	031	.089	.043	.100	194*	.131	.094	121	042	007	.073	.021	026	.063	.663**	
ours spent in the car (per week)	031	.050	.202*	.037	.047	600.	134	075	127	040	206*	157	.116	.145	028	.112
	sting blood glucose sting C-peptide bAIC OMA IR OMAβ-cell aily mean steps uurs spent in front of TV keo/DVD (S) ours spent in front of TV keo/DVD (S) ours spent in front of computer (S) ours spent in front of computer (W) ours spent in physical activity laying, nunning around) (S) ours spent in physical activity laying, nunning around) (S) ours spent in the car (per week) ours spent in the car (per week)	sting blood glucose	sting blood glucose 079 079 079 sting C-peptide 149 175 bAIC -062 -126 bAIC -062 -126 oMA IR 172 $-221*$ OMAβ-cell 040 072 ally mean steps $175*$ $-212*$ us spent in front of TV 135 -079 055 ours spent in front of TV 135 -027 055 ours spent in front of computer (%) -020 032 ours spent in physical activity -124 -102 ours spent in physical activity -031 089 ours spent in the car (per week) -031 030	sting blood glucose 079 079 $ 149$ 175 028 bMC -062 -126 -028 bMC MA IR 172 -062 -126 -028 $0MA$ IR 172 -07	sting blood glucose 079 079 $-$ sting C-peptide 149 175 028 $-$ bMIC -062 -126 028 016 bMIC -062 -126 028 016 bMIC -062 -126 028 016 bMA R 172 $221*$ 082 $991*$ OMA β-cell 040 072 $468*$ $831**$ oMA β-cell 040 072 $468*$ $831**$ oMA β-cell 040 072 $468*$ $831**$ oWA β-cell 040 072 $468*$ $831**$ oWA β-cell 040 072 $205*$ 006 006 own steps $-185*$ $-122*$ 053 006 063 own steps -133 $206*$ 053 006 073 006 own steps -020 023 -002 016 073 006 own steps -133 005 -135	sting blood glucose 079 079 $-$ sting C-peptide 149 175 028 $-$ bMIC -062 -126 028 $-$ bMIC -062 -126 028 016 $-$ bMIC -062 -126 028 016 $-$ bMIC -062 -126 028 016 $-$ bMIA 172 2214 082 9214 030 OMAAcell 040 072 4684 8314 030 off proved -785 2072 025 007 002 off proved -785 2072 025 002 012 or spent in front of computer (W) -022 002 016 -153 ours spent in front of computer (W) -022 002 016 -167 ours spent in front of computer (W) -022 012 003 004 009 ours spent in front of computer (W) -020 -1132 012 003 <td>sting Creptide 172 073 079</td> <td>sting blood glucose 7 20 2 20 2 20 2 20 2 20 2 20 2 20 2 2</td> <td>sting blood glucose 079 .75 .28 . Alt</td> <td>sting blood glucose 079 .77 .224 .28 . MAT R .72 .224 .28 .016 . MAT R .72 .224 .28 .016 . MAT R .72 .224 .28 .016 . MAT R .72 .224 .27 .297 . MAT R .72 .217 .297 .010 .299 . MAT R .72 .218 .297 .010 .010 . MAT R .72 .218 .297 .010 .010 .011 .051 . MAT R .717 .012 .012 .012 .012 .010 .011 .013 . MAT R .718 .233 .012 .012 .012 .011 .011 .013 . MAT R .718 .233 .012 .012 .012 .011 .011 .013 .013 .013 .013 .013 .013</td> <td>sting blood glucose 109 175 028</td> <td>sting blood glucose079079079.sting blood glucose.149.175.028.0bMC.402.126.028.0.bMC.402.216.028.0.bMC.402.216.028.0.bMC.402.216.028.0.bMC.402.217.028.005.209*bMC.416.72.214.002.209*bMA.412.214.072.209*.072sty mean steps.418*.212*.092.209*.071sty mean steps.418*.219*.214.017.011.013sty mean steps.418*.213*.185.072.219*.071.011sty mean steps.116.012.012.012.011.012.012.013sty mean steps.012.013.014.017.011.103.103.104sty mean steps.012.012.012.012.010.103.103.103sty steps.012.013.014.017.010.019.103.010.104sty steps.012.013.012.014.011.010.103.010.103sty steps.014.011.014.011.010.103.010.103.104sty steps.012.012.012.012.010.011.0</td> <td>sing blood gluceseDPDPsing C-peptide149175028MIC-002-126028016MIC-002-126028016MIC-002-126028016MIC-002-126028016MIC-012221*080072MIC-17221*081MIC-17221*081MIC-17221*092MIC-17MIC-17MIC-17MIC-17MIC-17MIC-17MIC-17MIC-17MIC-18MIC-18MIC-13MIC-13MICMIC<t< td=""><td>sing blood glucese9797.sing Cpephide149175028.bMC062.126028.028.016.bMC172221*082.016.DMA IR172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*.022.016.010.DMA Facell171.016.221*.017.016.DMA Facell171.016.210*.002.210*.002DMA Facell136.210*.023.012.210*.010DMA Facell136.210*.023.014.017.016DMA Facell136.210*.023.014.017.016DMA Facell.136.210*.023.014.017.016Lav DVUD (%).131.206.013.014.017.016Lav DVUD (%).131.016.013.017.016.025*.017Lav DVUD (%).131.016.016.017.016.024*.014Lav DVUD (%).131.016.016.017.016.025*.014Lav DVUD (%).131.016.016.016.017.016.019.016Lav DVUD (%</td><td>ang Cpopide 07</td><td>and glood glucose079079</td></t<></td>	sting Creptide 172 073 079	sting blood glucose 7 20 2 20 2 20 2 20 2 20 2 20 2 20 2 2	sting blood glucose 079 .75 .28 . Alt	sting blood glucose 079 .77 .224 .28 . MAT R .72 .224 .28 .016 . MAT R .72 .224 .28 .016 . MAT R .72 .224 .28 .016 . MAT R .72 .224 .27 .297 . MAT R .72 .217 .297 .010 .299 . MAT R .72 .218 .297 .010 .010 . MAT R .72 .218 .297 .010 .010 .011 .051 . MAT R .717 .012 .012 .012 .012 .010 .011 .013 . MAT R .718 .233 .012 .012 .012 .011 .011 .013 . MAT R .718 .233 .012 .012 .012 .011 .011 .013 .013 .013 .013 .013 .013	sting blood glucose 109 175 028	sting blood glucose079079079.sting blood glucose.149.175.028.0bMC.402.126.028.0.bMC.402.216.028.0.bMC.402.216.028.0.bMC.402.216.028.0.bMC.402.217.028.005.209*bMC.416.72.214.002.209*bMA.412.214.072.209*.072sty mean steps.418*.212*.092.209*.071sty mean steps.418*.219*.214.017.011.013sty mean steps.418*.213*.185.072.219*.071.011sty mean steps.116.012.012.012.011.012.012.013sty mean steps.012.013.014.017.011.103.103.104sty mean steps.012.012.012.012.010.103.103.103sty steps.012.013.014.017.010.019.103.010.104sty steps.012.013.012.014.011.010.103.010.103sty steps.014.011.014.011.010.103.010.103.104sty steps.012.012.012.012.010.011.0	sing blood gluceseDPDPsing C-peptide149175028MIC-002-126028016MIC-002-126028016MIC-002-126028016MIC-002-126028016MIC-012221*080072MIC-17221*081MIC-17221*081MIC-17221*092MIC-17MIC-17MIC-17MIC-17MIC-17MIC-17MIC-17MIC-17MIC-18MIC-18MIC-13MIC-13MICMIC <t< td=""><td>sing blood glucese9797.sing Cpephide149175028.bMC062.126028.028.016.bMC172221*082.016.DMA IR172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*.022.016.010.DMA Facell171.016.221*.017.016.DMA Facell171.016.210*.002.210*.002DMA Facell136.210*.023.012.210*.010DMA Facell136.210*.023.014.017.016DMA Facell136.210*.023.014.017.016DMA Facell.136.210*.023.014.017.016Lav DVUD (%).131.206.013.014.017.016Lav DVUD (%).131.016.013.017.016.025*.017Lav DVUD (%).131.016.016.017.016.024*.014Lav DVUD (%).131.016.016.017.016.025*.014Lav DVUD (%).131.016.016.016.017.016.019.016Lav DVUD (%</td><td>ang Cpopide 07</td><td>and glood glucose079079</td></t<>	sing blood glucese9797.sing Cpephide149175028.bMC062.126028.028.016.bMC172221*082.016.DMA IR172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*082.016.DMA Facell172.221*.022.016.010.DMA Facell171.016.221*.017.016.DMA Facell171.016.210*.002.210*.002DMA Facell136.210*.023.012.210*.010DMA Facell136.210*.023.014.017.016DMA Facell136.210*.023.014.017.016DMA Facell.136.210*.023.014.017.016Lav DVUD (%).131.206.013.014.017.016Lav DVUD (%).131.016.013.017.016.025*.017Lav DVUD (%).131.016.016.017.016.024*.014Lav DVUD (%).131.016.016.017.016.025*.014Lav DVUD (%).131.016.016.016.017.016.019.016Lav DVUD (%	ang Cpopide 07	and glood glucose079079

Table 4. Correlations within physical variables and questionnaire data

Pearsons orrelation Spearman correlation *significant level 0.05 ** significant level 0.01 (S) = school days (W) weekend days

DISCUSSION

Main methodological issues

All 21 700 parents-to-be in Southeast Sweden during Oct 1st 1997 until Oct 1st 1999 were invited to participate in the ABIS study. The questionnaire response rate at birth was 16 058 subjects. No selection was made by the researchers. At 1 year 10 836 families answered the questionnaire, at 5 years 7 356 families answered the questionnaire and at 8 years 3 952 parents and 3 837 children answered the questionnaire. One weakness of the ABIS study is the drop-out rate. It has been discussed how much loss that can be acceptable in prospective cohort studies, and some researchers have tried to put a number on it. Kristman et al propose that acceptable follow-up rates are 60%-70% ¹²⁵, others suggest 80% ¹²⁶. However, in a new study of Fewtrell et al ¹²⁷ they suggested that the most important is to give unambiguous information about the cohort throughout the study, to discuss potential bias, if the drop-out has affected the generalization of the findings and that appropriate sensitivity analyses are used ¹²⁷. In the papers in this thesis, the initial cohort is compared with the cohort which is still participating in the 5 year follow-ups in those variables that were used. There is no selection bias with regard to the parents' BMI and almost none with regard to the parents' age. However, there is a slight overrepresentation of parents with a high education, fewer parents born abroad, fewer single parents and fewer smoking parents at the 5 year follow-up.

All data in the ABIS study are self reported. Therefore, there is a possibility that some variables, like the food frequency consumption, to some extent can be underreported. The main objective of the ABIS study is to examine risk factors for T1D, and by tradition there is a belief among people that T1D is caused by or related to sugar consumption. Therefore, it can be an underreporting in these variables. There has also been an increased awareness of obesity among children and, therefore, some parents might have underreported the intake of fat ¹²⁸. However, the weight and height data were validated against those recorded at the Child Health Service. Eight different schools in the ABIS area where they had the children's records from the Child Health Service were included in this validation. The weight and height data on 145 children at 1 year and 5 years were registered from the child health records and compared with data from the parents. This validation showed a high correlation between

weight and height, having an intraclass correlation coefficient of P<0.001. Another study also shown that parents are good at estimating their children's weight ¹²⁹.

Main results

The incidence of T1D is increasing ³, and also the incidence of overweight and obesity among children is increasing ^{1, 67}. The reason for the increase in T1D is unknown, but one hypothesis discussed in recent years is the β -Cell Stress Hypothesis. This means that any phenomena which induce insulin resistance and/ or increased insulin demand and thereby add extra pressure on the β -cells should be regarded as a risk factor for T1D ^{5, 130}. These phenomena could be psychological stress, puberty and rapid growth ^{64, 130}. In this thesis phenomena which could contribute to the development of overweight/obesity in children and thereby to the development of diabetes have been studied (Figure 3).



Figure 3. Scheme of phenomena which could contribute to the β -cell stress and maybe to T1D.

Heredity

The parents' BMI is one factor associated with the children's BMI. There is an interaction of multiple genes which could be linked to obesity ⁷³, and in paper I it was found that children with heredity for overweight and T2D had a higher BMI at 5 years of age. Also the heredity for T2D are associated to a higher BMI at 5 years of age as shown in paper I.

Socioeconomic factors

Parents' education

The parents' education was the only socioeconomic factor in this study which was found to influence the children's weight (paper I). This is also confirmed in other studies ^{131, 132} where it also has been found that socioeconomic factors, such as smoking, education, and family income, influence the weight development in children.

Manner of living

Early nutrition

One factor which has been studied is breastfeeding. There are theories that breastfeeding protects against obesity. One theory is that children who are breastfed have a lower intake of protein and by that less risk for developing obesity 91 . The mothers in the ABIS breastfed for a long period, 78.4% breastfed exclusively for at least three months. In paper II no association was found between exclusive breastfeeding (< 4 month) and the children's BMI at 5 years when adjusting for other factors.

In paper III food frequencies are used to find associations between early nutrition and the child's BMI at 5 years of age. The food frequency questionnaire was one way to measure how often the child consumed a certain type of food. It gives a rather rough estimation and says nothing about portion size. However, the advantage with the food frequency questionnaire in the ABIS study is that it reflects the dietary habits for a long time even if it is influenced by attitudes and knowledge of the parents. Food frequencies do not give any simple explanation for overweight and obesity at 5 years. Porridge at 2.5 years may be

protecting against overweight/obesity and lemonade may contribute to overweight/obesity at 5 years. These results are confirmed by a study from the US where it was found that consumption of sugar-sweetened drinks is associated with obesity in children. The risk for overweight/ obesity increased by each additional glass of sugar-sweetened drink the child consumed every day ¹³³.

Physical activity/ inactivity

Physical activity is measured by an objective method; step counters (paper V) and by questionnaires to the parents (paper IV and V). This is associated not only to the child's BMI, but also to their WC and to blood glucose control and insulin sensitivity. When measuring physical activity, it was found that lower physical activity is associated with higher BMI and larger WC. Another finding was that children who were more physically inactive, spent more time in front of TV/video, and had more time spent in the car have higher fasting blood glucose. Those with low physical activity measured by step counters also have higher fasting C-peptide and increased HOMA IR index which indicate insulin resistance.

A life style with high energy intake and low levels of physical activity may in fact already at these low ages increase the β -cell stress and could, according to the β -cell stress hypothesis ⁵, contribute to an increasing incidence of T1D in children.

Individual factors of the child

BMI in early age

One of the purposes of this thesis was to study factors contributing to the development of overweight and obesity among children. One of the most important factors is the children's own BMI. Early overweight/obesity has to be taken seriously. Overweight and obesity at 1 and 2.5 years of age increase the risk of being overweight at 5 years (paper I) and obesity in childhood leads to obesity later in adult life ⁷⁷.

Overweight and obesity among children are increasing when comparing the children in the ABIS study with children born a few years earlier in the same area ¹³⁴. There are few effective treatments for childhood obesity ^{135, 136} and, therefore, the need of identification of risk factors is important to be able to prevent childhood obesity.

CONCLUSIONS

This thesis has shown:

- A strong factor for the development of childhood overweight and obesity is the child's own BMI at a very early age, the parents' BMI and the child's heredity for T2D and the child's heredity for overweight.
- If the parents have a low education (no university degree) this is a risk factor for childhood overweight and obesity.
- Physical activity is associated with the children's BMI.
- Low physical activity is associated with higher fasting C-peptide and decreased insulin sensitivity, which may stress the β-cells and could, according to the β-cell stress hypothesis, contribute to an increasing incidence of T1D in children.
- Early nutrition did not show any obvious relation to overweight/ obesity at 5 years of age. The foods that showed a correlation in the early nutrition of the child were sweet lemonade which could cause overweight/obesity and porridge which was found to protect against overweight/obesity.

Clinical implications

These factors, the child's BMI, the parents' BMI, heredity for T2D and obesity together with the parents' education, can be identified at the CHS and, maybe, it could be possible to start prevention early. Prevention could be advice about the importance of physical activity as a prevention of later overweight/ obesity and also advice about the importance of early nutrition, such as the intake of sweet lemonade. The advice should be about how important it is with physical activity and also how important it is to avoid physical inactivity. At the CHS, advice could be given on limiting the children's screen time. If the parents could do that, the children spontaneously would be more physically active and as a result overweight and obesity would be prevented. Also, when the children get a little older, it is important to promote physical activity in school from the beginning and to give advice about nutrition in school, not only regarding the school meals but also other intakes at school.

ACKNOWLEDEMENTS

I would like to express my gratitude to all who have helped, supported and encouraged me in my work. I am especially grateful to:

All families that participate in the ABIS study - a prospective cohort study that included repeated biological samples and extensive questionnaires for the parents.

Johnny Ludvigsson, Professor, my main supervisor, for inspiring me to become a researcher, for supporting me during this work, and for interesting scientific discussions.

Jonas Ludvigsson, PhD, my co-supervisor, for scientific guidance especially in the field of statistics, for constructive criticism and for encouragement.

Karin Enskär, PhD, my co-supervisor, for guiding me in to the world of research, for interesting scientific discussions and for always being there when needed, and for being a very good friend.

Hilde Brekke, PhD, my co-author in study III, for stimulating discussions and guidance in the field of nutrition and also for being a good friend.

Anders Raustorp, PhD, my co-author in study V, for constructive comments and interesting discussions.

Ann-Christine Gilmore-Ellis, for help and support with administration and for always being kind.

Iris Franzen and Christina Larsson, ABIS nurses, for their practical work in the ABIS project.

Annelie Sepa, PhD, for checking the ABIS data and for constructive criticism and for encouragement.

Anna-Greta Huus, mother in law, for language checking, support and encouragement.

All doctoral students and colleges at the department of pediatrics, Linköping University, for support, helpful criticism during seminaries, and stimulating discussions during this period.

The colleagues and friends at the University College of Health Sciences, Jönköping, who inspired me to begin and encouraged me to carry on. Thanks for all support.

The Child group at Jönköping University for helpful criticism during seminaries, for support and friendship.

Fredrik, my husband for being understanding and encouraging.

Hampus, Linus and Kasper our wonderful children who bring us happiness.

At last, but not at least my family, my father and the rest of my family, for support and for believing in me. All my friends outside the scientific world who are very important to me and who all have supported and encouraged me during these years.

This work was generously supported by the JDRF-Wallenberg Foundation, the Swedish Medical Research Council (MFR, Vetenskapsrådet), the Swedish Child Diabetes Foundation (Barndiabetesfonden), the Swedish diabetes Association, Novo Nordisk Foundation, Swedish Milk (Svensk Mjölk), The Health Research Council of South-east Sweden (FORSS), the Swedish Diary Association R&D, and the Diabetes Research Centre in Linköping.

REFERENCES

- 1. WHO. Obesity: preventing and managing the global epidemic. *Report from WHO consultation. World health organisation.* 2000;Technical report series 894.
- 2. Popkin BM. The world is fat. *Sci Am.* Sep 2007;297(3):88-95.
- **3.** Soltesz G, Patterson CC, Dahlquist G. Worldwide childhood type 1 diabetes incidence--what can we learn from epidemiology? *Pediatr Diabetes*. Oct 2007;8 Suppl 6:6-14.
- **4.** Wilkin TJ. The accelerator hypothesis: weight gain as the missing link between Type I and Type II diabetes. *Diabetologia*. Jul 2001;44(7):914-922.
- 5. Ludvigsson J. Why diabetes incidence increases--a unifying theory. *Ann N Y Acad Sci*. Oct 2006;1079:374-382.
- 6. Pickup J. WG. Textbook of diabetes, Second edition Blackweel Science, United Kingdom. 1997;1 (Chapter 1, The history of diabetes mellitus).
- 7. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. Jan 2007;30 Suppl 1:S42-47.
- **8.** WHO. Definition, Diagnosis and Classification of Diabetes Mellitus and it's complications: Report of a WHO consultation. *Geneva, World Health Org.* 1999.
- **9.** WHO/IDF. Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia. *Report of a WHO/IDF Consultation*. 2006.
- **10.** ADA. Gestational Diabetes Mellitus, American Diabetes Association. *Diabetes Care*. 2003;26 (Supplement 1).
- **11.** Atkinson MA, Eisenbarth GS. Type 1 diabetes: new perspectives on disease pathogenesis and treatment. *Lancet*. Jul 21 2001;358(9277):221-229.
- **12.** Knip M. Natural course of preclinical type 1 diabetes. *Horm Res.* 2002;57 Suppl 1:6-11.
- **13.** Harrison LC. Vaccination against self to prevent autoimmune disease: the type 1 diabetes model. *Immunol Cell Biol*. Feb 2008;86(2):139-145.
- 14. Onengut-Gumuscu S, Concannon P. Recent advances in the immunogenetics of human type 1 diabetes. *Curr Opin Immunol*. Oct 2006;18(5):634-638.
- **15.** Rich SS, Concannon P, Erlich H, et al. The Type 1 Diabetes Genetics Consortium. *Ann N Y Acad Sci.* Oct 2006;1079:1-8.
- **16.** Leslie RD, Atkinson MA, Notkins AL. Autoantigens IA-2 and GAD in Type I (insulin-dependent) diabetes. *Diabetologia*. Jan 1999;42(1):3-14.
- **17.** Taplin CE, Barker JM. Autoantibodies in type 1 diabetes. *Autoimmunity*. Feb 2008;41(1):11-18.
- **18.** LaGasse JM, Brantley MS, Leech NJ, et al. Successful prospective prediction of type 1 diabetes in schoolchildren through multiple defined autoantibodies: an 8-year follow-up of the Washington State Diabetes Prediction Study. *Diabetes Care.* Mar 2002;25(3):505-511.
- **19.** Ziegler AG, Hummel M, Schenker M, Bonifacio E. Autoantibody appearance and risk for development of childhood diabetes in offspring of parents with type 1 diabetes: the 2-year analysis of the German BABYDIAB Study. *Diabetes*. Mar 1999;48(3):460-468.

- **20.** Baekkeskov S, Nielsen JH, Marner B, Bilde T, Ludvigsson J, Lernmark A. Autoantibodies in newly diagnosed diabetic children immunoprecipitate human pancreatic islet cell proteins. *Nature*. Jul 8 1982;298(5870):167-169.
- **21.** Strebelow M, Schlosser M, Ziegler B, Rjasanowski I, Ziegler M. Karlsburg Type I diabetes risk study of a general population: frequencies and interactions of the four major Type I diabetes-associated autoantibodies studied in 9419 schoolchildren. *Diabetologia.* Jun 1999;42(6):661-670.
- 22. Ludvigsson J, Faresjo M, Hjorth M, et al. GAD treatment and insulin secretion in recent-onset type 1 diabetes. *N Engl J Med.* Oct 30 2008;359(18):1909-1920.
- **23.** Gullstrand C, Wahlberg J, Ilonen J, Vaarala O, Ludvigsson J. Progression to type 1 diabetes and autoantibody positivity in relation to HLA-risk genotypes in children participating in the ABIS study. *Pediatr Diabetes*. Jun 2008;9(3 Pt 1):182-190.
- 24. Borg H, Marcus C, Sjoblad S, Fernlund P, Sundkvist G. Insulin autoantibodies are of less value compared with islet antibodies in the clinical diagnosis of autoimmune type 1 diabetes in children older than 3 yr of age. *Pediatr Diabetes*. Sep 2002;3(3):149-154.
- **25.** Bonifacio E, Genovese S, Braghi S, et al. Islet autoantibody markers in IDDM: risk assessment strategies yielding high sensitivity. *Diabetologia*. Jul 1995;38(7):816-822.
- 26. Seissler J, Scherbaum WA. Autoimmune diagnostics in diabetes mellitus. *Clin Chem Lab Med.* 2006;44(2):133-137.
- 27. Holmberg H, Vaarala O, Sadauskaite-Kuehne V, Ilonen J, Padaiga Z, Ludvigsson J. Higher prevalence of autoantibodies to insulin and GAD65 in Swedish compared to Lithuanian children with type 1 diabetes. *Diabetes Res Clin Pract.* Jun 2006;72(3):308-314.
- **28.** Williams AJ, Norcross AJ, Dix RJ, Gillespie KM, Gale EA, Bingley PJ. The prevalence of insulin autoantibodies at the onset of Type 1 diabetes is higher in males than females during adolescence. *Diabetologia*. Oct 2003;46(10):1354-1356.
- **29.** Marciulionyte D, Williams AJ, Bingley PJ, Urbonaite B, Gale EA. A comparison of the prevalence of islet autoantibodies in children from two countries with differing incidence of diabetes. *Diabetologia*. Jan 2001;44(1):16-21.
- **30.** Johansson C, Samuelsson U, Ludvigsson J. A high weight gain early in life is associated with an increased risk of type 1 (insulin-dependent) diabetes mellitus. *Diabetologia.* Jan 1994;37(1):91-94.
- **31.** Roche EF, Menon A, Gill D, Hoey H. Clinical presentation of type 1 diabetes. *Pediatr Diabetes*. Jun 2005;6(2):75-78.
- **32.** Defronzo RA. DH, Keen H., Zimmet P. International textbook of diabetes mellitus, Third edition. *Chichester, West sussex. England.* 2004.
- **33.** Montgomery EL, Batch JA. Cataracts in insulin-dependent diabetes mellitus: sixteen years' experience in children and adolescents. *J Paediatr Child Health*. Apr 1998;34(2):179-182.
- **34.** Rewers A, Klingensmith G, Davis C, et al. Presence of diabetic ketoacidosis at diagnosis of diabetes mellitus in youth: the Search for Diabetes in Youth Study. *Pediatrics*. May 2008;121(5):e1258-1266.
- **35.** EURODIAB ACE Study Group. Variation and trends in incidence of childhood diabetes in Europe. *Lancet.* Mar 11 2000;355(9207):873-876.
- **36.** Karvonen M, Viik-Kajander M, Moltchanova E, Libman I, LaPorte R, Tuomilehto J. Incidence of childhood type 1 diabetes worldwide. Diabetes Mondiale (DiaMond) Project Group. *Diabetes Care*. Oct 2000;23(10):1516-1526.

- **37.** Dahlquist G, Mustonen L. Analysis of 20 years of prospective registration of childhood onset diabetes time trends and birth cohort effects. Swedish Childhood Diabetes Study Group. *Acta Paediatr*. Oct 2000;89(10):1231-1237.
- **38.** Pundziute-Lycka A, Dahlquist G, Urbonaite B, Zalinkevicius R. Time trend of childhood type 1 diabetes incidence in Lithuania and Sweden, 1983-2000. *Acta Paediatr.* Nov 2004;93(11):1519-1524.
- **39.** Pundziute-Lycka A, Dahlquist G, Nystrom L, et al. The incidence of Type I diabetes has not increased but shifted to a younger age at diagnosis in the 0-34 years group in Sweden 1983-1998. *Diabetologia*. Jun 2002;45(6):783-791.
- **40.** Wennick A, Hallstrom I. Swedish families' lived experience when a child is first diagnosed as having insulin-dependent diabetes mellitus: An ongoing learning process. *J Fam Nurs.* Nov 2006;12(4):368-389.
- **41.** Wennick A, Hallstrom I. Families' lived experience one year after a child was diagnosed with type 1 diabetes. *J Adv Nurs*. Nov 2007;60(3):299-307.
- **42.** Huus K, Enskar K. Adolescents' experience of living with diabetes. *Paediatr Nurs.* Apr 2007;19(3):29-31.
- **43.** Miller S. Hearing from children who have diabetes. *J Child Health Care*. Spring 1999;3(1):5-12.
- 44. Ory MG, Kronenfeld JJ. Living with juvenile diabetes mellitus. *Pediatr Nurs*. Sep-Oct 1980;6(5):47-50.
- **45.** Hyoty H, Taylor KW. The role of viruses in human diabetes. *Diabetologia*. Oct 2002;45(10):1353-1361.
- **46.** Tauriainen S, Salminen K, Hyoty H. Can enteroviruses cause type 1 diabetes? *Ann N Y Acad Sci.* Nov 2003;1005:13-22.
- **47.** Gamble DR, Taylor KW. Seasonal incidence of diabetes mellitus. *Br Med J.* Sep 13 1969;3(5671):631-633.
- **48.** Viskari HR, Koskela P, Lonnrot M, et al. Can enterovirus infections explain the increasing incidence of type 1 diabetes? *Diabetes Care.* Mar 2000;23(3):414-416.
- **49.** Viskari H, Ludvigsson J, Uibo R, et al. Relationship between the incidence of type 1 diabetes and enterovirus infections in different European populations: results from the EPIVIR project. *J Med Virol*. Apr 2004;72(4):610-617.
- 50. Hyoty H. Enterovirus infections and type 1 diabetes. Ann Med. 2002;34(3):138-147.
- **51.** Kolb H, Elliott RB. Increasing incidence of IDDM a consequence of improved hygiene? *Diabetologia*. Jul 1994;37(7):729.
- **52.** Bach JF. Protective role of infections and vaccinations on autoimmune diseases. *J Autoimmun.* May 2001;16(3):347-353.
- **53.** Wahlberg J, Vaarala O, Ludvigsson J. Dietary risk factors for the emergence of type 1 diabetes-related autoantibodies in 21/2 year-old Swedish children. *Br J Nutr.* Mar 2006;95(3):603-608.
- **54.** Bognetti E, Meschi F, Malavasi C, et al. HLA-antigens in Italian type 1 diabetic patients: role of DR3/DR4 antigens and breast feeding in the onset of the disease. *Acta Diabetol.* 1992;28(3-4):229-232.
- 55. Saukkonen T, Virtanen SM, Karppinen M, et al. Significance of cow's milk protein antibodies as risk factor for childhood IDDM: interactions with dietary cow's milk intake and HLA-DQB1 genotype. Childhood Diabetes in Finland Study Group. *Diabetologia.* Jan 1998;41(1):72-78.
- **56.** Rosenbauer J, Herzig P, Giani G. Early infant feeding and risk of type 1 diabetes mellitus-a nationwide population-based case-control study in pre-school children. *Diabetes Metab Res Rev.* Mar-Apr 2008;24(3):211-222.

- 57. Virtanen SM, Knip M. Nutritional risk predictors of beta cell autoimmunity and type 1 diabetes at a young age. *Am J Clin Nutr.* Dec 2003;78(6):1053-1067.
- **58.** Lamb MM, Myers MA, Barriga K, Zimmet PZ, Rewers M, Norris JM. Maternal diet during pregnancy and islet autoimmunity in offspring. *Pediatr Diabetes*. Apr 2008;9(2):135-141.
- **59.** Myers M, Zimmet P. Halting the accelerating epidemic of type 1 diabetes. *Lancet*. May 24 2008;371(9626):1730-1731.
- **60.** Stene LC, Ulriksen J, Magnus P, Joner G. Use of cod liver oil during pregnancy associated with lower risk of Type I diabetes in the offspring. *Diabetologia*. Sep 2000;43(9):1093-1098.
- **61.** Fronczak CM, Baron AE, Chase HP, et al. In utero dietary exposures and risk of islet autoimmunity in children. *Diabetes Care*. Dec 2003;26(12):3237-3242.
- **62.** Hagglof B, Blom L, Dahlquist G, Lonnberg G, Sahlin B. The Swedish childhood diabetes study: indications of severe psychological stress as a risk factor for type 1 (insulin-dependent) diabetes mellitus in childhood. *Diabetologia*. Aug 1991;34(8):579-583.
- **63.** Sepa A, Ludvigsson J. Psychological stress and the risk of diabetes-related autoimmunity: a review article. *Neuroimmunomodulation*. 2006;13(5-6):301-308.
- **64.** Sepa A, Frodi A, Ludvigsson J. Mothers' experiences of serious life events increase the risk of diabetes-related autoimmunity in their children. *Diabetes Care.* Oct 2005;28(10):2394-2399.
- **65.** Wilkin TJ. Diabetes: 1 and 2, or one and the same? Progress with the accelerator hypothesis. *Pediatr Diabetes*. Jun 2008;9(3 Pt 2):23-32.
- 66. Kissebah AH, Krakower GR. Regional adiposity and morbidity. *Physiol Rev.* Oct 1994;74(4):761-811.
- **67.** Marild S, Bondestam M, Bergstrom R, Ehnberg S, Hollsing A, Albertsson-Wikland K. Prevalence trends of obesity and overweight among 10-year-old children in western Sweden and relationship with parental body mass index. *Acta Paediatr.* Dec 2004;93(12):1588-1595.
- **68.** Kohn M, Rees JM, Brill S, et al. Preventing and treating adolescent obesity: a position paper of the Society for Adolescent Medicine. *J Adolesc Health.* Jun 2006;38(6):784-787.
- **69.** Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *Bmj.* May 6 2000;320(7244):1240-1243.
- **70.** Wang Y. Epidemiology of childhood obesity--methodological aspects and guidelines: what is new? *Int J Obes Relat Metab Disord*. Nov 2004;28 Suppl 3:S21-28.
- **71.** Lee S, Bacha F, Gungor N, Arslanian SA. Waist circumference is an independent predictor of insulin resistance in black and white youths. *J Pediatr*. Feb 2006;148(2):188-194.
- 72. Wadden TA, Foster GD. Behavioral treatment of obesity. *Med Clin North Am*. Mar 2000;84(2):441-461, vii.
- 73. Comuzzie AG, Allison DB. The search for human obesity genes. *Science*. May 29 1998;280(5368):1374-1377.
- 74. Erlanson-Albertsson C, Zetterstrom R. The global obesity epidemic: snacking and obesity may start with free meals during infant feeding. *Acta Paediatr.* Nov 2005;94(11):1523-1531.
- 75. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. Aug 10 2002;360(9331):473-482.

- **76.** Reilly JJ, Methven E, McDowell ZC, et al. Health consequences of obesity. *Arch Dis Child*. Sep 2003;88(9):748-752.
- 77. Whitaker RC, Wright JA, Pepe MS, Seidel KD, Dietz WH. Predicting obesity in young adulthood from childhood and parental obesity. *N Engl J Med.* Sep 25 1997;337(13):869-873.
- **78.** Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*. Apr 5 2006;295(13):1549-1555.
- **79.** Lobstein T, Frelut ML. Prevalence of overweight among children in Europe. *Obes Rev.* Nov 2003;4(4):195-200.
- **80.** Neovius M, Janson A, Rossner S. Prevalence of obesity in Sweden. *Obes Rev.* Feb 2006;7(1):1-3.
- **81.** Sjoberg A, Lissner L, Albertsson-Wikland K, Marild S. Recent anthropometric trends among Swedish school children: evidence for decreasing prevalence of overweight in girls. *Acta Paediatr.* Jan 2008;97(1):118-123.
- **82.** Guo SS, Wu W, Chumlea WC, Roche AF. Predicting overweight and obesity in adulthood from body mass index values in childhood and adolescence. *Am J Clin Nutr.* Sep 2002;76(3):653-658.
- **83.** de Beer M, Hofsteenge GH, Koot HM, Hirasing RA, Delemarre-van de Waal HA, Gemke RJ. Health-related-quality-of-life in obese adolescents is decreased and inversely related to BMI. *Acta Paediatr.* May 2007;96(5):710-714.
- **84.** Pinhas-Hamiel O, Singer S, Pilpel N, Fradkin A, Modan D, Reichman B. Healthrelated quality of life among children and adolescents: associations with obesity. *Int J Obes (Lond)*. Feb 2006;30(2):267-272.
- **85.** Araujo CL, Victora CG, Hallal PC, Gigante DP. Breastfeeding and overweight in childhood: evidence from the Pelotas 1993 birth cohort study. *Int J Obes (Lond)*. Mar 2006;30(3):500-506.
- **86.** Oddy WH, Scott JA, Graham KI, Binns CW. Breastfeeding influences on growth and health at one year of age. *Breastfeed Rev.* Mar 2006;14(1):15-23.
- **87.** Toschke AM, Martin RM, von Kries R, Wells J, Smith GD, Ness AR. Infant feeding method and obesity: body mass index and dual-energy X-ray absorptiometry measurements at 9-10 y of age from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Am J Clin Nutr.* Jun 2007;85(6):1578-1585.
- **88.** Weyermann M, Rothenbacher D, Brenner H. Duration of breastfeeding and risk of overweight in childhood: a prospective birth cohort study from Germany. *Int J Obes* (*Lond*). Aug 2006;30(8):1281-1287.
- **89.** Burke V, Beilin LJ, Simmer K, et al. Breastfeeding and overweight: longitudinal analysis in an Australian birth cohort. *J Pediatr*. Jul 2005;147(1):56-61.
- **90.** Burdette HL, Whitaker RC, Hall WC, Daniels SR. Breastfeeding, introduction of complementary foods, and adiposity at 5 y of age. *Am J Clin Nutr.* Mar 2006;83(3):550-558.
- **91.** Rolland-Cachera MF, Deheeger M, Akrout M, Bellisle F. Influence of macronutrients on adiposity development: a follow up study of nutrition and growth from 10 months to 8 years of age. *Int J Obes Relat Metab Disord*. Aug 1995;19(8):573-578.
- **92.** Whitehead RG. For how long is exclusive breast-feeding adequate to satisfy the dietary energy needs of the average young baby? *Pediatr Res.* Feb 1995;37(2):239-243.
- **93.** Lucas A, Sarson DL, Blackburn AM, Adrian TE, Aynsley-Green A, Bloom SR. Breast vs bottle: endocrine responses are different with formula feeding. *Lancet*. Jun 14 1980;1(8181):1267-1269.

- 94. Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. *Pediatrics*. Mar 1998;101(3 Pt 2):539-549.
- **95.** Flodmark CE, Lissau I, Moreno LA, Pietrobelli A, Widhalm K. New insights into the field of children and adolescents' obesity: the European perspective. *Int J Obes Relat Metab Disord*. Oct 2004;28(10):1189-1196.
- **96.** Burke V, Beilin LJ, Durkin K, Stritzke WG, Houghton S, Cameron CA. Television, computer use, physical activity, diet and fatness in Australian adolescents. *Int J Pediatr Obes.* 2006;1(4):248-255.
- **97.** Epstein LH, Roemmich JN, Robinson JL, et al. A randomized trial of the effects of reducing television viewing and computer use on body mass index in young children. *Arch Pediatr Adolesc Med.* Mar 2008;162(3):239-245.
- **98.** Gazzaniga JM, Burns TL. Relationship between diet composition and body fatness, with adjustment for resting energy expenditure and physical activity, in preadolescent children. *Am J Clin Nutr.* Jul 1993;58(1):21-28.
- **99.** Nguyen VT, Larson DE, Johnson RK, Goran MI. Fat intake and adiposity in children of lean and obese parents. *Am J Clin Nutr*. Apr 1996;63(4):507-513.
- **100.** Tucker LA, Seljaas GT, Hager RL. Body fat percentage of children varies according to their diet composition. *J Am Diet Assoc*. Sep 1997;97(9):981-986.
- **101.** Ricketts CD. Fat preferences, dietary fat intake and body composition in children. *Eur J Clin Nutr.* Nov 1997;51(11):778-781.
- **102.** Robertson SM, Cullen KW, Baranowski J, Baranowski T, Hu S, de Moor C. Factors related to adiposity among children aged 3 to 7 years. *J Am Diet Assoc.* Aug 1999;99(8):938-943.
- **103.** Atkin LM, Davies PS. Diet composition and body composition in preschool children. *Am J Clin Nutr.* Jul 2000;72(1):15-21.
- **104.** Davison KK, Birch LL. Child and parent characteristics as predictors of change in girls' body mass index. *Int J Obes Relat Metab Disord*. Dec 2001;25(12):1834-1842.
- 105. Skinner JD, Bounds W, Carruth BR, Morris M, Ziegler P. Predictors of children's body mass index: a longitudinal study of diet and growth in children aged 2-8 y. *Int J Obes Relat Metab Disord.* Apr 2004;28(4):476-482.
- **106.** Scaglioni S, Agostoni C, Notaris RD, et al. Early macronutrient intake and overweight at five years of age. *Int J Obes Relat Metab Disord*. Jun 2000;24(6):777-781.
- 107. Davies PS. Diet composition and body mass index in pre-school children. *Eur J Clin Nutr.* Jul 1997;51(7):443-448.
- **108.** Jago R, Baranowski T, Baranowski JC, Thompson D, Greaves KA. BMI from 3-6 y of age is predicted by TV viewing and physical activity, not diet. *Int J Obes (Lond)*. Jun 2005;29(6):557-564.
- **109.** Garemo M, Lenner RA, Strandvik B. Swedish pre-school children eat too much junk food and sucrose. *Acta Paediatr*. Feb 2007;96(2):266-272.
- **110.** Dehghan M, Akhtar-Danesh N, Merchant AT. Childhood obesity, prevalence and prevention. *Nutr J.* 2005;4:24.
- **111.** Tremblay MS, Willms JD. Is the Canadian childhood obesity epidemic related to physical inactivity? *Int J Obes Relat Metab Disord*. Sep 2003;27(9):1100-1105.
- 112. Swinburn B, Egger G. Preventive strategies against weight gain and obesity. *Obes Rev.* Nov 2002;3(4):289-301.
- **113.** Clemes SA, Matchett N, Wane SL. Reactivity: an issue for short-term pedometer studies? *Br J Sports Med.* Jan 2008;42(1):68-70.
- **114.** Vincent SDP, R. P. . Does Reactivity Exist in Children When Measuring Activity Levels with Pedometers? *Pediatric Exercise Science*. 2002;14:56-63.
- 115. SPSS. Version 11.5 for windows. SPSS inc. Chicago IL;USA. 2003.

- **116.** Kirkwood. B R. SAC. Medical Statistics. *Blackwell Publishing* 2007.
- 117. G AD. Practical statistics for medical research. London UK: Chapman & Hall; 1994.
- **118.** Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care*. Jun 2004;27(6):1487-1495.
- **119.** Reinard, ed. *Communication Research Statistics*. Thousland Oaks, Califonia; 2006. SAGE Publications.
- **120.** Beauchamp TL. CJ, ed. *Principles of Biomedical Ethics* Oxford University Press; 1989; No. Third Edition.
- **121.** Thompson . MK, Boyd KM., Horsburgh DH., ed. *Nursing Ethics*. Edinburgh, UK: Churchill Livingstone; 2006; No. Fifth Edition.
- **122.** Stolt UG, Liss PE, Svensson T, Ludvigsson J. Attitudes to bioethical issues: a case study of a screening project. *Soc Sci Med.* May 2002;54(9):1333-1344.
- **123.** Gustafsson Stolt U, Ludvigsson J, Liss PE, Svensson T. Bioethical theory and practice in genetic screening for type 1 diabetes. *Med Health Care Philos.* 2003;6(1):45-50.
- **124.** Huus K, Ludvigsson JF, Enskar K, Ludvigsson J. Risk factors in childhood obesityfindings from the All Babies In Southeast Sweden (ABIS) cohort. *Acta Paediatr.* Sep 2007;96(9):1321-1325.
- 125. Kristman V, Manno M, Cote P. Loss to follow-up in cohort studies: how much is too much? *Eur J Epidemiol.* 2004;19(8):751-760.
- **126.** Altman DG. Statistics in medical journals: some recent trends. *Stat Med.* Dec 15 2000;19(23):3275-3289.
- **127.** Fewtrell MS, Kennedy K, Singhal A, et al. How much loss to follow-up is acceptable in long-term randomised trials and prospective studies? *Arch Dis Child.* Jun 2008;93(6):458-461.
- **128.** Heitmann BL, Lissner L. Dietary underreporting by obese individuals--is it specific or non-specific? *Bmj.* Oct 14 1995;311(7011):986-989.
- **129.** Jackson J, Strauss CC, Lee AA, Hunter K. Parents' accuracy in estimating child weight status. *Addict Behav.* 1990;15(1):65-68.
- **130.** Sepa A, Wahlberg J, Vaarala O, Frodi A, Ludvigsson J. Psychological stress may induce diabetes-related autoimmunity in infancy. *Diabetes Care*. Feb 2005;28(2):290-295.
- **131.** Dubois L, Girard M. Early determinants of overweight at 4.5 years in a populationbased longitudinal study. *Int J Obes (Lond)*. Apr 2006;30(4):610-617.
- **132.** Bergmann KE, Bergmann RL, Von Kries R, et al. Early determinants of childhood overweight and adiposity in a birth cohort study: role of breast-feeding. *Int J Obes Relat Metab Disord*. Feb 2003;27(2):162-172.
- **133.** Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugarsweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet.* Feb 17 2001;357(9255):505-508.
- **134.** Angbratt M. EE, Funcke S., Nilsson U., Säterskog C., Söderlind M. Mapping of childrens weight and overweight development in Östergötland. *Centre of Public Health in Linköping.* 2003.
- **135.** Boon CS, Clydesdale FM. A review of childhood and adolescent obesity interventions. *Crit Rev Food Sci Nutr.* 2005;45(7-8):511-525.
- **136.** Plourde G. Preventing and managing pediatric obesity. Recommendations for family physicians. *Can Fam Physician*. Mar 2006;52:322-328.