Linköping University Postprint

Inter-species comparison of seed dormancy and germination of six annual *Asteraceae* weeds in an ecological context

Laila M. Karlsson, T. Tamado and Per Milberg

N.B.: When citing this work, cite the original article.

Original publication:

Laila M. Karlsson, T. Tamado and Per Milberg, Inter-species comparison of seed dormancy and germination of six annual *Asteraceae* weeds in an ecological context, 2008, Seed Science Research, (18), 35-45.

<http://dx.doi.org/10.1017/S0960258508888496>.

Copyright: Cambridge University Press, <http://journals.cambridge.org/>

Postprint available free at:

Linköping University E-Press: <http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-11879>

Version 19 September 2007

**Inter-species comparison of seed dormancy and germination of six annual
Asteraceae weeds in an ecological context**

Laila M. Karlsson¹, T. Tamado² and Per Milberg^{1*}

¹IFM Division of Ecology, Linköping University, SE-581 83 Linköping, Sweden

²Department of Plant Science, Haramaya University, Dire Dawa, and Woalita Sodo
University, P.O. Box 138, Ethiopia

*Correspondence:

Fax: +46 13 28 13 99

Email: permi@ifm.liu.se

Running head: Dormancy patterns of annual *Asteraceae*

Received: 5 February 2007

Accepted after revision:

Abstract

To understand germination timing in an ecological context, the response to environmental events that affect seed dormancy is central, and has to be combined with knowledge of germination responses to different circumstances. In this study, seed dormancy, germination, and emergence phenology of six annual co-occurring weedy *Asteraceae* species were investigated. Three pre-treatments (warm and cold stratification, and dry storage) were tested as possible dormancy affecting environmental events. Seeds were also sown outdoors. Species-specific differences were revealed in analyses. To facilitate general descriptions of dormancy patterns and germination preferences separately, condensed responses to the different possible dormancy affecting treatments and relative germination in different environments were plotted, giving species-specific patterns. Most species exhibited decreased dormancy to two or three pre-treatments. Dormancy was most effectively reduced by cold stratification for three species (*Guizotia scabra*, *Parthenium hysterophorus*, *Verbesina encelioides*), by warm stratification for two (*Bidens pilosa*, *Galinsoga parviflora*) and by dry storage for one (*Tagetes minuta*). All species germinated more when provided with light than in continuous darkness. Temperature levels most suitable for germination varied from low (15/5-20/10°C) for *Verbesina encelioides* to high (25/15-30/20°C) for *Bidens pilosa*. It is concluded that, even though the species have different dormancy patterns and germination preferences that suggest different possible distribution ranges, the species achieve similar emergence timing in the field in environments with a pronounced dry period after dispersal and small annual temperature fluctuations.

Keywords: *Bidens pilosa*, *Galinsoga parviflora*, *Guizotia scabra* ssp. *schimperi*, *Parthenium hysterophorus*, *Tagetes minuta*, tropical weeds, *Verbesina encelioides*

Introduction

To understand emergence timing in the field, as summarized and diagrammed by Merritt *et al.* (2007), it is necessary to include local environmental events affecting the degree of seed dormancy, as well as circumstances actually leading to germination. Unfortunately, "seed dormancy" is used with various definitions in the literature (e.g. Harper, 1957, 1977; Lang, 1987; Baskin and Baskin, 2004): referring to the circumstances that directly promote/prevent germination and, or solely, to the circumstances (including embryo and seed morphology) that make seeds delay or prevent germination even though subjected to suitable germination environments. The definition used in this paper, aiming at studying dormancy and germination from an ecological perspective, is that seed dormancy is a seed characteristic that prevents germination, even if suitable germination conditions prevail. Neither embryo or seed morphology, nor germination mechanisms, are involved in this definition.

To describe dormancy and germination patterns of species, we use (1) "dormancy pattern", i.e. environmental events that reduce and, if applicable, induce dormancy; (2) "germination preferences", i.e. environments that are (or become during dormancy reduction) suitable for germination; and (3) "dormancy strength" (Karlsson and Milberg, 2007a, b, in press). To be ecologically meaningful, seed dormancy should be regarded as a continuous property of a seed batch (even though it is not definitely known whether or not it is a continuum or an on-off property for an individual seed).

Germination preference alludes to a dynamic characteristic, not to a requirement for germination or to a theoretical optimum, e.g. light is a germination preference for several species, even though such species frequently do not have light as a requirement for germination in all circumstances (*e.g.* Karlsson and Milberg, 2007c). Dormancy strength, referring to the general pattern, is described as strong-weak, and the extent of dormancy present at any specific moment is referred to as "degree of dormancy".

The model by Merritt *et al.* (2007) addressed emergence timing of Australian species in Australia. For species occurring worldwide, one can suspect that different environmental events are important for emergence timing in different parts of the world. To evaluate the importance of different environmental circumstances, to foresee possible distributional ranges, and to compare species according to their response to specific situations and in general, we regard it as desirable to describe dormancy and germination separately through parameters that can be measured on continuous scales. The dormancy sorting systems suggested in the literature are based on discrete sorting; Harper (1957, 1977) used a system where seeds could be placed in different groups depending on the circumstances instantaneously preventing germination; Lang (1987) regarded the within-organism location of physiological factors preventing germination or the presence of an unsuitable environment as the most important sorting factors, and Baskin and Baskin (2004) first sorted according to morphology and then into subgroups that could include species with different responses to warm and cold stratification. Vleeshouwers *et al.* (1995) underlined the sorting and definition problem when they attempted to integrate physiology and ecology with respect to seed dormancy, and decided simply to exclude, before beginning their reasoning, all species that have small embryos (in relation to the endosperm), and those that do not imbibe water when fresh.

The extent and rate of response to pre-treatments are important parameters for evaluating the effect of different, possible dormancy-affecting, environmental events. These two factors can be measured by fitting germination results from tests over time to a logistic function (Karlsson and Milberg, 2007c). By condensing such measurements and plotting them together with germination preferences, species-specific patterns might arise, aiding in evaluation of emergence patterns and in comparisons of species. To evaluate this, we used six annual weedy *Asteraceae* that co-occur in eastern Ethiopia: *Bidens pilosa* L., *Galinsoga parviflora* Cav., *Guizotia scabra* (Vis.) Chiov. ssp. *schimperi* (Sch.Bip.) Baagoe (hereafter referred to as *Guizotia scabra* and included in the term "species"), *Parthenium hysterophorus* L., *Tagetes minuta* L. and *Verbesina encelioides* (Cav.) A.Gray.

Materials and methods

Seed collection

Ripe seeds of *Bidens pilosa*, *Galinsoga parviflora*, *Guizotia scabra*, *Parthenium hysterophorus*, *Tagetes minuta*, and *Verbesina encelioides* were collected on 7 and 8 November 2005. All species were collected at two sites: around Haramaya town (N 09°23' E 42°01') and at cultivated areas on the Haramaya University campus (N 09°25' E 42°02'; previously named Alemaya University). The sites are *ca.* 5 km apart, and are located in eastern Ethiopia, just above 2000 m a.s.l. All sites used were agricultural fields, including the margins. At collection, flower heads were lightly shaken in a bucket, and achenes (hereafter called seeds) that fell off were used. Seeds were used in the experiments as they are dispersed in nature, i.e. we did not remove the pappus, wings, enclosing petals or other structures. Experiments in the laboratory and in the

field commenced about 10 d after collection, and until then, seeds were stored dry at natural fluctuating temperatures.

Germination experiments

Seeds were tested for germination at the beginning of experiments ("fresh") and after three different pre-treatments: cold or warm stratification, or dry storage, for 2.5, 5, 10, 20 or 30 weeks. Germination tests were performed both in light during daytime ("light") and in continuous darkness ("darkness"). Germination tests were terminated after 4 weeks; germination in dishes in light was checked after 2 and 4 weeks, and dishes in darkness only at termination. When terminating a dish, ungerminated seeds were counted. Seeds that were soft and/or overgrown with mould were considered dead.

All pre-treatments were performed in continuous darkness. Average daily min-max temperatures were 4.5-5.4 (SD 0.17-0.15) °C in the cold environment, and 22.0-23.2 (SD 0.5-0.7) °C in the environment used for warm stratification and dry storage. The dry stored seeds were subjected to a relative humidity of 28.8-31.5 (average daily min-max, SD 3.3-4.1) %, which was achieved by putting in open bowls with moist sand and KCl(s), when considered appropriate, to adjust for seasonal variation in indoor . Germination tests were performed in incubators (Rubarth Apparatebau, Laatzen, Germany), with alternating 10/10 h (day/night) temperatures of 15/5, 20/10, 25/15 or 30/20°C with a 2 h, linear transition in between. The incubators were illuminated for 12 h at a photon flux density (400-700 nm) of 28-62 $\mu\text{mol m}^{-2} \text{s}^{-1}$ [measured with a SKP 200 photometer and SKP 215 sensor (Skye Instruments Ltd, Wales); measurement precision 1 $\mu\text{mol m}^{-2} \text{s}^{-1}$], and a 660/730 nm photon fluence ratio of 3.0-3.5 [(SKR 100

photometer with a SKR 110 sensor (Skye Instruments Ltd)]. Dishes in light were randomly re-arranged in the incubators at least once a week.

For germination tests, *ca.* 100 (average 116 ± 31) seeds were distributed in 3 to 4 Petri dishes (50 mm diameter), depending on seed size. The substrate was 10 ml of quartz sand (0.35 mm grain size, Baskarpsand 35, AB Baskarpsand, Habo, Sweden), moistened with 3.7 ml deionized water, plus enough water for imbibition; all dishes were sealed with Parafilm™. For dark treatments, dishes were wrapped in aluminium foil, and the dishes for warm stratification treatments were also put in plastic bags.

For all four incubation temperatures, the dishes with fresh seeds tested in light were subjected to continuous incubation for an additional 30 weeks after terminating the initial test, and also a set of dishes in darkness was incubated for 34 weeks. In addition, seeds were subjected to a 40 week treatment, comprising an initial dry period of 12 weeks, followed by 8 weeks of moist incubation in light at the four temperatures, 4 weeks of dryness and an additional 16 weeks of moist incubation at the same temperature as before, resembling the dry and rainy periods in eastern Ethiopia after harvest. The dishes in light that were kept 34 and 40 weeks were checked at least every 6 weeks.

Seedling emergence phenology

Soil was sampled at the Haramaya University campus on 17 October 2005; the soil on the site was well drained loam. Soil was sterilized by 100°C water vapour for at least 8 h. Pots (9 cm diameter), with a non-woven glass fibre sheet in the bottom, were filled with *ca.* 250 ml soil, and 75-100 seeds, depending on seed size, were placed on the soil surface in each pot. Two pots, individually covered with mosquito netting, were used

for each seed batch, and were buried in soil, at a cultivated area on the Haramaya University campus at a depth such that the soil surface in pots was at the same level as the surrounding soil surface. Seeds were then subjected to local temperatures and precipitation. Pots were checked for seedlings at least every week, and emerged seedlings were counted and removed. Temperature and precipitation were measured at the Haramaya University weather station.

Calculations and analyses

For the seedling emergence phenology experiment, emergence was calculated as the fraction of seeds sown in each seed batch. In germination tests, the fraction germinated was calculated from the number of seeds used, excluding seeds considered dead.

For all analyses, Statistica (StatSoft, Inc., 2004) was used. Analyses of variance were done on (1) arcsine-transformed germination data and (2) the difference between each adjacent test occasion as weekly averages. The two seed batches of each species were considered replicates of the species in both analyses. Categorical predictors were species, pre-treatment, germination test temperature and light condition. Time in pre-treatment was treated as a continuous variable in the analysis of germination. In the analysis of differences, each time span ("moment") between adjacent test occasions was treated as a categorical predictor. For interpretation, special attention was paid to the factor "species" and interactions including "species", to evaluate possible differences between species.

To investigate the general, relative, difference between seed batches and species, a principal component analysis (PCA) was done on arcsine-transformed germination

results, together with absolute differences between each adjacent test occasion. In the PCA, results for seed batches were not individually standardized.

The average extent of, and time to, final response for each pre-treatment were calculated. A logistic function (Equation 1) was fitted to each of the 288 series of six points in time from germination tests, i.e. one per seed batch, pre-treatment and germination test environment. In Equation 1, $f(t)$ is germination, t is time in pre-treatment, a , b , c and d are constants, where $0 \leq d \leq 1$ and $0 \leq (c+d) \leq 1$, and e (approximated to 2.71828) is the base of the natural logarithm.

$$f(t) = \frac{e^{a+bt} \cdot c}{1 + e^{a+bt}} + d \quad (1)$$

Calculations were performed using non-linear estimation and user-specified regression. The loss function, with quadratic difference between observed and predicted values, and the weight function (on the number of seeds tested at each occasion), with the pre-selected quasi-Newton estimation, were used. Initial values for a , b and c were set to zero and d to the actual germination of fresh seeds. Time for achieving major response to pre-treatment was calculated as the time between $t = 0$ and the right extreme point (hereafter referred to as $f'''_{\max}$) of the third derivative of Equation 1 (Equation 2, Figure 1).

$$f'''(t) = \frac{b^3 c e^{a+bt} (1 + (e^{a+bt})^6 - 9(e^{a+bt})^4 - 16(e^{a+bt})^3 - 9(e^{a+bt})^2)}{(1 + e^{a+bt})^8} \quad (2)$$

The average calculated final germination [$f(30)$ from Equation 1, $\pm$ SD] and weighted average response time [Equation 3, $\pm$ weighted SD (Equation 4)] for the

$$\bar{t}_w = \frac{\sum tr}{\sum r} \quad (3)$$

$$SD_w = \sqrt{\frac{\sum (r(t - \bar{t}_w))^2}{\sum r - \frac{\sum r^2}{N}}} \quad (4)$$

Figure 1. An example of the logistic function $f(t) = e^{a+bt} \cdot c / (1 + e^{a+bt}) + d$ fitted to germination data. $f(t)$ is germination, t is time in pre-treatment, a , b , c , d are constants, and e is the base of the natural logarithm. To obtain response time (i.e. the time from $t = 0$ to the point when major germination response is considered achieved), the corresponding t for the shown maximum of $f'''(t)$ can be calculated.

different pre-treatments were calculated. For each batch and test environment, t is the time obtained from Equation 2, and r is the extent of response (i.e. the absolute difference between $f(0)$ and $f(30)$ from Equation 1). Because of the weighting, each calculated time to achieve a major response contributes to the weighted average time, and corresponding SD, in direct proportion to magnitude of the response. Functions without any change at all (29 and 14 functions continuously at 0% and 100%, respectively) or with $f'''_{\max}$ outside $0 \leq t \leq 30$ (36 functions) were excluded from calculation of weighted average time to major response.

Together with average germination for fresh seeds in all environments, average calculated final germination was plotted against average response time (Equation 3). Further, average overall germination at all test occasions per species and test environment was calculated, and the relative average germination in each environment was plotted, for each environment, as the fraction of germination in the environment where most germination occurred for each species.

Results

Responses to test environments and pre-treatments

Fresh seeds germinated less than 50% in all test environments, with the exception of *Tagetes minuta*, which germinated 100% in light at higher temperature regimes (Fig. 2). All three pre-treatments tested led to increased germination for all species, except dry storage for *Parthenium hysterophorus* and warm incubation for *Verbesina encelioides*, for which there was no observable effect (Fig. 2).

Species was a significant explanatory factor for both germination and for differences between test occasions (Tables 1, 2). Overall, *Parthenium hysterophorus* germinated the least and *Tagetes minuta* the most (Fig. 2). *Bidens pilosa* and *Guizotia scabra* had the largest, and *Parthenium hysterophorus* and *Tagetes minuta* the smallest, overall differences between test occasions (Fig. 2).

Second order interactions, including the factor "species", were significant in all cases for germination (Table 1), and in all cases except "species*temperature" for difference between occasions (Table 2). For germination, this reflects species-specific differences in the extent of germination following different pre-treatments, different test temperatures and light requirements for germination (Fig. 2). For differences, the

Table 1. ANOVA on germination (arcsine transformed) of six *Asteraceae* species, each collected from two sites. Seeds were subjected to germination tests in eight different environments when fresh and after three different pre-treatments. Time was treated as a continuous variable. Each seed batch was considered one replicate in the analysis.

Factor	df	F	p
Intercept	1	858.4	0.000
{S} Species ¹	5	340.8	0.000
Time ²	1	379.8	0.000
{P} Pre-treatment ³	2	21.7	0.000
{T} Temperature ⁴	3	21.9	0.000
{L} Light ⁵	1	1779.1	0.000
Species*Pre-treatment	10	15.1	0.000
Species*Temperature	15	8.0	0.000
Species*Light	5	270.7	0.000
Pre-treatment*Temperature	6	0.5	0.818
Pre-treatment*Light	2	17.6	0.000
Temperature*Light	3	29.7	0.000
S*P*T	30	0.9	0.571
S*P*L	10	4.9	0.000
S*T*L	15	3.2	0.000
P*T*L	6	0.2	0.978
S*P*T*L	30	0.7	0.896
Error	1391		

1) *Bidens pilosa*, *Galinsoga parviflora*, *Guizotia scabra*, *Parthenium hysterophorus*, *Tagetes minuta* or *Verbesina encelioides*.

2) Germination tests after 0, 2.5, 5, 10, 20 and 30 weeks in pre-treatment.

3) Cold or warm stratification, or dry storage.

4) Germination test at 15/5, 20/10, 25/15 or 30/20°C day/night.

5) Germination test in light during daytime or in continuous darkness.

interaction "species*moment" (Table 2) shows species-specific differences in time in pre-treatment before most of the changes in germination occurred. The other significant second order interactions in the analysis of the differences resulted from species-specific differences in the extent of response to different pre-treatments, and in the extent to which a response occurred in darkness (Table 2, Fig. 2).

Third and higher order interactions, including the factor "species", were of minor importance for explaining the overall variation (Tables 1, 2), despite being statistically significant in some cases.

Table 2. ANOVA on weekly average difference between each adjacent germination test occasions when subjected to three different pre-treatments. Six *Asteraceae* species were included; each was collected from two sites and each seed batch was considered one replicate. Results including the factor "species" are in bold.

Factor	df	F	p
Intercept	1	286.0	0.000
{S} Species¹	5	10.3	0.000
{M} Moment ²	4	20.1	0.000
{P} Pre-treatment ³	2	7.0	0.001
{T} Temperature ⁴	3	1.1	0.334
{L} Light ⁵	1	86.9	0.000
Species*Moment	20	11.2	0.000
Species*Pre-treatment	10	4.3	0.000
Species*Temperature	15	1.5	0.107
Species*Light	5	8.2	0.000
Moment*Pre-treatment	8	10.9	0.000
Moment*Temperature	12	2.1	0.018
Moment*Light	4	7.5	0.000
Pre-treatment*Temperature	6	0.9	0.472
Pre-treatment*Light	2	4.1	0.017
Temperature*Light	3	0.6	0.585
S*M*P	40	5.8	0.000
S*M*T	60	1.5	0.016
S*M*L	20	5.3	0.000
S*P*T	30	0.5	0.992
S*P*L	10	1.0	0.411
S*T*L	15	1.3	0.199
M*P*T	24	1.2	0.269
M*P*L	8	3.0	0.003
M*T*L	12	1.1	0.335
P*T*L	6	0.2	0.985
S*M*P*T	120	1.2	0.097
S*M*P*L	40	3.1	0.000
S*M*T*L	60	1.6	0.004
S*P*T*L	30	0.2	1.000
M*P*T*L	24	1.2	0.235
S*M*P*T*L	120	1.1	0.183
Error	720		

1) *Bidens pilosa*, *Galinsoga parviflora*, *Guizotia scabra*, *Parthenium hysterophorus*, *Tagetes minuta* or *Verbesina encelioides*.

2) Moment in time between test occasions: 0-2.5, 2.5-5, 5-10, 10-20, or 20-30 weeks in pre-treatment.

3) Cold or warm stratification, or dry storage.

4) Germination test at 15/5, 20/10, 25/15 or 30/20°C day/night.

5) Germination test in light during daytime or in continuous darkness.

Figure 2. Average germination of two seed batches of six Asteraceae species. Seeds were subjected to three different pre-treatments (cold or warm stratification, or dry storage), and germination was tested at four daily fluctuating temperatures, both in light during daytime and in continuous darkness. Seeds of each species were collected from crop fields at two sites, ca. 5 kilometres apart, in eastern Ethiopia.

Differences between batches and species

In the PCA (Fig. 3), 57.6% and 20.1% of variation were explained by PC1 (the first Principal Component) and PC2, respectively. The two populations of each of *Galinsoga parviflora*, *Guizotia scabra*, *Parthenium hysterophorus* and *Tagetes minuta* clustered close to each other, while there were larger differences between the two batches of each of *Bidens pilosa* and *Verbesina encelioides* (Fig. 3).

Figure 3. Principal component analysis (PCA) of the combination of germination and weekly differences in response to pre-treatment of six Asteraceae species. Seeds were subjected to three different pre-treatments (cold or warm stratification, or dry storage), and germination was tested in eight environments after 0, 2.5, 5, 10, 20 and 30 weeks. Seeds were collected from crop fields at two sites, Haramaya University campus (open symbols) and Haramaya town (filled symbols), ca. 5 kilometres apart in eastern Ethiopia. Of the total variation, 57.6% and 20.1% are explained by PC1 and PC2, respectively.

Guizotia scabra, *Parthenium hysterophorus* and *Verbesina encelioides*

germinated faster and to a greater extent to cold stratification than to other pre-treatments, which either elicited a less pronounced response or no response (Fig. 4).

Galinsoga parviflora responded faster and to a higher extent to warm stratification than

Figure 4. Dormancy pattern and germination preferences of six annual weedy Asteraceae species collected in eastern Ethiopia. The left graphs show dormancy patterns as average germination after pre-treatment (cold or warm stratification, or dry storage) plotted against average time to a major response achieved. Average time and distribution bars on response time were weighted against the extent of each response. The right graphs show overall germination preferences to eight test environments (four daily fluctuating temperature regimes, with light during daytime or continuous darkness) as fractions of the environment that gave most germination. See text for detailed explanations.

to the other pre-treatments; *Bidens pilosa* responded fastest to warm stratification, but to the highest extent to cold stratification; and *Tagetes minuta* showed a response only to dry storage (Fig. 4). *Galinsoga parviflora* and *Verbesina encelioides* had the overall fastest responses, and *Bidens pilosa* and *Parthenium hysterophorus* the slowest (Fig. 4).

Germination preferences differed between species (Table 1). *Bidens pilosa* and *Galinsoga parviflora* germinated most at the highest temperatures tested, *Guizotia scabra* and *Parthenium hysterophorus* at 25/15°C, *Verbesina encelioides* at the lower temperatures, and *Tagetes minuta* had no clear temperature preference within the studied range (Fig. 4).

Seedling emergence and germination over time

After some rain at sowing in November, there was a dry period of nearly three months (Fig. 5). Most emergence during the 11-month observation period occurred during the middle of February, i.e. following the first precipitation (Fig. 5). Only *Guizotia scabra* did not emerge substantially to the first precipitation, but had an emergence peak during intense rains in late March and early April, as did *Bidens pilosa* and *Parthenium hysterophorus* (Fig. 5). *Guizotia scabra* and *Bidens pilosa* were the only species that emerged during May or later (Fig. 5).

When continuously moist-incubated in the light, *Tagetes minuta* germinated to nearly 100% at all temperatures within 8 weeks, and the few remaining seeds at 15/5°C germinated during 10 additional weeks (Fig. 6). *Verbesina encelioides* did not germinate at 30/20°C, and more slowly at 25/15°C than at 20/10 or 15/5°C, and *Parthenium hysterophorus* germinated mostly at 25/15°C (Fig. 6). *Guizotia scabra* and *Parthenium hysterophorus* germinated little at 15/5°C (Figs 4 and 6), while *Bidens pilosa* and *Galinsoga parviflora* germinated mostly after 18 weeks of incubation at

15/5°C; *Bidens pilosa* had its germination peak after 14 weeks of incubation at all temperatures (Fig. 6). When incubated for 34 weeks in darkness, only *Bidens pilosa* subjected to 30/20°C had germinated more than after four weeks of incubation (data not shown).

Figure 5. Cumulative average emergence of six Asteraceae species. Seeds were collected from crop fields at two sites, ca. 5 kilometres apart, in eastern Ethiopia. Seeds were sown in pots outdoors in Ethiopia, at ca. 2000 m a.s.l. on 20 November 2005, and were subjected to natural temperature and precipitation. Pots were checked at least every week; the first emergence was observed on 21 February 2006. Daily precipitation (bars), and maximum and minimum temperature (crosses) are shown.

When subjected to alternating periods of dryness and incubation, total germination after 24 weeks was higher than after continuous incubation for 34 weeks (Fig. 6). Compared to continuous incubation, *Guizotia scabra* and *Verbesina encelioides* increased germination at 30/20°C, and *Guizotia scabra* also increased at 15/5°C, after the second dry period.

Figure 6. Average germination of two seed batches of six Asteraceae species that were subjected to continuous incubation or to dry periods, alternating with incubation at different temperatures. Germination was checked at least every 6 weeks. Seeds of each species were collected from crop fields at two sites, ca. 5 kilometres apart, in eastern Ethiopia.

Discussion

Intra-species variation

Differences in dormancy and germination patterns within species may depend on genetic differences, local weather during growth of the mother plants, soil quality, seed position on the mother plant, or differences between seeds with different morphology. By collecting seeds available at a specific point in time, irrespective of variation within fruit heads, we describe the natural variation of two populations that have grown relatively close to each other in similar environments.

For *Galinsoga parviflora*, *Guizotia scabra*, *Parthenium hysterophorus* and *Tagetes minuta*, the overall relative difference within species was relatively small, while *Bidens pilosa* and *Verbesina encelioides* differed along PC1 (Fig. 3). Despite the occurrence of intra-species differences, "species" was an important explanatory factor for both germination and differences between test occasions (Tables 1, 2). Consequently, we conclude that it was meaningful to compare the general dormancy patterns and germination preferences of the species.

Comparison of species

When attempting to study differences in germination ecology between closely related species, groups within species and/or species with similar emergence patterns in the field (e.g. Keller and Kollman, 1999; Copete *et al.*, 2005; Bischoff *et al.*, 2006), dormancy classification systems are usually not used, but differences have been detected by ANOVA with seed batches as replicates (e.g. Tables 1 and 2; Pezzani and Montaña, 2006; Karlsson and Milberg, 2007c), by ordination (e.g. Fig. 3; Keller and

Kollman, 1999) or by comparing germination using variations within batches (e.g. Copete *et al.*, 2005; Bischoff *et al.*, 2006). These methods can detect differences and the magnitude of differences between compared units, but are not suitable tools for general, species/ecotype-specific descriptions. For making comparisons that are not dependent on specific testing points in time, logistic functions can be used to generalize germination results; this method allows descriptions of the principal response over time, rather than paying attention to seemingly random variations (e.g. Kettenring *et al.*, 2006; Karlsson and Milberg, 2007c). The selected point for measuring time to achieve a major response (Fig. 1) is the point where $f(t)$ has the highest "un-curving rate", i.e. when major change is fulfilled and the curve begins to flatten toward its upper asymptote. Thus, the amount of germination does not limit the calculation of response time.

When investigating dormancy and germination ecology, especially in cases when differences between batches are relatively small, we perceive a need for a method that summarises results from many tests (combinations of pre-treatment and incubation) and highlights possible differences. From an ecological point of view, such differences in responses between species as shown in Fig. 2 are of ecological interest, but this kind of result is difficult to use for comparisons and interpretations. Our intention was to visualize dormancy patterns and germination preferences in a condensed way, using continuous scales (Fig. 4). Positions on the x-axis show the weighted averages of response time, and positions on the y-axis average $f(30)$, i.e. the final magnitude of germination from Equation 1. In addition, average germination data for fresh seeds were plotted on the y-axis. The weighting of response time data (Equation 3) was done to allow the inclusion of all response times without giving too much influence to cases

where the response was weak (e.g., a change from 1% when fresh to 10% final germination was considered less important than an analogous change from 1% to 100%). The dispersal bars shown are not a tool for statistical comparisons, but are a way to give an impression of variation in the data. Points are included regardless of the extent of change in y-level; of course, if there is no obvious difference in relation to fresh seeds, the time to response is of no or minor ecological importance. For this kind of description to be meaningful, it is necessary that the experimental setup includes test environments where germination is possible, but does not always lead to full germination, thus allowing expressions of dormancy changes for all species. As shown in Fig. 2, this requirement was fulfilled for all species in the present study.

The dormancy patterns of *Bidens pilosa* and *Galinsoga parviflora* were quite similar regarding positive response to all pre-treatments, and they both responded faster to warm stratification than to the other pre-treatments (Fig. 4). However, they differed in that *Galinsoga parviflora* reduced dormancy much quicker, *Bidens pilosa* germinated relatively less at the two lowest temperatures tested and *Galinsoga parviflora* germinated exclusively when provided with light (Fig. 4).

Guizotia scabra, *Parthenium hysterophorus* and *Verbesina encelioides* all responded more quickly and more distinctly to cold stratification than to other pre-treatments (Fig. 4). However, there were considerable differences in dormancy patterns among these species: *Guizotia scabra* responded nearly as well to dry storage and warm stratification as to cold stratification, *Parthenium hysterophorus* did not respond to dry storage, and *Verbesina encelioides* did not respond to warm stratification (Fig. 4). Both *Guizotia scabra* and *Parthenium hysterophorus* germinated the most at 25/15°C, while

Verbesina encelioides had a clear preference for the two coolest test environments (Fig. 4).

Tagetes minuta differed from the other species in that dormancy was only substantially reduced when subjected to dry pre-treatment (Fig. 4). The species was strongly favoured by light for germination (Fig. 4), but, in contrast to the other species, germinated to the same magnitude at all temperatures within the range studied (Fig. 6).

Germination and emergence ecology

Because five out of six species responded to two or three pre-treatments (Fig. 4), we conclude that time *per se* is as important as specific environmental events for dormancy reduction for these species. Furthermore, considering the relatively small seasonal temperature differences in the study area and in the areas of origin for the species, it is noteworthy that all but *Tagetes minuta* clearly responded to cold pre-treatment (Fig. 4). This suggests that this dormancy reduction pattern is an evolutionary conservative character. The response could remain from hypothetical ancestral populations for which dormancy reduction during cold periods was favourable. The response could also be a result of evolution of a seed dormancy pattern with distribution of germination of each cohort over time, regardless of current environment. Thus, dormancy reduction during a cold period may have been a response neither useful nor harmful during the evolution of these species.

Guizotia scabra originates from East Africa (Baagøe, 1974), and seems not to have been dispersed from that area. *Verbesina encelioides* originates from the region of Mexico and the southern U.S. (Parsons and Cuthbertson, 2001), *Parthenium hysterophorus* from the Caribbean (Parsons and Cuthbertson, 2001) and the other

species from South America (Holm *et al.*, 1997; Bromilow, 2001; Parsons and Cuthbertson, 2001). The six species co-occur in various combinations as annual weeds in crop fields in highland, eastern Ethiopia (personal observation). With the exception of *Verbesina encelioides*, the species were among the 15 most common of 102 weedy species investigated in eastern Ethiopia in 1998 (Tamado and Milberg, 2000), among the more abundant species in sugarcane in central Ethiopia (Firehun and Tamado, 2006), and were included among 107 important weeds in Ethiopia as a whole (Stroud and Parker, 1989). *Verbesina encelioides*, which was rare in the area in 1998, had increased considerably by 2005 (personal observation). *Bidens pilosa*, *Galinsoga parviflora*, *Parthenium hysterophorus*, *Tagetes minuta* and *Verbesina encelioides* occur as serious, principal or common weeds around the southern hemisphere and north to Mexico, and *Galinsoga parviflora* and *Tagetes minuta* are also present in the U.S. (Holm *et al.*, 1979, 1997; Parsons and Cuthbertson, 2001). *Galinsoga parviflora*, *Tagetes minuta* and *Verbesina encelioides* are reported from 60°N in Norway, but only *Galinsoga parviflora* is established there (Lid *et al.*, 2005).

Despite the different origins and present distribution of the species, they co-occur as weeds in highland eastern Ethiopia, with similar periods for seed dispersal and emergence (Fig. 5). The weather during the study period in Ethiopia (Fig. 5) was normal for that time of the year; thus, seeds dispersed in autumn are normally subjected to a relatively long dry period before being imbibed for a number of consecutive days. Most emergence in pots occurred soon after the first rains in February (Fig. 5), similar to the germination after 12 weeks of storage before the onset of incubation (Fig. 6). In the field, there was a second peak of emergence in late March to early April (Fig. 5). From comparisons of continuous incubation to incubation with dry periods within (Fig. 6),

that response seems to be more an effect of time than of the actual dry periods *per se*, even though *Guizotia scabra* and *Verbesina encelioides* increased germination at 15/5 and 30/20°C, respectively, when dry periods were present.

In the field, in climates without pronounced temperature differences during the year, emergence in irrigated fields and in fields subjected to dry periods is probably about the same, according to the total amount of seedlings for these species, but emergence is more extended in time when fields are continuously irrigated (Fig. 6). Dry periods can increase the fraction of seeds germinating in darkness [buried seeds in the field, e.g. *Tagetes minuta* (Fig. 4)], thus reducing the soil seed bank and momentarily increasing the number of seedlings, compared to continuously irrigated fields. Seed germination of these species was generally stimulated by light (Fig. 4), and therefore, soil disturbance affects emergence by bringing seeds to soil surface (favouring germination), or by burying seeds, preventing germination.

The emergence timing in the field where these seeds were collected and sown can be explained by dormancy reduction during a dry period for *Bidens pilosa*, *Galinsoga parviflora*, *Guizotia scabra* and *Verbesina encelioides*, even though a dry period was not the most efficient environmental circumstance for dormancy reduction for any of them (Fig. 4). For imbibed *Tagetes minuta*, the emergence pattern was a result of germination that readily occurred in the light at the temperatures studied, and for *Parthenium hysterophorus*, it was mostly a matter of elapsed time from first imbibition (Fig. 6). Thus, the six investigated species have different dormancy patterns (Fig. 4), but achieve similar emergence timing in the field following a dry period after dispersal (Fig. 5), leading to possible co-occurrence in environments similar to those studied.

The different dormancy patterns (Fig. 4) indicate differences in possible distribution ranges or emergence timing in environments with larger temperature differences and less pronounced dry periods than at the collection site (Fig. 5). From the dormancy patterns, one can suspect that both *Parthenium hysterophorus* and *Bidens pilosa*, and other species with similar dormancy pattern and germination preferences, are able to establish in colder climates than their present range without any further adaptation. Because of a relatively strong dormancy, a reduction of dormancy in response to cold periods, and restricted germination at lower temperatures (Fig. 4), a substantial part of a seed cohort, produced from an initial successful plant flowering during summer, could disperse in autumn, but not germinate at the prevailing low temperatures. The degree of dormancy would be reduced during winter, and seeds would germinate in early summer, thus becoming established.

Acknowledgements

We thank Abdulrazak Abdulahi, Almaz Alemu and co-workers, Daniel Wondimu, Mekonnen Takaro and Peter Reneby, for assistance. Funding was provided by Formas (The Swedish Research Council for Environment, Agricultural and Spatial Planning).

References

- Baagøe, J.** (1974) The genus *Guizotia* (Compositae) - A taxonomic revision. *Botanisk Tidsskrift* **69**, 1-39.
- Baskin, J.M. and Baskin, C.C.** (2004) A classification system for seed dormancy. *Seed Science Research* **14**, 1-16.
- Bischoff, A., Vonlanthen, B., Steinger, T. and Müller-Schärer, H.** (2006) Seed provenance matters – Effects on germination of four plant species used for ecological restoration. *Basic and Applied Ecology* **7**, 347-359.
- Bromilow, C.** (2001) *Problem plants of South Africa*. Pretoria, Briza Publications.
- Copete, M.A., Herranz, J.M. and Ferrandis, P.** (2005) Seed dormancy and germination in threatened Iberian *Coincya* (Brassicaceae) taxa. *Ecoscience* **12**, 257-266.
- Firehun, Y. and Tamado, T.** (2006) Weed flora in the Rift Valley sugarcane plantations of Ethiopia as influenced by soil types and agronomic practises. *Weed Biology and Management* **6**, 139-150.
- Harper, J.L.** (1957) The ecological significance of dormancy and its importance in weed control. *Proceedings of the 4th International Congress of Crop Protection* **1**, 415-420.
- Harper, J.L.** (1977) *Population biology of plants*. London, Academic Press.
- Holm, L., Pancho, J.V., Herberger, J.P. and Plucknett, D.L.** (1979) *A geographical atlas of world weeds*. New York, John Wiley & Sons.
- Holm, L., Doll, J., Holm, E., Pancho, J. and Herberger, J.** (1997) *World weeds: Natural histories and distribution*. New York, John Wiley & Sons.

- Karlsson, L.M. and Milberg, P.** (2007a) A comparative study of germination ecology of four *Papaver* taxa. *Annals of Botany* **99**, 935-946.
- Karlsson, L.M. and Milberg, P.** (2007b) Seed dormancy pattern and germination preferences of the South African annual *Papaver aculeatum*. *South African Journal of Botany* **73**, 422-428.
- Karlsson, L.M. and Milberg, P.** (2007c) Comparing after-ripening response and germination preferences of *Conyza canadensis* and *C. bonariensis* (Asteraceae) through logistic functions. *Weed Research* (in press).
- Karlsson, L.M. and Milberg, P.** (200X) Variation within species and inter-species comparison of seed dormancy and germination of four annual *Lamium* species. *Flora* (in press).
- Keller, M. and Kollman, J.** (1999) Effects of seed provenance on germination of herbs for agricultural compensation sites. *Agriculture, Ecosystems and Environment* **72**, 87-99.
- Kettenring, K.M., Gardner, G. and Galatowitsch, S.M.** (2006) Effect of light on seed germination of eight wetland *Carex* species. *Annals of Botany* **98**, 869-874.
- Lang, G.A.** (1987) Dormancy: A new universal terminology. *HortScience* **22**, 817-820.
- Lid, J., Lid, D.T. and Elven, R.** (2005) *Norsk flora* (7th edition). Oslo, Det Norske Samlaget, (in Norwegian).
- Merritt, D.J., Turner, S.R., Clarke, S. and Dixon, K.W.** (2007) Seed dormancy and germination stimulation syndromes for Australian temperate species. *Australian Journal of Botany* **55**, 336-344.
- Parsons, W.T. and Cuthbertson, E.G.** (2001) *Noxious weeds of Australia* (2nd edition) Collingwood, CSIRO Publishing.

- Pezzani, F. and Montaña, C.** (2006) Inter- and intraspecific variation in the germination response to light quality and scarification in grasses growing in two-phase mosaics of the Chihuahuan desert. *Annals of Botany* **97**, 1063-1071.
- StatSoft, Inc.** (2004) STATISTICA (data analysis software system), version 7.
www.statsoft.com.
- Stroud, A. and Parker, C.** (1989) *A weed identification guide for Ethiopia*. Rome, Food and Agriculture Organization of the United Nations.
- Tamado, T. and Milberg, P.** (2000) Weed flora in arable fields of eastern Ethiopia with emphasis on the occurrence of *Parthenium hysterophorus*. *Weed Research* **40**, 507-521.
- Vleeshouwers, L.M., Bouwmeester, H.J. and Karssen, C.M.** (1995) Redefining seed dormancy: an attempt to integrate physiology and ecology. *Journal of Ecology* **83**, 1031-1037.