# Surgical treatment in chronic aortic regurgitation

Timing, results, prognosis and left ventricular function

Éva Tamás



Division of Cardiovascular Medicine Department of Medical and Health Sciences Linköping University, Sweden

Linköping 2008

Cover: Éva Tamás & Dennis Netzell Published articles have been reprinted with the permission of the copyright holder. Printed by Liu-Tryck, Linköping, Sweden, 2008 ISBN 978-91-7393-857-0 ISSN 0345-0082 Copyright © 2008 by Éva Tamás No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanic, photocopying, recording or otherwise, without prior permission of the author.



# **CONTENTS**

ABSTRACT	1
LIST OF ORIGINAL PAPERS	3
ABBREVIATIONS	4
INTRODUCTION	5
The aortic root and the left ventricle	6
The aortic root	6
Left ventricle: anatomy & function	7
Methods of clinical examination	8
Ultrasound imaging	8
Radionuclide ventriculography	
with Multiple Gated Image Acquisition (MUGA)	10
Cardiopulmonary Exercise Testing (CPET)	11
Chronic aortic regurgitation	12
Etiology & prevalence	12
Natural history & timing of surgery	13
AIMS OF THE STUDY	17
METHODS	18
Patients	18
Paper I	18
Papers II & III	18
Paper IV	19
Echocardiography	20
MUGA	21
CPET	22
Statistical methods	23
Ethical considerations	24

RESULTS	25
Survival analysis (I)	25
Left ventricle (I-III)	26
Left ventricular diameters and volumes	26
Left ventricular function at rest & during exercise	27
Left ventricle and aortic valve surgery	29
Physical work capacity (III)	29
Echocardiographic anatomy of the normal aortic valve (IV)	30
DISCUSSION	32
Aspects of pathophysiology, prevalence & gender	32
Survival after surgery for chronic AR	
Compensated & decompensated left ventricle in chronic AR	33
Effects of AR on physical work capacity	35
Timing of surgery	37
Clinical anatomy of the aortic valve	37
CONCLUSIONS	40
CLINICAL IMPLICATIONS	41
ACKNOWLEDGEMENTS	42
REFERENCES	44

# **ABSTRACT**

Chronic aortic regurgitation (AR) of varying degree affects 13% of men and 8.5% of women. In persons with severe AR, the expected length of life and its quality are influenced. Some individuals remain asymptomatic for a long period, due to effective compensatory mechanisms, but dysfunction of the left ventricle (LV) usually begins before symptoms appear and can be irreversible by then. This thesis addresses questions of LV function and optimal time for operation of patients suffering from chronic AR. Moreover, detailed echocardiographic studies of the anatomy of the normal aortic valve have been performed to obtain a better understanding of the in vivo anatomic relations within the aortic root.

Patients with chronic AR, without concomitant cardiac disease, were studied both retrospectively (n=88) and prospectively (n=29) and the aortic valves of persons (n=32) free from cardiac disease were investigated.

For the retrospectively studied patients, survival was 82% at 10 years which is an improvement compared with previously published results. The majority of the patients, however, had LV dysfunction preoperatively. By patients prospectively by echocardiography, ventriculography (MUGA) and cardiopulmonary exercise testing (CPET) our aim was to evaluate the predictive value of measurements of LV function at rest and during exercise for postoperative outcome. LV diameters were markedly elevated prior to and diminished significantly after surgery. Patients with an abnormal exercise ejection fraction (EF) response by MUGA preoperatively, presented the same reaction postoperatively. This could not be predicted by LV function determination at rest, or by NYHA functional class. In spite of median NYHA class II, these patients had a low work capacity on CPET, which was neither improved 6 months postoperatively nor correlated to echocardiographic LV dimensions. Thus, both MUGA and CPET may be useful complements for timing of surgery in patients with chronic AR.

Assuming that patients would benefit from preservation of their native valves the normal aortic valve was studied to gain detailed information about the echocardiographic anatomy and relations within the normal aortic root. This extended examination of the aortic root may facilitate a better planning of aortic valve-preserving interventions in the future.

# LIST OF ORIGINAL PAPERS

This thesis is based on the following papers, referred to in the text by their Roman numerals

- I. Tamas E, Nylander E, Olin C. Are patients with isolated chronic aortic regurgitation operated in time? Analysis of survival data over a decade. *Clin Cardiol*. 2005 *Jul*;28(7):329-32.
- II. Tamas E, Broqvist M, Olsson E, Franzén S, Nylander E. Exercise radionuclide ventriculography for predicting postoperative left ventricular function in chronic aortic regurgitation. Submitted manuscript.
- III. Tamas E, Nielsen NE, Vanhanen I, Nylander E. Measurement of physical work capacity in patients with chronic aortic regurgitation: A potential improvement in patient management *Submitted manuscript*.
- IV. Tamas E, Nylander E. Echocardiographic Description of the Anatomic Relations within the Normal Aortic Root. J Heart Valve Dis 2007;16:240-246

# **ABBREVIATIONS**

AR Aortic Regurgitation
BMI Body Mass Index
BSA Body Surface Area

CPET Cardiopulmonary Exercise Testing
EF Left Ventricular Ejection Fraction

ΔEF Left ventricular Ejection Fraction Response to Exercise

ICD Intercommissural Distance

LC Left Coronary

LV Left Ventricular LVF Left Ventricular Function

LVED Left Ventricular End-Diastolic Diameter
LVES Left Ventricular End-Systolic Diameter
LVEDV Left Ventricular End-Diastolic Volume
LVESV Left Ventricular End-Systolic Volume

maxSiD Maximal Sinus Diameter

MUGA Multiple Gated Image Acquisition

NC Non-Coronary

NYHA New York Heart Association

OV Opening of the Valve

RC Right Coronary

RPE Rate of Perceived Exertion

SAD Subaortic Diameter STJ Sinotubular Junction

TEE Transesophageal Echocardiography

V O2 peak Peak Oxygen Uptake

# Introduction

"The heart ... is the beginning of life; the sun of the microcosm... for it is the heart by whose virtue and pulse the blood is moved, perfected, made apt to nourish, and is preserved from corruption and coagulation; it is the household divinity which, discharging its function, nourishes, cherishes, quickens the whole body, and is indeed the foundation of life, the source of all action."

William Harvey's lines from 1628 can be considered to mark the beginning of an era of extensive studies on the heart and the circulation. Much has been learned since then but we are still far away from being able to describe and understand anatomy and hemodynamics completely.

The studies presented in this thesis address questions of left ventricular function and optimal time for operation of patients suffering from chronic aortic regurgitation; moreover detailed echocardiographic anatomy of the normal aortic valve is presented.

#### The aortic root and the left ventricle

#### The aortic root

The aortic valve is located between the left ventricle and the aorta. Unfortunately, completely different opinions about the nomenclature of this region of the heart still remain<sup>2-4</sup>. The three semilunar cusps are usually referred to as the aortic valve. However, these cusps together with the upper part of the left ventricle and the aortic wall form a functional entity called the aortic root. The three half-moon formed cusps are connected to the aortic wall in such a manner that the deepest part of the pocket formed by the aortic wall and a cusp is in the left ventricular outflow tract. The lower border of the cusps together with the aortic wall forms a fibrous crown which can be interpreted as the annulus<sup>4</sup> in spite of the fact that the Latin word means 'ring'. (Fig.1) Others define the aortic annulus as an imaginary ring laid through the deepest points of the cusps. <sup>5</sup>



Figure 1
Aortic root. A: The crown shaped conception of the annulus is based on histopathological studies. B: Surgical interpretation of the annulus.

It is considered common knowledge that the aortic valve is competent when the upper segments of the cusps meet, and this is what prevents blood from regurgitating out of the aorta to the left ventricle. The band-like area where the cusps meet bears the name of coaptation area. The meeting post between two cusps on the aortic wall is called commissure. The aortic wall bulges between two commissures and the aortic sinus is formed. These sinuses have a role in coronary perfusion through a special pattern of blood flow due to their geometry. Leonardo DaVinci described and showed this pattern but the exact function and mechanism is still unknown. Hence, it is difficult to say how important the sinuses are. <sup>6-9</sup>

As the limit of the aortic root is the deepest point of the cusps towards the left ventricle, the upper limit is the sinotubular junction (STJ) where the sinuses and the commissures end and the aorta takes the shape of a tube.

#### Left ventricle: anatomy & function

The dominating tissue of the left ventricle is muscle. The fibres of the myocardium are organized in three layers, where the fibres run in different directions: longitudinally under the endocardium, circumferentially in the middle and obliquely close to the epicardium. The longitudinal fibres of the inner muscle layer perform the long axis movement of the heart.

The phase during which the left ventricle is filled with blood is called diastole (*Greek. 'act of dispending or dilating'*) while the emptying is called systole (*Greek. 'contraction'*). The original words describe very well what happens to the heart muscle during these phases. Intact contractility, in other words ability to perform contraction and have time to relax between contractions, is the basis of myocardial function. The total function of the left ventricle is a counteraction of systolic and diastolic function of the myocardial fibres, as dilatation is as important as contraction in order to be able to present good pump function and thereby sufficient circulation for the body. <sup>6,10</sup>

#### Methods of clinical examination

Measuring contractility in vivo in humans is difficult with standard diagnostic tools but evaluation of function of which contractility is an important component is possible with different methods. Ejection fraction (EF) is one measure of systolic function.

$$EF (\%) = \frac{LVEDV - LVESV}{LVEDV} \times 100$$

where LVEDV is left ventricular end-diastolic volume and LVESV is left ventricular end-systolic volume. Nevertheless, EF is influenced by e.g. ventricular volume and valvular regurgitation. Another tool for assessment of systolic function is the longitudinal function. <sup>10, 11</sup> The diastolic function can be estimated by measuring blood flow velocity and inflow pattern to the left heart. The ideal technique for examination should be non-invasive, informative, reproducible, reliable and cost-effective.

#### Ultrasound imaging

Sound is a pressure wave and its frequency is the number of cycles per second (Hertz; Hz). The range which has higher frequencies than are audible for humans is referred to as ultrasound (≥ 20 kHz). Imaging ultrasound frequencies range from 2 MHz to 40 MHz. Transducer crystals are used to generate ultrasound. An electric impulse causes expansion of the crystal and the adjacent tissues are compressed. After expansion, the crystal contracts and a pressure wave is produced that travels through the tissues by the transfer of the pressure wave from one vibrating particle to the next, in the direction of the wave-front. Ultrasound passes through homogenous tissue without change in velocity (1540 m/s in soft tissue). As the sound beam passes through tissue, it is reflected, refracted and absorbed. Reflection is utilized for creating images, hence the name for cardiac imaging is echocardiography (*Greek. echo – 'reflected sound'*).

Ultrasound technique was first used during World War II and documentation of medical application came first in 1952. Edler in Lund recorded cardiac motion and published this in 1953. However, only the expansion of computer technology made it possible to create compact and fast

equipment for echocardiography and its extensive use could begin in the middle of 1980s.

Two dimensional (2D) images show real-time anatomical images of cardiac structures in a scanned sector of the heart.

M-mode (time – motion) imaging is based on information collected along one scan line only. This results in a higher sampling rate and a very accurate presentation of movement for a particular structure. The position of the M-mode line is usually guided by a 2D image.

Doppler studies (continuous, pulsed wave and colour coded) are used to measure blood flow characteristics. Measurements are based on the Doppler-effect (J.C. Doppler, Austrian physicist, 1803-1853) which is described as a frequency shift when the radiation source is moving.

$$\Delta f = \frac{2vfcos\theta}{C}$$
 where v is the velocity of the blood, f is the emitted frequency,  $\theta$ 

is the angle between the blood flow and the ultrasound beam and C is the velocity of the ultrasound in soft tissue. The velocity can be calculated as

$$v = \frac{\Delta fC}{2f\cos\theta}$$
. Maximum value for  $\cos\theta$  is  $\cos0^\circ = 1$  which means that maximum

velocity is detected when blood flow is parallel to the ultrasound beam. A  $\theta$  = 20° gives a 6% difference in blood velocity which is accepted as the error limit. This means that the measured velocity is practically always less than the real value. Pulsed wave doppler is used when blood flow velocity is to be measured at a certain depth.




Figure 2
2D and colour flow doppler images of a regurgitant aortic valve.

Continuous wave doppler is used to measure high velocities. Measuring the velocities makes it possible to give a good approximation of pressure and thereby flow through regurgitant valves according to the simplified Bernoulli equation  $P = 4(v_2^2-v_1^2)$ , where P is the pressure drop,  $v_2$  is the peak flow through the valve and  $v_1$  is the flow just prior to the valve.

Colour coded doppler images give information about flow characteristics. The form, direction and magnitude of the jet are basic pieces of information in grading regurgitant valves but one has to bear in mind that this is still a qualitative method.<sup>10-12</sup>

Radionuclide ventriculography with Multiple Gated Image Acquisition (MUGA)

This method is based on the principle that radioactivity of a certain amount of injected isotope marked red blood cells in a blood pool is proportional to the volume of the blood pool. Radioactivity is detected by NaI crystals in a gamma camera. These crystals work as scintillation detectors and capture  $\gamma$ -radiation with high sensitivity and good resolution. In order to screen radiation beams not perpendicular to the detector, a plate of lead with wholes is placed in the way of the radiation. With absorption of radiation, the scintillation detector gives a light signal. This signal is captured by a photomultiplicator which transforms it into an electric signal and gives information about position in the X and Y directions. The position of a third position signal (Z) is captured by a pulse height analysator (PHA). The PHA accepts the energy signal only if it lies within a previously defined range. This makes it possible to filtrate radiation of unknown origin and thereby improve image contrast and geometrical resolution. The radioactive impulses on the detector are counted (counts), collected and stored by a computer. A certain number of counts are necessary in order to achieve a good image quality and to provide a reliable material for analysis of the blood pool in the imaged organ.

Digital images are stored in the form of a picture matrix, generally 64x64 pixels. Measurements in nuclear medicine are based on evaluation of relative activity within a certain region called region of interest (ROI). The ROI can be delineated manually by an operator or automatically by the computer. Background radiation from blood flow through other tissues than the organ of interest affects analysis of radioactivity therefore background activity is to be subtracted. Images are normalized and filtered to diminish variation and noise, and to enhance contrast.




Figure 3 Calculation of EF by MUGA.

## Cardiopulmonary Exercise Testing (CPET)

The primary function of the cardiopulmonary system is to provide oxygen (O<sub>2</sub>) supply and carbon dioxide (CO<sub>2</sub>) elimination. Under steady state conditions, the external respiration (lungs) is equivalent to internal (cell) respiration. Exercise testing using a treadmill or bicycle protocol is commonly used to test exercise capacity. Considerable amounts of additional information can be gained through registration and analysis of respiratory gases. CPET records data about the cardiovascular and respiratory systems under stress

simultaneously, which makes it possible to identify whether the reason for the patient's symptoms is respiratory or cardiac. Physical exercise needs a higher level of oxygen supply  $(\dot{V}O_{2\,peak})$  in order to be able to produce the necessary energy for muscle work and results in CO<sub>2</sub> accumulation, which must be eliminated to avoid acidosis.

# Chronic aortic regurgitation

#### Etiology & prevalence

Aortic regurgitation arises when the semilunar cusps between the left ventricle and aorta do not close effectively and a certain volume of blood regurgitates to the left ventricle. When this defect develops and is sustained for a longer period of time, it is called chronic aortic regurgitation. The possible etiology is presented in Table 1.

The prevalence of chronic aortic regurgitation or, as often also referred to in the literature, aortic insufficiency, is 13.0% in men and 8.5% in women, as reported in The Framingham Heart Study in 1999.<sup>13</sup>

The regurgitant blood volume causes volume overload in the left ventricle. The total stroke volume (sum of effective stroke volume and regurgitant volume) is increased. In order to be able to achieve a normal effective stroke volume the heart increases the LV end-diastolic volume. This causes elevation of LV end-diastolic pressure but, nevertheless, LV function can be normal at this stage and the patient is often free of any symptoms. As the illness progresses, a further increase of the LV end-diastolic pressure and impairment of LV function are to be expected. In order to be able to maintain stroke volume, the left ventricle compensates with eccentric hypertrophy. There is an increased demand on oxygen supply due to the volume overload, higher left ventricular pressure and hypertrophy and, at the same time, it is exactly these factors that are limiting coronary perfusion.

1.	Inflamm	ation	
	1.1.	Endocarditis lenta	
	1.2.	Ankylosing spondylitis	
	1.3.	Rheumatic disease	
	1.4.	Rheumatiod arthritis	
	1.5.	Giant cell aortitis	
	1.6.	Syphilitic aortitis	
2.	Tissue d	egeneration	
	2.1.	Marfan's syndrome (cystic medianecrosis)	
	2.2.	Ehlers-Danlos syndrome	
	2.3.	Osteogenesis imperfecta	
	2.4.	Reiters syndrome	
	2.5.	Myxomatous degeneration	
3.	Congeni	tal	
	3.1.	Bicuspid aortic valve	
	3.2.	Outlet supracristal VSD	
	3.3.	Discrete subaortic stenosis	
Į.	Other		
	4.1.	Chronic severe systemic hypertension	
	4.2.	Anorectic drugs	
	4.3.	Idiopathic aortic root dilatation	

Table 1 Etiology of chronic aortic regurgitation<sup>6, 14-17</sup>

#### Natural history & timing of surgery

Less than 6% of asymptomatic patients with normal LVF present with symptoms per year. Nevertheless, 3.5% per year develop LV dysfunction and 0.2 suffer sudden death. Of those who already have LV dysfunction, 25% progress to symptoms and 10% of those who became symptomatic die every year. 16, 17

In order to avoid irreversible deterioration of LVF, aortic valve replacement or reconstruction is the state of the art treatment for symptomatic patients. As surgery is unavoidable for these patients, natural history has not been studied on larger groups of patients after the introduction of prosthetic valve implantation in the aortic position in 1953.

Criteria for indication of surgery	Class	Level of evidence
1. Symptomatic patient 1.1. Severe AR irrespective of LVF	I	В
2. Asymptomatic patient		
2.1. Severe AR		
2.1.1. LV dysfunction (EF < 50%)	I	В
2.1.2. Normal LV function (EF > 50%)		
2.1.2.1. LVED > 75 mm or LVES > 55 mm	IIa*	B*
2.1.2.2. LVED > 70 mm or LVES > 50 mm	IIb*/IIa⁺	$C^*/C^t$
2.1.2.2.1. progressive LV dilatation*		
2.1.2.2.2. declining exercise tolerance*		
2.1.2.2.3. abnormal hemodynamic		
responses to exercise*		
(consider lower threshold values for patients of small stature		
of either gender)		
3. Asymptomatic and symptomatic		
3.1. Severe & moderate chronic AR while	I	С
undergoing CABG or surgery on the aorta or		
other heart valves		
3.2. Whatever/moderate severity of AR	IIb*	C*
undergoing surgery on the ascending aorta		
3.2.1. patients with Marfan's syndrome <sup>+</sup>	I	C
3.2.2. bicuspid valves <sup>†</sup>	IIa	C

Table 2
Indications of surgery for patients with chronic AR according to present guidelines from the American College of Cardiology\* and the European Society of Cardiology\*. <sup>16, 17</sup>

The above mentioned compensation by the left ventricle keeps the majority of the patients asymptomatic and may disguise a beginning important dysfunction. Therefore, regular follow-up of LVF is recommended

in order to be able to operate before irreversible changes occur. Table 2 shows the indications for surgery for patients with chronic aortic regurgitation.

Finding the optimal time for surgical intervention has remained an art<sup>18-21</sup> during the years of aortic valve surgery, in spite of the evolution of diagnostic methods. Scandinavian studies<sup>22-24</sup> have shown that the relative survival for patients with chronic aortic regurgitation was 67% after 10 years. This group of patients has not only high expected mortality but a relatively high morbidity as well.

Natural history of chronic AR<sup>6, 13, 25</sup> tells us that predominantly young and middle aged persons, with an otherwise long life expectancy, suffer from this disease. A well chosen time for operation of these patients could be essential for their clinical outcome e.g. expected length of life. The "heart of the matter" would seem to be how to determine that point on the timeline when the LVF starts to deteriorate and whether the standard methods of today have the capacity to provide us with these pieces of information.

# AIMS OF THE STUDY

I. To analyze the preoperative LVF and survival data in patients suffering from chronic AR without concomitant heart disease, operated on during a ten year interval, 1989-1999.

#### II. To study

- i) the changes in the EF from rest to exercise, as a measure of LVF both pre- and postoperatively in patients with chronic AR without any other cardiac disease.
- ii) whether any of the measured preoperative LV dimensional or functional parameters could reliably predict LVF after surgery.
- III. To evaluate and compare the pre- and postoperative functional capacity of asymptomatic to moderately symptomatic patients with chronic AR by CPET.

#### IV. To investigate

- i) the aortic valve in detail by TEE, in order to describe the anatomical and, if possible, physiological properties of healthy human aortic valves in vivo.
- ii) if application of echocardiography is reproducible and reliable for measurements of the fine details in the clinical anatomy of the aortic valves.

## **METHODS**

#### **Patients**

#### Paper I

A cohort of patients (n=88) operated on for chronic isolated aortic regurgitation during a ten-year period (1989-1999) was studied. Concomitant cardiac disease was an exclusion criterion. General patient characteristics are indicated in Table 3. Most of our patients were middle aged men with body surface area (BSA) and body mass index (BMI) within normal ranges. The women were somewhat older, had generally lower BSAs and were slightly overweight. The majority of this patient population was moderately symptomatic. All included patients had hemodynamically highly significant aortic regurgitation, judged to need surgical treatment based on clinical signs and echocardiography results.

#### Papers II & III

Consecutive patients (n=29) with severe chronic aortic regurgitation were included in this study. Patients with concomitant coronary artery disease and with other heart valve disease than chronic aortic regurgitation were excluded. Aortic stenosis was defined as aortic orifice area  $\leq 1.6~\rm cm^2$ . Moreover, patients who had any history of congenital heart disease or active endocarditis were excluded. Healed endocarditis was not an exclusion criterion. Patients were asymptomatic to moderately symptomatic in this population. Transthoracic echocardiography and MUGA, both at rest and during exercise, were performed 1-2 days before the planned surgery and 6 months postoperatively.

Twenty-six patients completed the study. The reasons for three patients being lost to follow up were: moved abroad, mild claustrophobic symptoms and abstained from the postoperative study, and one patient suffered perioperative stroke with incapacitating neurological symptoms. Preoperative data of these patients were analyzed. In Paper III, the patient in NYHA IV was not included.

Paper IV

Patients (n=32) referred for TEE for suspected cardiac sources of emboli, but diagnosed as normal, were included in this cross-sectional study. Exclusion criteria were concomitant heart valve disease, endocarditis (active as well as healed), history of myocardial infarction or any cardiac abnormality detected during the echocardiographic study, except for trivial valve regurgitation.

	Age (yrs)	BSA (m <sup>2</sup> )	BMI (kg/ m²)	NYHA
Paper I				
All patients	$56 \pm 13$	$1.9 \pm 0.2$	$21.0 \pm 2.0$	III (I-IV)
(n=88)				
Men (n=76)	$55 \pm 14$	$2.0 \pm 0.2$	20.6 ±1.7	II (I-IV)
Women (n=12)	$65 \pm 10$	$1.6 \pm 0.1$	$23.4 \pm 2.2$	III (I-IV)
Paper II				
Men (n=29)	$51 \pm 14$	$2.0 \pm 0.2$	$27.4 \pm 3.1$	II (I-IV)
Paper III				
Men (n=28)	$50 \pm 13$	$2.0 \pm 0.2$	$27.1 \pm 2.8$	II (I-III)
Paper IV				
All patients	$54 \pm 14$ .	$1.9 \pm 0.2$	$24.9 \pm 3.6$	-
(n=32)				
Men (n=20)	$55 \pm 13$	$2.0 \pm 0.1$	$24.7 \pm 3.1$	-
Women (n=12)	$53 \pm 17$	$1.7 \pm 0.2$	$25.3 \pm 4.3$	-

Table 3
Demographic data & NYHA functional class

# Echocardiography

Global LV function was evaluated using a 4-grade scale, where normal systolic function corresponds to an EF≥50%, slight dysfunction to EF 40-49%, moderate dysfunction to 30-39% and severely reduced function corresponds to EF<30%.<sup>26</sup> The LV diameter was measured in end-diastole and end-systole by M-mode. The aortic root diameter at the level of cusp closure was measured (M-mode) in Paper I.

For transthoracic ultrasound scans a GE Vingmed, Vivid 5 or a Vivid 7 echocardiograph was used in Papers II & III. Images were recorded, stored digitally and analysed off-line.

LV diameters in end-diastole, and in end-systole (LVED, LVES) and LV wall thickness in end-diastole were measured by M-mode and LV mass was calculated according to Devereaux's formula. LV end-diastolic volume (LVEDV), left ventricular end-systolic volume (LVESV) and EF according to Simpson's rule were calculated. Volume measurements, an average of three tracings in each registration, were performed by an experienced investigator who was blinded for the EF results of MUGA.

The velocity of E and A waves, E/A ratio and pulmonary venous systolic and diastolic velocity were analysed and used for an integrated description of diastolic function.<sup>27, 28</sup>

In Paper IV, the echocardiograms were recorded using a GE Vivid 5 instrument fitted with a multiplane transesophageal probe. Two-dimensional, electrocardiogram-synchronized cine loops of the aortic valve in the long axis (100-120°) and short axis (45-60°) views, both in mid-systole and end-diastole, were stored digitally and measurements carried out off-line. After scrolling to identify optimal images in end-diastole (defined as the beginning of the QRS) and mid-systole (when the valve was most wide open, coinciding with the top of the T-wave), parameters were measured using the inner edge technique. This method was chosen in order to ensure uniformity for both horizontal and vertical measurements. Anatomic parameters of the entire aortic apparatus, displayed in Fig. 4, were measured three times and the mean value used in the statistical analyses. Two investigators performed the same series of measurements, and this dataset was analyzed for repeatability and reproducibility.




Figure 4
Aortic root and cusp parameters.Panel A: long axis view, diastole, Panel B: long axis view, systole, Panel C: short axis view diastole, Panel D: short axis view, systole.

Identification numbers for parameters: 1, 2: subaortic diameter; 3, 4: maximal sinus diameter; 5, 6: sinotubular junction; 7: cusp separation (valve opening); 8: coaptation; 9, 10: left cusp height; 11, 12: right cusp height; 13, 14: left sinus height; 15, 16: right sinus height; 17, 18: intercommissural distance left coronary sinus; 19, 20: intercommissural distance right coronary sinus; 21, 22: intercommissural distance non-coronary sinus; 23, 24: left coronary cusp diameter; 25, 26: right coronary cusp diameter; 27, 28: non-coronary cusp diameter.

# **MUGA**

Erythrocytes were labelled with 10 MBq/kg Tc-99m pertechnetate using the modified in-vitro technique. Electrocardiogram gated imaging was performed using a General Electric XR/T gamma camera (General Electric Medical Systems, Milwaukee, USA) equipped with a general purpose, low energy collimator and a Genie ACQ collecting station. For optimal separation of the left and the right ventricles, left anterior oblique (LAO 45°) positioning was

used. Supine multigated blood pool imaging was performed first at rest. After this, the patients performed supine bicycle exercise at a pedal rate of 60/min. Besides continuous ECG and heart rate registration, blood pressure was also measured every 2 minutes. The initial workload was 50 W and was increased by 50 W at the next level.

Xeleris Functional Imaging Workstation, EF Analysis (General Electric Medical Systems, Milwaukee, USA) was used for image and data analysis. Five Mcounts in total were collected at rest and 2.5 Mcounts during steady state exercise, with 32 frames per heart cycle in a matrix size of 64 x 64. Left ventricular end-diastolic and end-systolic outlines were marked manually by each of two operators separately, who made 3 measurements each. The mean of these measurements was used. The automatic algorithms of the computer program were used for background (BG) correction. Left ventricular ejection fraction (EF) was calculated as

$$EF(\%) = \frac{\left(LV\ counts\ diastole - BG\ counts\right) - \left(LV\ counts\ systole - BG\ counts\right)}{\left(LV\ counts\ diastole - BG\ counts\right)} \times 100\ .$$

Cycle length windowing with dynamic arrhythmia filtration<sup>29</sup> with forward gating was used. A heart rate (cycle length) histogram was generated and observed for 1-3 min. When steady state heart rate was achieved a window was centred round peak heart rate ±10% for patients with sinus rhythm. Beats outside this interval were rejected. In patients with atrial fibrillation the window of acceptable beats was widened up to 15-20% to reduce imaging time.

Subgroups of patients were formed based on preoperative EF exercise response ( $\Delta$ EF) during MUGA and were categorized as decreasing:  $\Delta$ EF < -5%, unaltered: -5%  $\leq \Delta$ EF  $\leq$  5% and increasing:  $\Delta$ EF > 5%. A 5% or higher increase of EF was considered normal. These subgroups were defined considering the above limits and a 5% bias of the method.<sup>30-32</sup>

#### **CPET**

An incremental symptom limited bicycle exercise test using an electrically braked bicycle ergometer (eBike, CE-0459, Ergoline, GE Medical Systems, Freiburg, Germany) at a pedalling rate of 60/min was performed by patients in Paper III. The exercise test was started at 50 Watts (W) steady state for 6

minutes. Thereafter, the workload was increased by 10 or 20 W every min until exhaustion or appearance of any other limiting symptom.

Expired gas was continuously collected through a low resistance tightly fitted mouthpiece with the nostrils clamped. Samples were analysed by MedGraphics CPX (Medical Graphics Corp., St. Paul., Minnesota, US). Oxygen, carbon dioxide and flow sensors were calibrated immediately prior to each test.

A 12-lead electrocardiogram (ECG) was recorded continuously during the test. Systolic and diastolic blood pressure were recorded at baseline and systolic blood pressure was measured at 3 minute intervals during the test, followed by an assessment of dyspnoea, chest pain (Borg CR-10 scale)<sup>33, 34</sup> and Borg's rate of perceived exertion (RPE).<sup>33-35</sup>

The test was performed aiming to reach maximal exertion, "levelling-off" of the oxygen uptake and respiratory exchange ratios clearly above 1. The ECG recording was continued for 5 minutes after termination of exercise.

#### Statistical methods

Continuous data are presented as mean  $\pm$  standard deviation (SD), ordinal data as median and range. Significance was set as p < 0.05 throughout these studies. Computer analyses were carried out by STATISTICA<sup>TM</sup> 6.0 - 8.0, StatSoft®, Tulsa, Okla., USA.

Kaplan-Meier product-limit analysis was used to assess cumulative long-term survival. Risk analysis was performed by Cox regression model in Paper I. Comparison of pre- and postoperative data was performed by Wilcoxon's matched pairs test<sup>36</sup> and by sign test in Papers II & III respectively.

General discriminant analysis<sup>37</sup> (forward stepwise) model and general linear models, were used to verify connections among pre- and postoperative variables for data in Papers II & III.

The multivariate method of tree clustering<sup>37</sup> by Ward's method was applied to uncover relationships among data. Cluster analysis is an exploratory technique which organizes observed data into meaningful categories. Ward's method is based on analysis of variance as the linkage rule. The degree of association between variability of two parameters is maximal if they belong to the same group and minimal otherwise.

Multiple regression (forward stepwise regression) was performed to further specify relationships of significance among variables within a cluster formerly defined by tree clustering. Finally, these variables were plotted against each other in Paper IV.

Reproducibility and repeatability were assessed by correlation coefficient and Bland-Altman analysis<sup>38, 39</sup> in Paper III, and by gage process analysis<sup>40</sup> in a subgroup of 11 randomly chosen patients in Paper IV.

#### Ethical considerations


The presented studies were performed according to the principles of Human Rights in Declaration of Helsinki and were approved by the Ethics Committee of the Faculty of Health Sciences, University of Linköping. All the patients in the prospective studies (Papers II- IV) gave their informed consent.

# RESULTS

# Survival analysis (I)

In the group of the retrospectively studied patients 30-day mortality was 1%, due to acute heart failure (preoperative NYHA I and reduced LVF with corresponding EF 30-40%). Late mortality was 11%. Autopsy was performed in the case of early death and 2 of the cases of late death. Neither autopsy reports, nor clinical assessment indicated malfunction of the prostheses. The cause of death was cardiac in 5 cases: 2 cases of heart failure and 3 cases of myocardial infarction. Patients who died from heart failure were in NYHA classes III-IV preoperatively and those who suffered a late, fatal myocardial infarction were all in NYHA class III. Estimated preoperative EF of these five patients was > 50% in one case, 40-50% in two cases, and 30-40% in one case and < 30% in one case. Non-cardiac causes of death were insulin coma in 1, prostatic cancer in 1, pleuracarcinosis in 1 and cerebral bleeding in 2 patients, both of whom had mechanical prostheses and were treated with coumarine.

Α








Figure 5
Kaplan-Meier analysis. Panel A: Cardiac event free survival Panel B: Overall survival.

Survival analysis is shown in Figure 5. Gender, age and LV function could not be shown to be independent risk factors for impaired survival. Thrombosis, dysfunction of the prostheses, embolization or endocarditis were not reported postoperatively.

# Left ventricle (I-III)

Left ventricular diameters and volumes

In the retrospective study group, the mean aortic diameter was  $37.5 \pm 8$  mm at the level of the cusps. The end diastolic diameter of the LV was  $70 \pm 10$  mm while the end systolic diameter of the LV was  $49 \pm 10$  mm.

The prospectively studied patents also had a pronounced LV dilatation (LVED =  $68 \pm 6$  mm, LVES =  $49 \pm 7$  mm) with wall thickness within the upper normal range, which resulted in a high LV mass preoperatively. LV diameters diminished markedly, generally falling within normal limits postoperatively

and the overall comparison of pre- and postoperative LV masses and LV end-diastolic volumes showed significant reduction after surgery (Table 2, Paper II).

Preoperative NYHA class had no relation to pre- or postoperative left ventricular end-diastolic or end-systolic diameter (LVED, LVES) or LV mass. However, NYHA and the preoperative left ventricular volume at rest were significantly connected (p=0.02) but no other conclusions could be made concerning volumes.

#### Left ventricular function at rest & during exercise


Left ventricular function was normal, or only slightly impaired, in the majority of the retrospectively studied patients (Fig. 6A). However, 31 patients had moderate or severe left ventricular dysfunction preoperatively, i.e. estimated EF was ≤ 40%. Figures 6 B & C present the preoperative EF measured by echocardiography and MUGA in the prospective studies. In the same material, preoperatively calculated EF at rest was not a predictor for postoperative EF at rest.

By MUGA we found that both pre- and postoperative EF at maximal workload (EF<sub>max</sub>) could be predicted from preoperative EF<sub>rest</sub> (p < 0.001 and p=0.007 respectively).

Preoperative NYHA class was independent of both pre- and postoperative  $EF_{rest}$  and  $\Delta EF$ .


A

LVF - ECHO & NYHA for patients 1989-1999



В

#### LVF - ECHO & NYHA for patients 2002-2006



C

#### LVF - MUGA & NYHA for patients 2002-2006




Figure 6

Proportional distribution of NYHA functional class (I-IV) in the different preoperative EF categories at rest for both retrospectively and prospectively studied patients. Panel A: Patients operated on between 1989 and 1999. Panel B: Patients operated on between 2002 and 2006. Panel C: Patients operated on between 2002 and 2006. EF calculated by MUGA.

The semiquantitative description of LVF corresponds to the following EF values: Normal  $\geq$  50%, slight dysfunction 40-49%, moderate dysfunction 30-39%, severe dysfunction < 30%.

#### Left ventricle and aortic valve surgery

Preoperative LVED proved to be a predictor for both preoperative and postoperative  $\Delta$ EF (p=0.003 and 0.04 respectively) in general but did not have any predictive value in allocating the patients in the different subgroups based on EF response postoperatively. LVES, however, was independent of any exercise response. Preoperative left ventricular volumes could not be connected to EF at rest or during exercise.

Analyses of subgroups, formed according to  $\Delta EF$ , showed that patients with decreasing and unaltered EF preoperatively presented a significantly improved but still abnormal  $\Delta EF$  during exercise postoperatively. The group of patients with increasing EF preoperatively responded similarly to exercise postoperatively (Fig.3 & Table 2, Paper II). Preoperative  $\Delta EF$  proved to be a predictor for postoperative  $\Delta EF$  (p=0.02). Moreover, preoperative  $\Delta EF$  was the single and only predictor as to which subgroup (EF decreasing: p=0.03, unaltered: p=0.02, increasing: p=0.0008) patients were to be allocated postoperatively.

# Physical work capacity (III)

Patients performed the exercise test to maximal exhaustion both pre- and postoperatively. CPET data are shown in Table 3, Paper III. Patient's physical work capacity was classified according to Åstrand's reference values<sup>41</sup>, which are corrected for gender, age and weight. According to this classification, the pre- and postoperative work capacity was "low" in 17 cases and 16 cases, "fair" in 4 and 6 cases and "average" in 5 and 3 cases, for pre and post respectively. Figure 1, Paper III illustrates the different groups based on preoperative work capacity and their performance pre- and postoperatively.

Preoperative physical work capacity class according to Åstrand et al.<sup>41</sup> was a predictor of postoperative maximal oxygen uptake ( $\dot{V}$  O<sub>2 peak</sub>) per kg body weight (p=0.03): having a low physical work capacity according to Åstrand's classification implied a significant probability of remaining in the low performance class after surgery.

# Echocardiographic anatomy of the normal aortic valve (IV)

All aortic diameters changed by less than 1 mm between systole (s) and diastole (d). The subaortic diameter (SAD) showed no statistically significant connection to maximal sinus diameter, or to the STJ. However, the STJ could be described as a function of the mean maximal sinus diameter (maxSiD):

$$STJ = 3.7 + 0.76 \times \frac{maxSiD s + maxSiD d}{2}.$$

No connection was found between STJ and cusp or sinus length in the long-axis view.

Coaptation was observed in all patients, with length found to be independent of the aortic diameters but related to left coronary (LC) and right coronary (RC) cusp height in diastole;

coaptation = 
$$1.3 + 0.45 \times \frac{LC \text{ cusp height d} + RC \text{ cusp height d}}{2}$$
.

Neither LC sinus height nor RC sinus height in diastole alone showed any connection with coaptation. Their mean value, however, had a certain relation to coaptation, which was significant. The opening of the valve (OV) as a function of the subaortic diameter can be described as

$$OV = 1 + 0.84 \times SAD$$
.

The OV was also significantly related to the mean of the LC and RC cusp height in systole.

The intercommissural distances (ICD) in the right coronary (RC) and non-coronary (NC) sinuses measured in systole were equal and larger than that of the left coronary (LC) sinus. In diastole, the ICD in the RC sinus proved to be the largest, followed by LC sinus and NC sinus, when studied in the short-axis view (Table I, Paper IV). The ICD in the NC sinus changed more than that in either the LC or RC sinuses. The latter two changed equally between systole and diastole. These measurements showed the RC sinus to be deeper than the LC sinus, and the RC cusp to be longer than the LC cusp in the long-axis view. Sinus height was shortened by 2 mm in systole (Table I, Paper IV). The LC and RC cusp diameters were similar, and larger than the NC cusp diameter in

systole. However, this pattern changed in diastole, with LC and NC cusp diameters showing similarity and the RC cusp diameter being smallest (Table I & Fig. 2, Paper IV). Changes in cusp diameter and ICD were largest in the NC sinus and cusp, whereas changes in the LC and RC sinuses and cusps were smaller and of similar magnitude.

## **DISCUSSION**

Aspects of pathophysiology, prevalence & gender

When defining inclusion criteria for these studies, we aimed to form a population with chronic severe isolated AR requiring surgery, without coexistent cardiac disease.

The decision about surgery was then based on either progressive LV dilatation, indication of a declining LV function despite a normal EF, or the appearance of symptoms, together with an awareness not to delay surgery too long.

One fourth of the patients could be considered strictly asymptomatic, those who were NYHA I without medication. The rest of the patients were symptom free while on medication, or had mild to moderate symptoms.

That a few patients were in a higher NYHA class upon entry into the study was, in several of the cases, due to their AR not having been detected until then.

The patients were mainly middle-aged men, and this is influenced by the exclusion of patients with aneurysms, since aneurysm in connection with AR has been shown to be related to female gender.<sup>13, 42-44</sup> Chronic aortic insufficiency occurs predominantly in men between the ages of 20 and 80. The incidence and the prevalence of chronic aortic insufficiency<sup>13, 42</sup> explain the lack of women among patients in the prospective studies.

Previously published studies<sup>22-24</sup> did not, in contrast to our studies, define acute endocarditis as an exclusion criterion. Since acute surgery due to endocarditis was, however, extremely rare during the time period covered by these previous studies, our conclusions concerning comparison of survival data can still be assumed to be valid for the retrospectively studied patient population.

Survival after surgery for chronic AR

Although we have not found statistical comparisons to be justified, due to differences between populations for our retrospective data and results of earlier studies, our data from the last decade indicate a higher survival rate than that in the earlier Scandinavian samples. One Scandinavian study<sup>22</sup> covered the period 1969 to 1983 and showed a relative survival rate of 67% at 10 years in patients with AR despite a similar mean age at surgery to that of the patient population in our study. Other international studies<sup>42-46</sup> that covered the approximately 20-year period prior to our study have shown that the expected cardiac survival, as well as the overall survival for patients undergoing surgery for chronic aortic insufficiency was lower than that experienced with our patients. Turina et al.<sup>46</sup> published a study on 192 patients with chronic AR operated on between 1970 and 1983. Perioperative mortality was 2.6% and the survival rate after 10 years was 76%. They found that advanced age at the time of surgery and higher preoperative NYHA class were the main predictors of late mortality. However, Morris et al.<sup>45</sup> studied a group of patients who had both AR and aortic stenosis, and only a smaller group of their patients had pure AR.

During follow-up, one fifth of our patients had needed in-patient hospital care because of significant heart problems and a third had symptoms that could be of cardiac origin. The trend in the treatment of aortic regurgitation today is to recommend surgery before signs of obvious LV dysfunction develop; for example, asymptomatic patients often undergo surgery. Looking back one decade, this strategy had not completely penetrated, since the majority of our patients were moderately symptomatic, and substantial preoperative LV dysfunction was present in a third of the cases preoperatively in the retrospectively studied patients.

None of the prospectively studied patients died or suffered perioperative myocardial infarction.

#### Compensated & decompensated left ventricle in chronic AR

A leaking aortic valve causes a certain fraction of blood ejected at each systole to regurgitate back into the LV during diastole, causing a "pendulum volume". The total stroke volume increases, partly due to the regurgitant volume and partly because the LV increases its volume so as to provide an unaltered effective stroke volume. This leads to volume overload and consequently also to a pressure overload and, in order to compensate, the LV develops eccentric hypertrophy. Thus, an increased myocardial mass is inevitable for this compensation.

While the LV is able to compensate for the regurgitant lesion, LV diameters and mass increase are appropriate for meeting the increased needs

and EF remains within normal range. When reserves begin to languish, distinct alterations arise. A further dilatation of the LV and a drop in EF can be observed and, eventually, the patient becomes symptomatic. Current guidelines<sup>16, 17</sup> recommend operation based on LV diameters and EF to prevent further deterioration and preserve LVF (Table 2), and to avoid premature death.

There is general consensus<sup>16, 17, 47-51</sup> that patients with significant AR should be operated upon when signs of LV dysfunction appear, irrespective of symptoms. EF<sub>rest</sub> is considered to be an insensitive marker<sup>52-55</sup> and LV function during exercise has proved to be related to prognosis in AR patients. Consequently, even if the time point for surgery of a patient with chronic AR has been theoretically well defined and described<sup>16, 17</sup>, this theory often proves to be difficult to apply in clinical practice in asymptomatic or mildly symptomatic patients with normal EF<sub>rest</sub>.

Our prospectively studied patient population had an overall normal ejection fraction (Fig. 1, Paper II). We did not find any connection or correlation between NYHA functional class and EF, which is concordant with results from earlier studies. This confirms the earlier observations<sup>55, 56</sup> that lack of symptoms cannot be relied upon for judgement of LV function and timing of surgery in AR.

In order to detect imminent LV dysfunction in asymptomatic or mildly symptomatic patients, exercise testing is recommended in addition to follow-up of LV diameters and ejection fraction at rest for further differentiation. The predictive value is, however, considered to be unclear due to lack of sufficient evidence<sup>16, 17</sup>.

Previous studies of MUGA have shown that EF during exercise and exercise EF response have predictive value for the predicted length of the period until symptoms develop and there is an absolute indication for surgery.<sup>57, 58</sup> These studies were performed in an era when asymptomatic patients were not operated on. Furthermore, these variables had predictive value for the risk of sudden cardiac death in the natural history of the disease.<sup>59</sup> In the present study, preoperative EF<sub>rest</sub> was found to be significantly correlated to EF<sub>max</sub>, separately both pre- and postoperatively, which was in agreement with previous studies.<sup>30, 51, 52, 60</sup> However, preoperative EF<sub>rest</sub> was not related to postoperative EF increase upon exercise and was, therefore, not found to be informative for LV performance during exercise postoperatively. Moreover, we found that preoperative exercise response was not only a predictor for postoperative exercise response but that preoperative ΔEF predicted which type of exercise response the patient was going to have

postoperatively. This is an extension of results in accordance with earlier studies of cardiac catheterization<sup>57</sup> and echocardiography at rest and post-stress.<sup>58</sup>

Patients in the different subgroups defined by preoperative EF exercise response showed a similar exercise response pattern postoperatively. Decreasing EF during exercise preoperatively meant an improved but still not normally increasing EF during exercise on follow-up. The general pattern was that the group of patients with unaltered EF preoperatively responded to exercise normally after operation while those who had had normal  $\Delta EF$  had preserved it. This might indicate that at least those with decreasing EF should have been operated on earlier in order to preserve LVF, which had obviously deteriorated at the time of the operation in spite of normal EFrest. The importance of functional evaluation during exercise is emphasized by the left ventricular diameters lying within intervals, where guidelines recommend regular follow-up and consideration of hemodynamic response before a decision about surgery is recommended. LV volumes significantly diminished postoperatively in the whole study group, as expected and described in previous studies.61, 62 We found that EF calculations by MUGA and by ultrasound were in agreement, although echocardiography gave a lower level of EF in our study, as has been reported by others.<sup>63</sup> This indicates that careful choice and use of validated methods is important for follow-up over the time. Determination of EF by radionuclide ventriculography reproducibility, even during exercise<sup>64</sup> and therefore adds valuable information in borderline cases.

### Effects of AR on physical work capacity

During exercise, the fraction of the stroke volume that regurgitates into the LV, i.e. the regurgitant fraction, diminishes, due to the heart rate increase with a relatively more pronounced shortening of diastole than of systole. This reduction is reported to be approximately 20%.<sup>65</sup> Due to this reduced regurgitation, patients with significant AR show a reduction of both diastolic and systolic LV volumes during exercise.<sup>65, 66</sup> Thereby, if their LV function is normal, the EF of persons with AR increases upon exercise, even if total stroke volume does not. This hemodynamic reaction to exercise in AR provides an extremely adequate explanation as to, why the exercise capacity may remain normal for a long time despite severe AR. When symptoms occur, initial complaints are usually tiredness or dyspnoea upon exertion, indicating a

reduced capacity to increase effective cardiac output normally and/or elevated LV filling pressures.

During the last 1-2 decades, CPET has been extensively studied<sup>67-69</sup> and reported to be a valuable tool in the evaluation of patients with heart failure. A few studies have also pointed towards the possibility of using CPET to evaluate the results of cardiac rehabilitation after cardiac events or cardiac surgery.<sup>70-73</sup> These studies have shown that CPET is a reliable method and that VO2 peak is a useful predictor of heart failure patients' event free survival, as well as a good measure of the positive effect of cardiac rehabilitation by physical training.

In our patients with severe AR, there was a striking discrepancy between preoperative NYHA class, EF and physical work capacity. The patients as a group (median NYHA II) presented a surprisingly poor physical work capacity preoperatively and failed to show any improvement 6 months postoperatively and patients in NYHA III did not consistently have the lowest physical work capacity.

The expected improvement, as seen in previous studies in patients who entered a cardiac rehabilitation program <sup>71-73</sup>, did not occur in this specific group. This could be explained by the fact that our patients did not participate in a cardiac rehabilitation program or, simply, that the follow-up interval needs to be longer for patients with chronic AR<sup>74</sup>, assuming a longer recovery period than for patients with coronary disease or other valve disease.<sup>70-73</sup> Further studies are indicated as to whether having a prosthetic aortic valve may also per se impede the physical work capacity.

The lack of postoperative improvement in our patients could also be due to subclinical LV dysfunction that is not possible to identify by preoperative examinations at rest. However, the cardiac compensation by LV dilatation and eccentric hypertrophy, as well as systemic adaptations, may be insufficient and thereby manifested during CPET. Preoperative physical work capacity by Åstrand's classification, a  $\dot{V}$  O2 peak nomogram corrected for age, gender and weight, was a predictor for postoperative physical work capacity. As  $\dot{V}$  O2 peak is an indicator of aerobic work and thereby for the general status of the oxygen transport and utilisation capacity, it may represent potential for detecting latent heart muscle dysfunction at an earlier stage. This would implement a complementary examination to the routine follow-up of the patients with chronic AR who have normal EF and moderately elevated LV diameters and volumes.

### Timing of surgery

The treatment of severe chronic aortic regurgitation is surgical, and in the vast majority of cases, means implantation of a prosthetic valve. Although this can be done with low perioperative mortality, the postoperative morbidity is not negligible, e.g. postoperative heart failure, patient prosthesis mismatch<sup>75-78</sup> and bleeding complications due to coumarine treatment.

Timing of surgery for patients with chronic aortic regurgitation remains an art<sup>18, 19, 79, 80</sup> because our standard methods merely estimate LVF and do not provide exact measurement of myocardial contractility. We of course, have to keep trying to give the best possible estimation.

Our retrospective analysis showed that overall survival for patients operated for chronic AR during a defined interval was improved compared to previously published data on survival for chronic AR. One third of these patients had moderate or severe LV dysfunction at the time of the operation and an irreversible LV dysfunction may have affected their postoperative survival and morbidity.

The LVF, based on resting EF alone, for the prospective study group was better compared to the retrospective group (Fig. 6) but a majority of these patients presented abnormal EF reaction and low physical capacity both prior to and after surgery. These reactions may be used to provoke latent LV dysfunction and identify patients on the borderline of developing LV dysfunction and thereby indicate an operation earlier in the natural history of the disease.

As a treating physician, it is a difficult decision to recommend surgery to an asymptomatic person, who will not feel any improvement but rather, in the short term, may suffer complications. The decision to recommend surgery must then rest on the rationale that surgery is highly probable to be beneficial to the patient in the long term, to avoid irreversible LV dysfunction.

### Clinical anatomy of the aortic valve

Although supporting data are scarce as to whether patients with chronic AR should have better preserved LVF or have a more favourable outcome of rehabilitation after aortic valve sparing operation<sup>81-84</sup> compared to implantation of a prosthetic valve, it is conceivable that avoiding patient prosthesis mismatch per se would provide better function. Reconstructing a

failing valve requires knowledge about normally functioning valves. Several animal<sup>85-92</sup> and human<sup>93-97</sup> studies describing normal anatomy and function of the aortic valve and root have been performed. The former in vivo but the latter mostly consist of autopsy data. The aortic flow characteristics and opening patterns of the valve have been studied in human series of small size <sup>9, 85, 98-101</sup>

We have applied an established diagnostic method (TEE) and extended its use, in order to provide a more detailed description<sup>102-105</sup> of the in vivo physioanatomy of the normal aortic root in its entirety in contrast to previous studies.<sup>106-110</sup> Measurements of the aortic root showed overall larger dimensions compared to those found in cadaver studies and reported previously, when the same parameter was measured.<sup>93, 96, 97</sup> Postmortem studies are based on measuring inner size, as in the present study. However, the effect of blood pressure as a factor that can modify dimensions in vivo should also be taken into consideration. A difficulty emerges even when comparing our measurements to earlier echocardiographic studies, due to differences in measuring by inner edge to inner edge, trailing edge to leading edge, or leading edge to leading edge.

There is no ruling consensus about whether the sinuses are alike or if they are different in size.<sup>94, 96</sup> The statement, based on an autopsy material, by Berdajs et al.<sup>96</sup> of a pattern 'R > N > L', meaning that the right coronary sinus is the largest and the left coronary sinus the smallest, was not confirmed by the present data. Indeed, analysis of these data showed the NC sinus and cusp not to resemble the other two, although these relations were shown to depend on the cardiac cycle. Changes in the dimensions of both cusp diameter and intercommissural distance were largest in the NC area and less, but still alike, for the RC and LC parameters. The interpretation here is that the NC sinus and cusp have a more extensive movement than the other two, at least when studied using 2D TEE.

It was also found, somewhat surprisingly, that the subaortic diameter was not related to either the maximal sinus diameter or to the STJ. This may mean that this parameter is influenced by the structures and physiological events in the left ventricle, and that its changes have no impact on maximal width in the sinuses of Valsalva and the STJ. This observation is concordant with the observations of Sutton et al.<sup>111</sup> and Lansac et al.<sup>112</sup> If the subaortic diameter is related to the left ventricular outlet, then a statistically significant relation to the opening of the valve, as found in the present study, would be reasonable. However, as the constant in this equation is less than the standard deviation of the subaortic diameter, it can be stated that a normal valve opens to

approximately 80% of the dimension of the subaortic diameter in the transesophageal long-axis view. When choosing a graft for reconstructive surgery, this means that a graft with a diameter the same as the subaortic diameter would allow valve opening without contacting the graft. Hence, an accelerated leaflet degeneration and thickening caused by contact with the graft might be avoided.

The relation between the STJ and the maximal sinus diameter might be relevant in the reconstruction of aortic root geometry. The regression equation may be simplified to state that, in healthy persons, the STJ is three-quarters of the maximal sinus diameter.

Another parameter that is usually considered important is coaptation; however, such importance is based on common sense as its existence was rarely stated and its length was not measured in earlier studies. The finding that coaptation occurred in all of the patients and that all of the valves were competent, emphasizes its importance. The analysis showed a relation only between the cusps in diastole and the coaptation. The equation describing coaptation considering the standard deviation for coaptation and that the precision of the echocardiographic measurements is 1-2 mm, can be interpreted as, without oversimplifying or neglecting important details, that the length of coaptation was approximately half of the cusp height. It is noteworthy that, although there is frequently a coexistence of root dilatation and an absence of coaptation in AR patients, the length of coaptation in normal subjects was found to be independent of the aortic root diameters.

## **CONCLUSIONS**

### I. Retrospective analysis of left ventricular function

i) A reduced LV function preoperatively could be seen in a substantial number of patients operated on for chronic AR during the period 1989-1999.

### II. LV function at rest and during exercise

- i) Preoperative exercise response measured with radionuclide ventriculography can identify patients at risk of an abnormal postoperative LV function response during exercise.
- ii) It can predict what kind of exercise response (decreasing, unaltered or increasing EF during exercise) to expect after surgery.
- iii) An abnormal EF response to exercise might also occur in patients who do not fulfil the criteria for surgery based on LV dimensions by echocardiography or EF at rest.

#### III. Physical work capacity in patients with chronic AR

i) Echocardiographic parameters such as EF, LV diameters and volumes at rest did not reflect the preoperative physical work capacity in patients with chronic AR and neither did they predict the postoperative physical work capacity.

#### IV. Echocardiographic anatomy of the normal aortic valve

- i) The length of coaptation in normal subjects was found to be independent of the aortic root diameters and was approximately half of the cusp height.
- ii) The subaortic diameter was not related to either the maximal sinus diameter or to the STJ.
- iii) A normal valve opens to approximately 80% of the dimension of the subaortic diameter in the transesophageal long-axis view.

## **CLINICAL IMPLICATIONS**

- I. Efforts should be made to evaluate LV function and detect LV dysfunction, preferably by a non-invasive method, early in the natural history of patients with AR. The rationale for performing surgery earlier than previously will most probably be justified.
- II. Patients with significant chronic AR could benefit from follow-up of LV function during exercise and by considering surgery when the character of the exercise response is abnormal or changes character.
- III. We suggest that the use of CPET in preoperative follow up and decisionmaking as a complement to echocardiography may be a useful tool in identifying patients with poor physical work capacity indicating insufficient compensation and latent LV dysfunction in chronic AR.
- IV. Detailed mapping of the aortic root by echocardiography might help to improve techniques for restoring normal functional anatomic relations in the pathologically altered aortic root.

# **ACKNOWLEDGEMENTS**

Christian Olin, my former tutor, taught me a lot about cardiac surgery. He encouraged and helped me to start this work. However, Árpád Péterffy had made it possible for me to start to work with cardiothoracic surgery at all. Therefore, I am deeply indebted for these gentlemen for the rest of my life.

When expressing gratitude is in order I feel that I have to thank everyone who contributed to this work by attending to patients, conducting examinations, taking good care of administrative work and logistics and scrutinizing the manuscript - all the help and encouragements are dearly remembered.

Eva Nylander was my tutor while the majority of the work on this thesis was done. I'm afraid I have no words to properly tell how much fun it is to discover new things with her help, how much I appreciate her patience when I'm slow in understanding and how much I enjoy the freedom she has given me in my work yet guiding me always wisely and steadily. It is a privilege to work with her.

Compliments should be paid to Mats Broqvist, co-tutor and my co-author as well as to Eva Olsson, Stefan Franzén, Niels-Erik Nielsen and Ingemar Vanhanen, my other co-authors, who contributed a lot through taking part in designing and realizing these studies and creatively criticising the results and the manuscripts during enjoyable discussions.

... and Ilona and Gábor, my amiable parents, who already know very well how grateful I am - for everything.

Tamás Éva May 2008 The studies were performed at the Dept of Cardiothoracic and Vascular Surgery and at the Dept of Clinical Physiology and Nuclear Medicine, Heart Centre in Östergötland, University Hospital Linköping.

The work was financially supported by:

The Swedish Heart - Lung Foundation
The County Council of Östergötland
The Research Foundations of the Heart Centre in Östergötland
Lions Foundation for Research at Linköping University

## REFERENCES

- 1. Harvey W. Exercitatio Anatomica de Motu Cordis et Sanguinis in Animalibus (Anatomical Motion of the Heart and Blood); 1628.
- 2. Mercer JL, Benedicty M, Bahnson HT. The geometry and construction of the aortic leaflet. J Thorac Cardiovasc Surg 1973;65:511-8.
- 3. Swanson M, Clark RE. Dimensions and geometric relationships of the human aortic valve as a function of pressure. Circ Res 1974;35:871-82.
- 4. Andersson R. Clinical anatomy of the aortic root. Heart 2000;84:670-3.
- 5. El Khoury G, Glineur D, Rubay J, et al. Functional classification of aortic root/valve abnormalities and their correlation with etiologies and surgical procedures. Curr Opin Cardiol 2005;20:115-21.
- 6. Zipes DP, Libby P, Bonow RO, Braunwald E. Braunwald's Heart Disease. 7th ed; 2005.
- 7. Kirklin JW, Barratt-Boyes BG. Cardiac Surgery. 2nd ed: Churchill Livingstone; 1993.
- 8. Piwnica A, Westaby S. Surgery for required Aortic Valve Disease. 1st ed: Isis Medical Media Ltd.; 1997.
- Kvitting JP, Ebbers T, Wigström L, Engvall J, Olin CL, Bolger AF. Flow patterns in the aortic root and the aorta studied with time-resolved, 3-dimensional, phase-contrast magnetic resonance imaging: implications for aortic valve-sparing surgery. J Thorac Cardiovasc Surg 2004;127:1602-7.
- 10. Jonson B, Westling H, White T, Wollmer P. Klinisk Fysiologi. 1st ed: Liber; 1998.
- 11. Walsh CA, Wilde P. Practical echocardiography. 1st ed: Oxford University Press; 1999.
- 12. Hammarström E. Doppler ekokardiografi: Bok & Tryck AB; 1996.
- 13. Singh JP, Evans JC, Levy D, et al. Prevalence and clinical determinants of mitral, tricuspid, and aortic regurgitation (the Framingham Heart Study). Am J Cardiol 1999;83:897-902.
- 14. Bonow RO. Aortic Regurgitation. Curr Treat Options Cardiovasc Med 2000;2:125-32.
- 15. Maurer G. Aortic regurgitation. Heart 2006;92:994-1000.
- 16. Bonow RO, Carabello BA, Kanu C, et al. ACC/AHA 2006 guidelines for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (writing committee to revise the 1998 Guidelines for the Management of Patients With Valvular Heart Disease): developed in collaboration with the Society of Cardiovascular Anesthesiologists: endorsed by the Society for Cardiovascular Angiography and Interventions and the Society of Thoracic Surgeons. Circulation 2006;114:e84-231.
- 17. Vahanian A, Baumgartner H, Bax J, et al. Guidelines on the management of valvular heart disease: The Task Force on the Management of Valvular Heart Disease of the European Society of Cardiology. Eur Heart J 2007;28:230-68.
- 18. Bonow RO. Chronic aortic regurgitation. Role of medical therapy and optimal timing for surgery. Cardiol Clin 1998;16:449-61.
- Hicks GL, Jr., Massey HT. Update on indications for surgery in aortic insufficiency. Curr Opin Cardiol 2002;17:172-8.

- 20. Katz AS, Devereux RB. Timing of surgery in chronic aortic regurgitation. Echocardiography 2000;17:303-11.
- 21. Tarasoutchi F, Grinberg M, Filho JP, et al. Symptoms, left ventricular function, and timing of valve replacement surgery in patients with aortic regurgitation. Am Heart J 1999;138:477-85.
- 22. Lindblom D, Lindblom U, Qvist J, Lundström H. Long-term relative survival rates after heart valve replacement. J Am Coll Cardiol 1990;15:566-73.
- 23. Abdelnoor M, Hall KV, Nitter-Hauge S, et al. Risk factors of morbidity and mortality in surgically treated chronic aortic valvular heart disease. Rev Epidemiol Sante Publique 1988;36:89-98.
- 24. Abdelnoor M, Nitter-Hauge S, Trettli S. Relative survival of patients after heart valve replacement. Eur Heart J 1990;11:23-8.
- Bonow RO, Rosing DR, McIntosh CL, et al. The natural history of asymptomatic patients with aortic regurgitation and normal left ventricular function. Circulation 1983;68:509-17.
- 26. Alehagen U, Eriksson H, Nylander E, Dahlström U. Heart failure in the Elderly: Characteristics of a Swedish Primary Health Care Population. Heart Drug 2002;2:211-20.
- 27. Wallentin Guron C, Bech-Hanssen O, Wikh R, Rosengren A, Hartford M, Caidahl K. The E/e filling index and right ventricular pressure in relation to applied international Doppler recommendations of left ventricular filling assessment. Eur J Echocardiogr 2005;6:419-28.
- 28. Cheitlin MD, Armstrong WF, Aurigemma GP, et al. ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography--summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). J Am Coll Cardiol 2003;42:954-70.
- 29. Wallis JW, Wu-Connolly L, Rocchini AP, Juni JE. Dynamic arrhythmia filtration for gated blood-pool imaging: validation against list mode technique. J Nucl Med 1986;27:1347-52.
- 30. Bassand JP, Faivre R, Berthout P, et al. Factors influencing the variations of ejection fraction during exercise in chronic aortic regurgitation. Eur J Nucl Med 1987;13:419-24.
- 31. Anand IS, Florea VG, Solomon SD, Konstam MA, Udelson JE. Noninvasive assessment of left ventricular remodeling: concepts, techniques, and implications for clinical trials. J Card Fail 2002;8:S452-64.
- 32. Iskandrian AS, Heo J. Radionuclide angiographic evaluation of left ventricular performance at rest and during exercise in patients with a regurgitation. Am Heart J 1986;111:1143-9.
- 33. Borg G, Holmgren A, Lindblad I. Quantitative evaluation of chest pain. Acta Med Scand Suppl 1981;644:43-5.
- 34. Borg G. Psychophysical scaling with applications in physical work and the perception of exertion. Scand J Work Environ Health 1990;16 Suppl 1:55-8.
- 35. Borg GA. Psychophysical bases of perceived exertion. Med Sci Sports Exerc 1982;14:377-81
- 36. Altman DG. Practical statistics for medical research. 1st ed: Chapman & Hall; 1991.
- 37. Manly B. Multivariate Statistical Methods. A primer. 2nd ed: Chapman & Hall 2000.
- 38. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. Lancet 1986;1:307-10.

- 39. Bland JM, Altman DG. Comparing methods of measurement: why plotting difference against standard method is misleading. Lancet 1995;346:1085-7.
- 40. Duncan A. Random-effects models and mixed models In: Quality Control and Industrial Statistics. 4th ed: RD Irwin, Homewood, IL; 1974:716-34.
- 41. Åstrand I. Aerobic work capacity in men and women with special reference to age. Acta Physiol Scand Suppl 1960;49:1-92.
- 42. Carabello BA. Aortic regurgitation in women. Does the measuring stick need a change? Circulation 1996;94:2355-7.
- 43. Klodas E, Enriquez-Sarano M, Tajik AJ, Mullany CJ, Bailey KR, Seward JB. Surgery for aortic regurgitation in women. Contrasting indications and outcomes compared with men. Circulation 1996;94:2472-8.
- 44. McDonald ML, Smedira NG, Blackstone EH, Grimm RA, Lytle BW, Cosgrove DM. Reduced survival in women after valve surgery for aortic regurgitation: effect of aortic enlargement and late aortic rupture. J Thorac Cardiovasc Surg 2000;119:1205-12.
- 45. Morris JJ, Schaff HV, Mullany CJ, et al. Determinants of survival and recovery of left ventricular function after aortic valve replacement. Ann Thorac Surg 1993;56:22-9; discussion 9-30.
- 46. Turina J, Milincic J, Seifert B, Turina M. Valve replacement in chronic aortic regurgitation. True predictors of survival after extended follow-up. Circulation 1998;98:II100-6; discussion II6-7.
- 47. Bonow RO. Asymptomatic aortic regurgitation: indications for operation. J Card Surg 1994;9:170-3.
- 48. Borer JS, Herrold EM, Hochreiter CA, et al. Aortic regurgitation: selection of asymptomatic patients for valve surgery. Adv Cardiol 2002;39:74-85.
- Borer JS, Supino PG, Hochreiter C, et al. Valve surgery in the asymptomatic patient with aortic regurgitation: current indications and the effect of change rates in objective measures. Adv Cardiol 2004;41:36-47.
- 50. Henry WL, Bonow RO, Borer JS, et al. Observations on the optimum time for operative intervention for aortic regurgitation. I. Evaluation of the results of aortic valve replacement in symptomatic patients. Circulation 1980;61:471-83.
- 51. Lindsay J, Jr., Silverman A, Van Voorhees LB, Nolan NG. Prognostic implications of left ventricular function during exercise in asymptomatic patients with aortic regurgitation. Angiology 1987;38:386-92.
- 52. Bonow RO, Borer JS, Rosing DR, et al. Preoperative exercise capacity in symptomatic patients with aortic regurgitation as a predictor of postoperative left ventricular function and long-term prognosis. Circulation 1980;62:1280-90.
- 53. Bonow RO, Lakatos E, Maron BJ, Epstein SE. Serial long-term assessment of the natural history of asymptomatic patients with chronic aortic regurgitation and normal left ventricular systolic function. Circulation 1991;84:1625-35.
- 54. Bonow RO, Picone AL, McIntosh CL, et al. Survival and functional results after valve replacement for aortic regurgitation from 1976 to 1983: impact of preoperative left ventricular function. Circulation 1985;72:1244-56.
- 55. Borer JS, Bacharach SL, Green MV, et al. Exercise-induced left ventricular dysfunction in symptomatic and asymptomatic patients with aortic regurgitation: assessment with radionuclide cineangiography. Am J Cardiol 1978;42:351-7.

- 56. Kraus F, Dacian S, Hall D, Klein U, Rudolph W. Relationship between symptoms and hemodynamics associated with regurgitant lesions of the aortic or mitral valve. Z Kardiol 1986;75 Suppl 2:137-40.
- 57. Wahi S, Haluska B, Pasquet A, Case C, Rimmerman CM, Marwick TH. Exercise echocardiography predicts development of left ventricular dysfunction in medically and surgically treated patients with asymptomatic severe aortic regurgitation. Heart 2000;84:606-14.
- 58. Tabayashi K, Kagawa Y, Abe Y, Yokoyama A, Onodera H, Horiuchi T. Long-term follow-up of patients who have undergone valve replacements for acquired valvular disease: a clinical and hemodynamic study. Tohoku J Exp Med 1982;138:209-22.
- 59. Borer JS, Hochreiter C, Herrold EM, et al. Prediction of indications for valve replacement among asymptomatic or minimally symptomatic patients with chronic aortic regurgitation and normal left ventricular performance. Circulation 1998;97:525-34.
- Kawanishi DT, McKay CR, Chandraratna PA, et al. Cardiovascular response to dynamic exercise in patients with chronic symptomatic mild-to-moderate and severe aortic regurgitation. Circulation 1986;73:62-72.
- 61. Bonow RO, Dodd JT, Maron BJ, et al. Long-term serial changes in left ventricular function and reversal of ventricular dilatation after valve replacement for chronic aortic regurgitation. Circulation 1988;78:1108-20.
- Jin XY, Pepper JR, Gibson DG, Yacoub MH. Early changes in the time course of myocardial contraction after correcting aortic regurgitation. Ann Thorac Surg 1999;67:139-45.
- 63. van Royen N, Jaffe CC, Krumholz HM, et al. Comparison and reproducibility of visual echocardiographic and quantitative radionuclide left ventricular ejection fractions. Am J Cardiol 1996;77:843-50.
- 64. Klocke FJ, Baird MG, Lorell BH, et al. ACC/AHA/ASNC guidelines for the clinical use of cardiac radionuclide imaging--executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASNC Committee to Revise the 1995 Guidelines for the Clinical Use of Cardiac Radionuclide Imaging). J Am Coll Cardiol 2003;42:1318-33.
- 65. Thompson R, Ross I, Leslie P, Easthope R. Haemodynamic adaptation to exercise in asymptomatic patients with severe aortic regurgitation. Cardiovasc Res 1985;19:212-8.
- 66. Iskandrian AS, Hakki AH, Amenta A, Mandler J, Kane S. Regulation of cardiac output during upright exercise in patients with aortic regurgitation. Cathet Cardiovasc Diagn 1984:10:573-82.
- 67. Ades PA, Savage PD, Brawner CA, et al. Aerobic capacity in patients entering cardiac rehabilitation. Circulation 2006;113:2706-12.
- 68. Ingle L. Theoretical rationale and practical recommendations for cardiopulmonary exercise testing in patients with chronic heart failure. Heart Fail Rev 2007;12:12-22.
- 69. Koike A, Koyama Y, Itoh H, Adachi H, Marumo F, Hiroe M. Prognostic significance of cardiopulmonary exercise testing for 10-year survival in patients with mild to moderate heart failure. Jpn Circ J 2000;64:915-20.
- 70. Kavanagh T. Exercise in cardiac rehabilitation. Br J Sports Med 2000;34:3.
- 71. Kavanagh T, Mertens DJ, Hamm LF, et al. Prediction of long-term prognosis in 12 169 men referred for cardiac rehabilitation. Circulation 2002;106:666-71.
- 72. Kavanagh T, Mertens DJ, Hamm LF, et al. Peak oxygen intake and cardiac mortality in women referred for cardiac rehabilitation. J Am Coll Cardiol 2003;42:2139-43.

- 73. Ueshima K, Kamata J, Kobayashi N, et al. Effects of exercise training after open heart surgery on quality of life and exercise tolerance in patients with mitral regurgitation or aortic regurgitation. Jpn Heart J 2004;45:789-97.
- Borer JS, Rosing DR, Kent KM, et al. Left ventricular function at rest and during exercise after aortic valve replacement in patients with aortic regurgitation. Am J Cardiol 1979;44:1297-305.
- 75. Blais C, Dumesnil JG, Baillot R, Simard S, Doyle D, Pibarot P. Impact of valve prosthesis-patient mismatch on short-term mortality after aortic valve replacement. Circulation 2003;108:983-8.
- 76. Pibarot P, Dumesnil JG. Prevention of valve prosthesis--patient mismatch before aortic valve replacement: does it matter and is it feasible? Heart 2007;93:549-51.
- 77. Pibarot P, Dumesnil JG. Hemodynamic and clinical impact of prosthesis-patient mismatch in the aortic valve position and its prevention. J Am Coll Cardiol 2000;36:1131-41.
- 78. Pibarot P, Dumesnil JG, Lemieux M, Cartier P, Metras J, Durand LG. Impact of prosthesis-patient mismatch on hemodynamic and symptomatic status, morbidity and mortality after aortic valve replacement with a bioprosthetic heart valve. J Heart Valve Dis 1998;7:211-8.
- 79. Katz AS DR. Timing of surgery in chronic aortic regurgitation. Echocardiography 2000;17:303-11.
- 80. Tarasoutchi F GM, Filho JP, Izaki M, Cardoso LF, Pomerantezeff P, Nuschbacher A, da Luz PL. Symptoms, left ventricular function, and timing of valve replacement surgery in patients with aortic regurgitation. Am Heart J 1999;Sep;138:477-85.
- 81. David TE, Armstrong S, Ivanov J, Feindel CM, Omran A, Webb G. Results of aortic valve-sparing operations. J Thorac Cardiovasc Surg 2001;122:39-46.
- 82. David TE, Feindel CM. An aortic valve-sparing operation for patients with aortic incompetence and aneurysm of the ascending aorta. J Thorac Cardiovasc Surg 1992;103:617-21; discussion 22.
- 83. Pepper J, Yacoub M. Valve conserving operation for aortic regurgitation. J Card Surg 1997;12:151-6.
- 84. Yacoub MH, Gehle P, Chandrasekaran V, Birks EJ, Child A, Radley-Smith R. Late results of a valve-preserving operation in patients with aneurysms of the ascending aorta and root. J Thorac Cardiovasc Surg 1998;115:1080-90.
- 85. Robicsek F, Thubrikar MJ. Role of sinus wall compliance in aortic leaflet function. Am J Cardiol 1999;84:944-6, A7.
- 86. Thubrikar M. The Aortic Valve. 1st ed: CrC Press, Inc.; 1990.
- 87. Thubrikar M, Bosher LP, Nolan SP. The mechanism of opening of the aortic valve. J Thorac Cardiovasc Surg 1979;77:863-70.
- 88. Thubrikar M, Harry R, Nolan SP. Normal aortic valve function in dogs. Am J Cardiol 1977;40:563-8.
- 89. Thubrikar M, Nolan SP, Bosher LP, Deck JD. The cyclic changes and structure of the base of the aortic valve. Am Heart J 1980;99:217-24.
- 90. Thubrikar M, Piepgrass WC, Shaner TW, Nolan SP. The design of the normal aortic valve. Am J Physiol 1981;241:H795-801.
- 91. Brewer RJ, Deck JD, Capati B, Nolan SP. The dynamic aortic root. Its role in aortic valve function. J Thorac Cardiovasc Surg 1976;72:413-7.

- 92. Brewer RJ, Mentzer RM, Jr., Deck JD, Ritter RC, Trefil JS, Nolan SP. An in vivo study of the dimensional changes of the aortic valve leaflets during the cardiac cycle. J Thorac Cardiovasc Surg 1977;74:645-50.
- 93. Westaby S, Karp RB, Blackstone EH, Bishop SP. Adult human valve dimensions and their surgical significance. Am J Cardiol 1984;53:552-6.
- 94. Silver MA, Roberts WC. Detailed anatomy of the normally functioning aortic valve in hearts of normal and increased weight. Am J Cardiol 1985;55:454-61.
- 95. Yacoub MH, Kilner PJ, Birks EJ, Misfeld M. The aortic outflow and root: a tale of dynamism and crosstalk. Ann Thorac Surg 1999;68:S37-43.
- 96. Berdajs D, Lajos P, Turina M. The anatomy of the aortic root. Cardiovasc Surg 2002;10:320-7.
- 97. Kunzelman KS, Grande KJ, David TE, Cochran RP, Verrier ED. Aortic root and valve relationships. Impact on surgical repair. J Thorac Cardiovasc Surg 1994;107:162-70.
- 98. Markl M, Draney MT, Miller DC, et al. Time-resolved three-dimensional magnetic resonance velocity mapping of aortic flow in healthy volunteers and patients after valve-sparing aortic root replacement. J Thorac Cardiovasc Surg 2005;130:456-63.
- 99. De Paulis R, De Matteis GM, Nardi P, Scaffa R, Bassano C, Chiariello L. Analysis of valve motion after the reimplantation type of valve-sparing procedure (David I) with a new aortic root conduit. Ann Thorac Surg 2002;74:53-7.
- 100. Leyh RG, Schmidtke C, Sievers HH, Yacoub MH. Opening and closing characteristics of the aortic valve after different types of valve-preserving surgery. Circulation 1999;100:2153-60.
- 101. Maselli D, Montalto A, Santise G, Minardi G, Manzara C, Musumeci F. A normogram to anticipate dimension of neo-sinuses of valsalva in valve-sparing aortic operations. Eur J Cardiothorac Surg 2005;27:831-5.
- 102. Roman MJ, Devereux RB, Kramer-Fox R, O'Loughlin J. Two-dimensional echocardiographic aortic root dimensions in normal children and adults. Am J Cardiol 1989;64:507-12.
- 103. Sheil ML, Jenkins O, Sholler GF. Echocardiographic assessment of aortic root dimensions in normal children based on measurement of a new ratio of aortic size independent of growth. Am J Cardiol 1995;75:711-5.
- 104. Cohen GI, White M, Sochowski RA, et al. Reference values for normal adult transesophageal echocardiographic measurements. J Am Soc Echocardiogr 1995;8:221-30.
- 105. Drexler M, Erbel R, Muller U, Wittlich N, Mohr-Kahaly S, Meyer J. Measurement of intracardiac dimensions and structures in normal young adult subjects by transesophageal echocardiography. Am J Cardiol 1990;65:1491-6.
- 106. Francis GS, Hagan AD, Oury J, O'Rourke RA. Accuracy of echocardiography for assessing aortic root diameter. Br Heart J 1975;37:376-8.
- 107. Vasan RS, Larson MG, Levy D. Determinants of echocardiographic aortic root size. The Framingham Heart Study. Circulation 1995;91:734-40.
- 108. Espinal M, Fuisz AR, Nanda NC, Aaluri SR, Mukhtar O, Sekar PC. Sensitivity and specificity of transesophageal echocardiography for determination of aortic valve morphology. Am Heart J 2000;139:1071-6.
- 109. Stoddard MF, Liddell NE, Vogel RL, Longaker RA, Dawkins PR. Comparison of cardiac dimensions by transesophageal and transthoracic echocardiography. Am Heart J 1992;124:675-8.

- 110. Schnittger I, Gordon EP, Fitzgerald PJ, Popp RL. Standardized intracardiac measurements of two-dimensional echocardiography. J Am Coll Cardiol 1983;2:934-8.
- 111. Sutton JP, 3rd, Ho SY, Anderson RH. The forgotten interleaflet triangles: a review of the surgical anatomy of the aortic valve. Ann Thorac Surg 1995;59:419-27.
- 112. Lansac E, Lim HS, Shomura Y, et al. A four-dimensional study of the aortic rootdynamics. Eur J Cardiothorac Surg 2002;22:497-503.