COMMENTARY

REDOX SIGNALLING BY TRANSCRIPTION FACTORS NF- κ B AND AP-1 IN LYMPHOCYTES

KLAUS SCHULZE-OSTHOFF,*† MAREK LOS‡ and PATRICK A. BAEUERLE*

*Institute of Biochemistry, Albert-Ludwigs-University, D-79104 Freiburg, Germany; and ‡Division of Molecular Oncology, German Cancer Research Center, D-69120 Heidelberg, Germany

Key words: NF- κ B; AP-1; transcription; reactive oxygen intermediates; redox regulation; lymphocytes

When molecular oxygen accumulated in our atmosphere more than 2 billion years ago, aerobic organisms adapted using oxygen as an electron acceptor in respiration. Although respiration is, by far, one of the most efficient ways of generating energy in biological systems, it has one harmful side-effect: incomplete reduction of dioxygen results in the formation of ROIs.[§] These include the superoxide anion ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), and the hydroxyl radical ($OH\cdot$). Superoxide anions originate from a single electron transfer on dioxygen [1]. This ROI has a short diffusion radius inside the cell, but it is eventually converted by SODs into H_2O_2 , the most stable and diffusible ROI. Transition metals, such as iron or copper ions, convert $O_2^{\cdot-}$ and H_2O_2 to an extremely reactive hydroxyl radical via the Fenton or Haber/Weiss reaction.

While low amounts of ROIs can be considered as physiological side-products of various electron transfer reactions and are tolerated by the cell, increased levels of ROIs, referred to as oxidative stress, are linked to pathological phenomena. Many adverse conditions, including ionizing and UV irradiation, heat shock, viral and bacterial infections, as well as environmental pollutants cause oxidative stress. In principle, ROIs can react with all classes of biological macromolecules, resulting in lipid peroxidation, inactivation of proteins, and strand breakage in nucleic acids [1].

During evolution, cells of the immune system, in particular, have learned to utilize ROIs in two ways: first, as chemical weapons and, second, as signalling molecules triggering a variety of biological processes, such as gene expression and proliferation control. How macrophages and neutrophils produce high amounts of ROIs as a first line of defense against invading microorganisms has been reviewed extensively [2, 3]. However, the roles of ROIs as signalling molecules are just beginning to be discovered. On the one hand, immune cells but also

many other types of cells can transiently produce low levels of ROIs in response to distinct extracellular stimuli. These ROI levels do not suffice to kill other cells, but may have a hormone-like function. On the other hand, cells can respond to slightly enhanced ROI levels and use them as second messengers in signal transduction processes. ROIs can trigger the activation of protein kinases and phosphatases, regulate gene expression, and control cell proliferation and death. In this commentary, we will review very recent studies showing that ROIs and oxidation/reduction (redox) processes are critically involved in controlling cellular functions, particularly in T cells.

Mechanisms protecting from ROIs

With the use of dioxygen as electron acceptor in respiration, it was mandatory to coevolve mechanisms protecting aerobic life from the detrimental effects of ROIs. A first line of defense was the development of effective antioxidative molecules, that either detoxify ROIs (GSH, SODs, catalase, thioredoxin), block free radical chain reactions (tocopherol, ascorbate) or bind transition metals (transferrin, coeruloplasmin). Among the most important antioxidants is GSH (L- γ -glutamyl-L-cysteinylglycine). GSH reduces peroxides by the action of glutathione peroxidase and is regenerated by NADPH-dependent glutathione reductase [4, 5]. Inside the cell, millimolar amounts of GSH are present mainly in the reduced form, although even small changes in the ratio of GSH to GSSG may have significant biological effects. GSH/GSSG ratios can vary enormously between intracellular compartments, being the highest (100:1) in the nucleus and the lowest (2:1) in the endoplasmic reticulum [6]. Cytoplasmic GSH levels are determined largely by the availability of its precursor, L-cysteine. The most important source of L-cysteine in the blood plasma is L-cystine, the disulfide-linked oxidized dimer. In contrast to most other cell types, T lymphocytes have a very low transport activity for L-cystine and, hence, a low baseline supply with the GSH precursor [7, 8]. As a consequence, T cells are highly susceptible toward changes in the intracellular redox state.

Another important antioxidative molecule, whose role in immunoregulatory processes is becoming increasingly elucidated, is thioredoxin [9–11]. Thioredoxin is a relatively small protein with two redox active sulfhydryl groups. It is largely cytoplasmic, but it can be secreted and taken back up into the cytosol by lymphocytes under certain conditions. Although both the GSH and thioredoxin systems catalyze the reduction of disulfides, there

† Corresponding author: Dr. Klaus Schulze-Osthoff, Institute of Biochemistry, Albert-Ludwigs-University, Hermann-Herder-Str. 7, D-79104 Freiburg, Germany. Tel. 49-761-203-5221; FAX 49-761-203-5257.

§ Abbreviations: AP-1, activator protein-1; ATL, adult T cell leukemia; EGF, epidermal growth factor; GSH, glutathione; GSSG, glutathione disulfide; IL, interleukin; HIV, human immunodeficiency virus; LPS, lipopolysaccharide; LTR, long terminal repeat; MHC, major histocompatibility complex; NF-AT, nuclear factor activated T cells; NF- κ B, nuclear factor- κ B; PDTC, pyrrolidine dithiocarbamate; ROI, reactive oxygen intermediate; SOD, superoxide dismutase; SRE, serum response element; SRF, serum response factor; TCF, ternary complex factor; TCR, T cell receptor; and TNF, tumor necrosis factor.

is one significant difference. GSH is more effective in reducing small disulfides, whereas thioredoxin is more efficient in reducing (poly)peptides with exposed disulfides. The principal role of thioredoxin appears to be its protein oxidoreductase activity.

The oxidant stress response

When cells are treated with H_2O_2 , they become resistant toward subsequent higher amounts of ROIs that would be lethal without pretreatment [12]. This observation indicates that cells can activate an adaptive genetic program against oxidative stress [13]. The genetic response to oxidative stress has been studied intensively in bacteria [reviewed in Refs. 14–16]. In *Escherichia coli*, the synthesis of some 80 proteins is induced upon exposure to H_2O_2 or $O_2^{\cdot-}$ -generating chemicals. Two transcription factor systems, OxyR and SoxR/S, have been identified that activate expression of genes whose products are involved in either the protection from oxidative stress or the repair of ROI-mediated damages. Intriguingly, the two transcription factors respond to different ROI species. OxyR is activated by H_2O_2 , whereas SoxR/S is selectively induced by superoxide anions. Transcription factors that sense changes in dioxygen or ROI concentration have also been found in yeast and plants. In *Saccharomyces cerevisiae*, binding of a heme group, whose synthesis requires oxygen, activates the transcription factor HAP-1, which controls transcription of the catalase and SOD gene [17]. In plants, salicylic acid (SA) is known as a mediator for systemic acquired resistance, an equivalent of the vertebrate immune response. Recently, the SA-binding protein has been shown to contain catalase activity, which decreases significantly upon ligand binding and thereby results in the accumulation of H_2O_2 [18]. Hence, the plant immune response involves generation of ROI messengers by ligand-induced inhibition of an ROI-eliminating enzyme.

In mammalian cells, the mechanisms by which ROIs are sensed or inducibly produced are less well understood, and transcription factors that are exclusively activated by ROIs or that selectively control expression of ROI-protective and repair enzymes have not been identified. In T lymphocytes, most of the genes induced by oxidative stress can be equally activated by more biological signals, such as cytokines or T cell mitogens. The overlapping effect of both chemical and biological signals may be explained by the fact that some cytokines and T cell mitogens seem to initiate ROI production themselves and utilize ROIs as intracellular signalling molecules. In the past years, increasing evidence has suggested that ROI-induced gene expression plays a widespread role in inflammation, injury and infection.

NF- κ B: A redox effector of the immune system

NF- κ B was the first eukaryotic transcription factor shown to respond directly to oxidative stress [19]. NF- κ B plays a key role in the regulation of numerous genes involved in pathogen responses and cellular defense mechanisms. Many immunologically relevant genes, in particular those encoding cytokines and cytokine receptors, growth factors, and cell adhesion molecules, contain functional NF- κ B binding sites in their promoter and enhancer regions [for actual lists, see Refs. 20 and 21]. In T lymphocytes, gene expression of IL-2 and the IL-2 receptor α chain have been shown to be controlled

by NF- κ B. Furthermore, some retroviruses, such as HIV-1, contain κ B binding sites in their regulatory LTR region and thus employ the transcriptional activity of this host factor for induction of proviral expression and viral replication [22].

NF- κ B activation does not require new protein synthesis. A great variety of mainly pathogenic or pro-inflammatory stimuli, such as LPS, viral infection, UV and γ -irradiation, and the cytokines IL-1 and TNF, activate the transcription factor within minutes. In addition, virtually all inducers of T cell activation, such as ligation of CD3, CD2, and CD28 and infection with T cell lymphotropic viruses, such as HTLV-1 and HIV-1 and -2, activate NF- κ B [21].

By measurement of ROI production, GSH depletion, and other indicators of oxidative stress, it has been shown that basically all inducers of NF- κ B also induce a transient ROI formation. These observations indicated that ROIs are presumably a common messenger required for NF- κ B activation. Indeed, several studies have reported that activation of NF- κ B by all stimuli known to date is blocked by various antioxidants, including GSH precursors (*N*-acetylcysteine, L-cysteine), metal chelators, thiols, dithiocarbamates, and vitamin E and derivatives [reviewed in Ref. 13]. Further evidence for an essential role of ROIs came from experiments using exogenously added prooxidants. In Jurkat and other T cell lines, addition of 100–300 μ M H_2O_2 or butylperoxide resulted in the rapid activation of NF- κ B [19, 23]. Incubation with various superoxide, hydroxyl radical, or NO-generating compounds, however, failed to induce activation, suggesting that, like the bacterial transcription factor OxyR, NF- κ B activation is selectively mediated by peroxides. Compelling genetic evidence for an involvement of H_2O_2 came from a recent study with catalase- and Cu/Zn superoxide dismutase (Cu/Zn-SOD)-overexpressing cell lines. In a catalase-overexpressing cell line, NF- κ B activation was substantially suppressed compared with the parental cell line, whereas it was superinduced in a Cu/Zn-SOD-overexpressing cell line [24]. Likewise, overexpression of thioredoxin or addition of recombinant thioredoxin to the cell culture medium prevented NF- κ B activation by the protein kinase C activator phorbol 12-myristate 13-acetate [25].

Inducible ROI production

How do the diverse stimuli of NF- κ B activation induce ROI formation? Since a great variety of agents activate the transcription factor, it is very likely that oxidative stress is elicited by more than one means. In the case of TNF, a very potent activator of NF- κ B, it has been suggested that the respiratory chain of mitochondria is the site of ROI formation, which is important for the cytotoxicity as well as NF- κ B activation by TNF [26, 27]. In cells lacking a functional mitochondrial respiratory chain as the result of drug action or organelle depletion, NF- κ B activation was suppressed significantly.

In T cells, a potent NF- κ B activation usually requires two signals, one triggering the TCR and another one, a co-stimulatory molecule [28]. It was observed recently that triggering CD28, a major co-stimulatory molecule, causes the rapid decrease of GSH within minutes and enhances ROI formation, which is maintained for 16–24 hr after cell stimulation [29, 30]. This long-lasting prooxidant stimulus may explain why, in contrast to many other cell types, NF- κ B activation in T cells is

generally observed for a rather prolonged period. Inhibitor studies have demonstrated that the dioxygenase activity of 5-lipoxygenase is involved in inducing the prooxidant signal observed upon CD28 ligation [30].

Another principle of ROI formation has been demonstrated for the Tat protein, a major transactivating factor of HIV-1. The potentiation of NF- κ B activation by Tat has been attributed to an increased ROI formation by Tat-mediated transcriptional inhibition of the manganese-dependent SOD gene [31]. This situation is reminiscent of the plant system where a redox signal is produced by inhibition of catalase, another ROI-detoxifying enzyme.

ROI-induced intracellular signal transduction

Some steps in the signal transduction pathway of NF- κ B activation were elucidated recently, but the precise mode of how ROIs activate NF- κ B is still largely unknown. Activated nuclear NF- κ B is frequently a heterodimer of a 50 kDa DNA-binding (p50) and a 65 kDa DNA-binding and transactivating subunit (p65 or RelA). In activated T cells, NF- κ B complexes are often composed of a p50/c-Rel heterodimer, whereas in unstimulated cells a transcriptionally inactive p50 homodimer can be found that occupies κ B sites and thus may inhibit access of active complexes to DNA [32]. In most cell types, NF- κ B is present in the cytoplasm as an inactive complex bound to a third inhibitory subunit, called I κ B. Activation of cells results in the rapid release of I κ B, which allows NF- κ B to translocate to the nucleus and to bind to its cognate DNA sequences. It has been demonstrated that NF- κ B activation coincides with the rapid proteolytic loss of I κ B. The degradation of I κ B is absolutely necessary because various protease inhibitors prevent NF- κ B activation [33–35]. The degradation of I κ B- α is preceded by phosphorylation on serines 32 and 36 [36]. This modification is not sufficient to release I κ B from NF- κ B but dramatically increases the turnover of I κ B, presumably by enhancing the conjugation with ubiquitin [37].

Where in these steps of NF- κ B activation do ROIs play a role? Unlike the bacterial peroxide sensor OxyR that can be activated by oxidation *in vitro*, oxidation of purified NF- κ B–I κ B does not activate but even causes a loss of DNA-binding activity [38]. A conserved cysteine residue in the DNA-binding domain of NF- κ B subunits seems to be responsible for this oxidative inhibition [39]. However, since GSH is highly reduced in the nucleus, the physiological significance of this mechanism remains unclear.

A recent study demonstrated that the phosphorylation of I κ B- α is blocked by the antioxidant PDTC [35]. This hinted at the possibility that either an I κ B kinase was activated or that an I κ B phosphatase, which constantly counteracts a constitutive I κ B phosphorylation, was inactivated. In light of the observation that serines 32 and 36 constitute sites for the constitutive casein kinase type II, the latter scenario appears more likely. It is very possible that the kinase or phosphatase sensing changes in ROI levels is not directly involved in modifying I κ B but is an upstream component of a more complex signaling pathway. There are several kinases that have been reported to be activated by H₂O₂ and other inducers of oxidative stress, including the MAP kinase-related Jun kinases (JNKs) or the tyrosine kinases LTK, EGF receptor and Src kinases [40–45].

Various studies indicated that NF- κ B activation involves a cascade of membrane-bound and cytoplasmic protein kinases. For instance, UV irradiation rapidly activates the tyrosine kinase activity of c-Src, an event followed by the activation of the membrane-bound GTP binding protein H-Ras and the Ser/Thr-specific Raf-1 kinase [45]. Overexpression of transdominant negative mutants of these molecules inhibited UV-induced NF- κ B activation. Likewise, overexpression of v-Src activated NF- κ B in a T cell line [46].

In T lymphocytes, treatment with H₂O₂ results in the rapid and extensive tyrosine phosphorylation of multiple proteins. At least two Src kinases, p56^{lck} and p59^{lyn}, are activated by oxidizing agents [43, 44]. Although it is unclear how these kinases are coupled to the NF- κ B pathway, they may provide a useful model for ROI-induced signal transduction. We have observed recently that, similar to NF- κ B, p56^{lck} and p59^{lyn} are not directly activated by H₂O₂ in a cell-free system or using recombinant kinases. Strong activation, however, could be seen by the addition of GSSG, indicating that, presumably, the formation of a mixed disulfide, a reaction called S-thiolation, may be involved in oxidant-induced kinase activation (Schenk H, Dröge W and Schulze-Osthoff K, unpublished results). Since several inducers of T cell activation lead to an accumulation of GSSG, S-thiolation of a critical cysteine residue may be a physiological mechanism controlling activation of Src-related kinases and other kinases or phosphatases.

Direct redox regulation of transcription factors

NF- κ B is not the only transcription factor responding to changes in ROI concentrations. As seen with NF- κ B, several unrelated factors with different DNA-binding domains contain also cysteine residues whose oxidation state affects DNA binding. Such redox-sensitive cysteines have been found in the basic leucine zipper proteins c-Fos and c-Jun, the helix-loop-helix transcription factor USF, the zinc finger protein Egr-1, and other factors, such as c-Myb and p53 [reviewed in Ref. 13]. These proteins are inactivated by oxidation *in vitro*, and mutation of the critical cysteine residue to a serine residue creates proteins that are resistant toward oxidative inactivation. Since the reduced state of the conserved cysteines is critical for DNA binding, it has been speculated that redox modification of the DNA-binding domain may be a secondary mechanism of controlling transcriptional activity. However, all present data concerning the direct modification of transcription factors by cysteine oxidation rely on cell-free experiments, and there are only vague hints supporting a physiological role. A ubiquitous nuclear protein, called Ref-1, has been identified, which restores the DNA-binding activity of oxidized Fos and Jun proteins *in vitro* [47]. Ref-1 catalyzes the reduction of thiol groups in these proteins, using thioredoxin as a cofactor.

Antioxidant-induced signal transduction

Heterodimers of members of the Jun and Fos transcription factor family constitute the transcription factor AP-1, for which DNA-binding motifs (consensus sequence: 5'-TGANTCA-3') can be found in several cytokine genes [48]. Many conditions inducing oxidative stress result in the increased transcription of the *c-fos* and *c-jun* genes. In contrast, however, only a very moderate induction of AP-1 DNA-binding and transactiva-

tion activity is usually observed [49, 50]. Thus, the appearance of mRNAs for Jun and Fos does not necessarily correlate with the activation of AP-1 DNA binding. Exposure of some T cell lines to H_2O_2 results in virtually no change of AP-1 DNA binding activity, while NF- κ B is strongly activated [23]. Surprisingly, a strong activation of AP-1 could be observed after treatment of cells with several structurally unrelated antioxidants, including PDTC, *N*-acetyl cysteine, and butylated hydroxyanisole [25, 50]. A significant increase in AP-1 DNA-binding and transactivating activity was also detected after transient overexpression or addition of recombinant thioredoxin, while NF- κ B activation was inhibited under the same conditions. The physiological equivalent to the antioxidant treatment may be hypoxia. We* and others [51] have observed that HeLa cells strongly activate AP-1 in a biphasic response under reduced oxygen pressure. In contrast to a previous report [52], this does activate NF- κ B. We have observed a very strong activation of NF- κ B only upon subsequent reoxygenation of cell cultures.

The induction of AP-1 by antioxidant treatments requires *de novo* protein synthesis of the c-Jun and c-Fos proteins, indicating that yet other transcription factors are responsible for the activation of AP-1 in response to cellular redox changes. In the case of the *c-fos* gene, this pre-existing factor is TCF/Elk-1; in the case of the *c-jun* gene, it is a c-Jun homodimer or a c-Jun/ATF-2 heterodimer. TCF/Elk-1 forms a ternary complex with the SRF on the SRE [53, 54]. Phosphorylation of the transactivation domain of TCF/Elk-1 by mitogen-activated kinases (MAPKs or ERKs) induces the transactivating potential of the complex and leads to *c-fos* gene expression. We have found that the SRE is responsible for the activation of *c-fos* expression in response to both antioxidant and prooxidant stimulation of cells [50]. Such treatments cause an increased TCF/Elk-1 phosphorylation in the cell nucleus, which is preceded by novel phosphorylation and activation of MAP kinases.† TCF/Elk-1 can be considered as a nuclear sensor of the cellular redox imbalance. It is currently unknown which kinases (or phosphatases) in the cascade cause phosphorylation of MAP kinases in response to antioxidants or oxidants. One candidate for an H_2O_2 -responsive upstream kinase is the EGF receptor tyrosine kinase [42].

Preexisting homodimers of c-Jun and heterodimers of c-Jun and ATF-2, which induce the *c-jun* gene in response to prooxidant stimuli and other stresses, are activated by *de novo* phosphorylation of c-Jun in its N-terminal domain [55]. One important kinase is JNK, which activates c-Jun in response to UV light, presumably, depending on redox changes [40]. It is currently unknown what activates *c-jun* transcription in response to antioxidant stimuli. A candidate factor is NF-Jun [56]. Since strong AP-1 activation is observed under antioxidant but not under prooxidant conditions, additional posttranscriptional control steps seem to be involved.

The expression of most genes is controlled by the concerted action of multiple distinct transcription factors. For instance, the promoter of the IL-2 gene contains

DNA-binding sites for at least four distinct factors: NF- κ B, AP-1, NF-AT, and Oct factors [28, 57]. Some of these factors can physically interact among each other when bound to DNA and thereby synergistically promote their DNA-binding and transactivating activity. NF- κ B was found to interact directly with AP-1 subunits [58]. NF-ATp, an NF- κ B/Rel-related transcription factor activated by the calcium-dependent phosphatase calcineurin, also binds AP-1 and recruits the factor to DNA [59]. These physical interactions with redox-controlled factors will also engage other transcription factors in redox regulation.

Redox-sensitive T cell functions

Since the redox-controlled transcription factors NF- κ B and AP-1 are used extensively for signalling and gene induction in immune cells, it is mandatory that various T cell functions are redox dependent. Indeed, while processes occurring early during T cell activation are facilitated by mild or transient prooxidant conditions, late events appear to require reducing or antioxidant conditions. In contrast to bacteria, the ROI response in T cells does not primarily involve cellular defense or repair mechanisms but is associated with more physiological processes, such as signal transduction, proliferation, cytokine synthesis, and other immune regulatory processes. This indicates that during evolution T cells did not only acquire the capability to sense the redox state of the environment but also to use it for the control of specific cellular responses.

The role of ROIs in early T cell activation was studied by the regulation of cytokine and cytokine receptor gene expression. Many gene products produced by activated T cells, including IL-2, GM-CSF, TNF- α , IL-6, IL-2R α chain, and MHC class I genes, require NF- κ B for their inducible transcriptional control [reviewed in Ref. 21]. It has been reported that micromolar concentrations of H_2O_2 potentially activate NF- κ B in T cell lines [19], as well as the transcription of the IL-2 and IL-2R α chain genes, which is conferred by κ B regulatory sites in these genes [23, 60]. Since neutrophils and macrophages secrete large amounts of ROIs in inflamed tissues, exogenously produced ROIs are likely to play a hormone- or mediator-like role in early T cell activation. Furthermore, activation of cultured T cells by stimulation of the TCR or co-stimulatory molecules rapidly induces an intracellular production of ROIs, as measured by the induction of H_2O_2 production or GSH depletion [23, 29, 30]. The regulatory role of this ROI production is shown by the strong inhibitory effect of distinct antioxidants on the expression of cytokine genes following T cell activation. IL-2 and the IL-2R α chain are examples of genes inhibited by antioxidants in peripheral blood lymphocytes [60, 61]. Antioxidants also have been reported to cause the arrest of thymocyte differentiation, an event associated with a profound decrease of NF- κ B and TCF-1 activation [62]. The requirement of ROIs for the expression of NF- κ B-regulated cytokine and other genes is not restricted to T cells. It was also observed for endothelial cells, various tumor cell lines, fibroblasts, keratinocytes, macrophages, and pre-B cells.

ROIs seem to be further involved in early IL-2-independent steps of T cell proliferation. Studies of mixed lymphocyte cultures have revealed that several agents that inhibit the effects of ROIs also suppress lymphocyte proliferation [63, 64]. In some T cell lines, a short ex-

* Rupec R and Baeuerle PA, Manuscript submitted for publication.

† Müller J, Cahill M, Baeuerle PA and Nordheim A, Manuscript submitted for publication.

posure to low amounts of H_2O_2 causes a marked increase in DNA synthesis [23, 65]. The mitogenic effect of ROIs is activation dependent and only observed in the absence of TCR stimulation. This indicates that a strong mitogenic stimulus either bypasses the requirement of ROIs or induces ROI formation by itself.

ROIs seem to be required only for the early phases of the T cell cycle. Antioxidants strongly inhibit the proliferation of mitogenically activated PBLs in the first 2 hr but not at later time points after mitogenic stimulation, indicating that preferentially the transition from the G_0 to the G_1 phase is blocked by these agents [66]. Moreover, prolonged prooxidant treatment is lethal to T cells. In fact, the most optimal mitogenic stimulation by H_2O_2 is achieved by the subsequent addition of catalase [23].

Later stages of the T cell cycle are promoted by reducing/antioxidant conditions. Thiols, such as 2-mercaptoethanol, increase proliferation in response to T cell mitogens and are therefore routinely added to the culture medium. Thioredoxin, also called "ATL-derived factor," has been identified as an autocrine growth factor produced by ATL cells [11]. Thioredoxin exerts co-stimulatory activity and increases cytokine expression in response to phorbol esters or TCR stimulation. Experiments using GSH depletion have shown that a number of T cell functions are correlated with their thiol levels. Even a partial depletion of the intracellular GSH pool has been found to inhibit IL-2-dependent cytotoxic T cell activity and the generation of lymphokine-activated killer (LAK) cells [67–69]. Therefore, administration of antioxidants may provide a measure to increase the antitumor efficiency in IL-2/LAK cell therapy.

Conclusions

Changes in the cellular redox state toward either prooxidant or antioxidant conditions have profound effects on cellular functions. T lymphocytes are particularly well-studied in this respect. It appears that early T cell functions with a proinflammatory character (such as cytokine gene induction) are promoted by prooxidant conditions, whereas later functions, such as proliferation and maintained expression of genes, are inhibited by prooxidant but promoted by antioxidant conditions. This suggests that there is a requirement of delicate cellular redox shifts for effective T cell activation. Given our detailed knowledge about the roles and characteristics of transcription factors NF- κ B and AP-1 (in conjunction with NF-AT) in T cell function, the following scenario emerges. Immediate-early T cell functions require a transient production of ROIs in response to stimulation of TCR and co-stimulatory molecules. This condition is required to activate transcription factor NF- κ B, which will initiate the transcription of cytokine (IL-2, TNF) and receptor genes (IL-2R α chain). For progression to later T cell functions, it seems mandatory that the potentially harmful prooxidant phase is only transient and restored to normoxic or even antioxidant conditions. It is under these latter conditions that the transcription factor AP-1 becomes activated by new synthesis. AP-1 is composed of proto-oncogene products and is involved in the positive control of proliferation. It is thus tempting to speculate that the requirement of reducing conditions for later T cell functions is partly due to the optimal activation of AP-1 under these conditions. An interesting possibility is that generally inflammatory processes rely on

prooxidant conditions and NF- κ B, whereas proliferative processes depend on antioxidant conditions and AP-1.

Acknowledgements—The authors thank W. Dröge for fruitful collaboration and helpful discussions.

REFERENCES

- Halliwell B and Gutteridge JMC, The role of free radicals and catalytic metal ions in human disease: An overview. *Methods Enzymol* **186**: 1–85, 1990.
- Rossi F, The O_2^- -forming NADPH oxidase from phagocytes: Nature, mechanism of activation and function. *Biochim Biophys Acta* **853**: 65–89, 1986.
- Morel F, Doussiere J and Vignais PV, The superoxide-generating oxidase of phagocytic cells. Physiological, molecular and pathological aspects. *Eur J Biochem* **201**: 523–546, 1991.
- Meister A and Anderson ME, Glutathione. *Annu Rev Biochem* **52**: 711–760, 1983.
- Dröge W, Schulze-Osthoff K, Mihm S, Galter D, Schenk H, Eck H-P, Roth S and Gmünder H, Functions of glutathione and glutathione disulfide in immunology and immunopathology. *FASEB J* **14**: 1131–1138, 1994.
- Hwang C, Sinskey AJ and Lodish HF, Oxidized redox state of glutathione in the endoplasmic reticulum. *Science* **257**: 1496–1502, 1992.
- Ishii T, Sugita Y and Bannai S, Regulation of glutathione levels in mouse spleen lymphocytes by transport of cysteine. *J Cell Physiol* **133**: 330–336, 1987.
- Gmünder H, Eck H-P and Dröge W, Low membrane transport activity for cystine in resting and mitogenically stimulated human lymphocyte preparations and human T cell clones. *Eur J Biochem* **201**: 113–117, 1991.
- Holmgren A, Thioredoxin. *Annu Rev Biochem* **54**: 237–271, 1985.
- Holmgren A, Thioredoxin and glutaredoxin systems. *J Biol Chem* **264**: 13963–13966, 1989.
- Yodoi J and Uchiyama T, Diseases associated with HTLV-I virus, IL-2 receptor dysregulation and redox regulation. *Immunol Today* **13**: 405–411, 1992.
- Janssen YMW, Van Houten B, Borm PJA and Mossman BT, Biology of disease. Cell and tissue responses to oxidative damage. *Lab Invest* **69**: 261–274, 1993.
- Schulze-Osthoff K and Baeuerle PA, Redox regulation in gene expression. In: *Oxyradicals in Medical Biology* (Ed. McCord J), JAI Press, Greenwich, CT, in press.
- Storz G, Tartaglia LA, Farr SB and Ames BN, Bacterial defenses against oxidative stress. *Trends Genet* **6**: 363–368, 1990.
- Demple B and Amabile-Cuevas CF, Redox redux: The control of oxidative stress responses. *Cell* **67**: 837–839, 1991.
- Pahl HL and Baeuerle PA, Oxygen and the control of gene expression. *Bioessays* **16**: 497–502, 1994.
- Zitomer RS and Lowry CV, Regulation of gene expression by oxygen in *Saccharomyces cerevisiae*. *Microbiol Rev* **56**: 1–11, 1992.
- Chen Z, Silva H and Klessing DF, Active oxygen species in induction of plant systemic acquired resistance by salicylic acid. *Science* **262**: 1883–1886, 1993.
- Schreck R, Rieber P and Baeuerle PA, Reactive oxygen species intermediates as apparently widely used messengers in the activation of the NF- κ B transcription factor and HIV-1. *EMBO J* **10**: 2247–2258, 1991.
- Grilli M, Chiu JJ-S and Lenardo MJ, NF- κ B and Rel: Participants in a multifactorial transcriptional regulatory system. *Int Rev Cytol* **143**: 1–62, 1993.
- Baeuerle PA and Henkel T, Function and activation of NF- κ B in the immune system. *Annu Rev Immunol* **12**: 141–179, 1994.
- Nabel G and Baltimore D, An inducible transcription factor

- activates expression of human immunodeficiency virus in T cells. *Nature* **326**: 711–713, 1987.
23. Los M, Dröge W, Stricker K, Baeuerle PA and Schulze-Osthoff K, Hydrogen peroxide as a potent activator of T-lymphocyte functions. *Eur J Immunol* **25**: 159–165, 1995.
 24. Schmidt KN, Amstad P, Cerutti P and Baeuerle PA, The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF- κ B. *Chem Biol* **2**: 13–22, 1995.
 25. Schenk H, Klein M, Erdbrügger W, Dröge W and Schulze-Osthoff K, Distinct effects of thioredoxin and other antioxidants on the activation of NF- κ B and AP-1. *Proc Natl Acad Sci USA* **91**: 1672–1676, 1994.
 26. Schulze-Osthoff K, Bakker AC, Vanhaesebroeck B, Beyaert R, Jacobs WA and Fiers W, Cytotoxic activity of tumor necrosis factor is mediated by early damage of mitochondrial functions. Evidence for the involvement of mitochondrial radical generation. *J Biol Chem* **267**: 5317–5323, 1992.
 27. Schulze-Osthoff K, Beyaert R, Vandevoorde V, Haegeman G and Fiers W, Depletion of the mitochondrial electron transport abrogates the cytotoxic and gene-inductive effects of TNF. *EMBO J* **12**: 3095–3104, 1993.
 28. Crabtree GR and Clipstone NA, Signal transmission between the plasma membrane and nucleus of T lymphocytes. *Annu Rev Biochem* **63**: 1045–1083, 1994.
 29. Costello R, Lipcey C, Algarte M, Cerdan C, Baeuerle PA, Olive D and Imbert J, Activation of primary human T lymphocytes through CD2 plus CD28 adhesion molecules induces long-term nuclear expression of NF- κ B. *Cell Growth Differ* **4**: 329–339, 1993.
 30. Los M, Schenk H, Hexel K, Baeuerle PA, Dröge W and Schulze-Osthoff K, IL-2 gene expression and NF- κ B activation through CD28 requires reactive oxygen production by 5-lipoxygenase. *EMBO J*, in press.
 31. Westendorp MO, Shatrov VA, Schulze-Osthoff K, Frank R, Kraft M, Los M, Krammer PH, Dröge W and Lehmann V, HIV-1 Tat potentiates TNF-induced NF- κ B activation and cytotoxicity by altering the cellular redox state. *EMBO J* **14**: 546–554, 1995.
 32. Kang S-M, Tran A-C, Grilli M and Lenardo MJ, NF- κ B subunit regulation in nontransformed CD4⁺ T lymphocytes. *Science* **256**: 1452–1456, 1992.
 33. Henkel T, Machleidt T, Alkalay I, Krönke M, Ben-Neriah Y and Baeuerle PA, Rapid proteolysis of I κ B- α is necessary for activation of transcription factor NF- κ B. *Nature* **365**: 182–185, 1993.
 34. Palombella VJ, Rando OJ, Goldberg AL and Maniatis T, The ubiquitin-proteasome pathway is required for processing the NF- κ B precursor protein and the activation of NF- κ B. *Cell* **78**: 773–785, 1994.
 35. Traenckner EB-M, Wilk S and Baeuerle PA, A proteasome inhibitor prevents activation of NF- κ B and stabilizes a newly phosphorylated form of I κ B- α that is still bound to NF- κ B. *EMBO J* **13**: 5433–5441, 1994.
 36. Traenckner EB-M, Pahl HL, Schmidt KN, Wilk S and Baeuerle PA, Phosphorylation of human I κ B- α on serines 32 and 36 controls I κ B- α proteolysis and NF- κ B activation in response to diverse stimuli. *EMBO J* **14**: 2876–2883, 1995.
 37. Traenckner EB-M and Baeuerle PA, Appearance of ubiquitin-conjugated forms of I κ B- α during its phosphorylation-induced degradation. *J Cell Sci*, in press.
 38. Toledano MB and Leonard WJ, Modulation of transcription factor NF- κ B binding by oxidation-reduction *in vitro*. *Proc Natl Acad Sci USA* **88**: 4328–4332, 1991.
 39. Kumar S, Rabson AB and Gélinas C, The RxxRxxC motif conserved in all Rel κ B proteins is essential for the DNA-binding activity and redox regulation of the v-Rel oncoprotein. *Mol Cell Biol* **12**: 3094–3106, 1992.
 40. Kyriakis JM, Banerjee P, Nikolakaki E, Dai T, Rubie EA, Ahmad MF, Avruch J and Woodgett JR, The stress-activated protein kinase subfamily of c-Jun kinases. *Nature* **369**: 156–160, 1994.
 41. Bauskin AR, Alkalay I and Ben-Neriah Y, Redox regulation of a protein tyrosine kinase in the endoplasmic reticulum. *Cell* **66**: 685–696, 1991.
 42. Gamou S and Shimizu N, Hydrogen peroxide preferentially enhances the tyrosine phosphorylation of epidermal growth factor receptor. *FEBS Lett* **357**: 161–164, 1995.
 43. Schieven GL, Kirihaara JM, Myers DE, Ledbetter JA and Uckun FM, Reactive oxygen intermediates activate NF- κ B in a tyrosine kinase-dependent mechanism and in combination with vanadate activate the p56^{lck} and p59^{lyn} tyrosine kinases in human lymphocytes. *Blood* **82**: 1212–1220, 1993.
 44. Nakamura K, Hori T, Sato N, Sugie K, Kawakami T and Yodoi J, Redox regulation of a src family protein tyrosine kinase p56^{lck} in T cells. *Oncogene* **8**: 3133–3139, 1993.
 45. Devary Y, Rosette C, DiDonato JA and Karin M, NF- κ B activation by ultraviolet light not dependent on a nuclear signal. *Science* **261**: 1442–1445, 1993.
 46. Eicher DM, Tan T-H, Rice NR, O'Shea JJ and Kennedy ICS, Expression of v-src in T cells correlates with nuclear expression of NF- κ B. *J Immunol* **152**: 2710–2719, 1994.
 47. Xanthoudakis S, Miao G, Wang F, Pan Y-CE and Curran T, Redox activation of Fos-Jun DNA-binding activity is mediated by a DNA repair enzyme. *EMBO J* **11**: 3323–3335, 1992.
 48. Angel P and Karin M, The role of Jun, Fos, and the AP-1 complex in cell-proliferation and transformation. *Biochim Biophys Acta* **1072**: 129–157, 1991.
 49. Devary Y, Gottlieb RA, Laus LF and Karin M, Rapid and preferential activation of the c-jun gene during the mammalian UV response. *Mol Cell Biol* **11**: 2804–2811, 1991.
 50. Meyer M, Schreck R and Baeuerle PA, H₂O₂ and antioxidants have opposite effects on activation of NF- κ B and AP-1 in intact cells: AP-1 as secondary antioxidant-responsive factor. *EMBO J* **12**: 2005–2015, 1993.
 51. Yao K-S, Xanthoudakis S, Curran T and O'Dwyer PJ, Activation of AP-1 and of a nuclear redox factor, Ref-1, in the response of HT29 colon cancer cells to hypoxia. *Mol Cell Biol* **14**: 5997–6003, 1994.
 52. Koong AC, Chen EY and Giaccia AJ, Hypoxia causes the activation of nuclear factor κ B through phosphorylation of I κ B- α on tyrosine residues. *Cancer Res* **54**: 1425–1430, 1994.
 53. Treisman R, The serum response element. *Trends Biochem Sci* **17**: 423–426, 1992.
 54. Janknecht R and Nordheim A, Gene regulation by Ets proteins. *Biochim Biophys Acta* **1155**: 346–356, 1993.
 55. Binétruy B, Smeal T and Karin M, Ha-Ras augments c-Jun activity and stimulates phosphorylation of its activation domain. *Nature* **351**: 122–127, 1991.
 56. Brach MA, Herrmann F, Yamada H, Baeuerle PA and Kufe DW, Identification of NF-jun, a novel inducible transcription factor that regulates c-jun gene transcription. *EMBO J* **11**: 1479–1486, 1992.
 57. Serfling E, Barthelmäs R, Pfeuffer I, Schenk B, Zarius S, Swoboda R, Mercurio F and Karin M, Ubiquitous and lymphocyte-specific factors are involved in the induction of the mouse interleukin 2 gene in T lymphocytes. *EMBO J* **8**: 465–473, 1989.
 58. Stein B, Baldwin AS Jr, Ballard DW, Greene WC, Angel P and Herrlich P, Cross-coupling of the NF- κ B p65 and Fos/Jun transcription factors produces potentiated biological function. *EMBO J* **12**: 3879–3891, 1993.
 59. Rao A, NF-ATp: A transcription factor required for the co-ordinate induction of several cytokine genes. *Immunol Today* **15**: 274–281, 1994.
 60. Roth S and Dröge W, Regulation of interleukin 2 production, interleukin 2 mRNA expression and intracellular glutathione levels in *ex vivo* derived T lymphocytes by lactate. *Eur J Immunol* **21**: 1933–1937, 1991.
 61. Dornand J and Gerber M, Inhibition of murine T-cell re-

- sponses by anti-oxidants: The targets of lipo-oxygenase pathway inhibitors. *Immunology* **68**: 384–391, 1989.
62. Ivanov V, Merckenschlager M and Ceredig R, Antioxidant treatment of thymic organ cultures decreases NF- κ B and TCF1 α transcription factor activities and inhibits $\alpha\beta$ T cell development. *J Immunol* **151**: 4694–4704, 1993.
63. Novogradsky A, Ravid A, Rubin AL and Stenzel KH, Hydroxyl radical scavengers inhibit lymphocyte mitogenesis. *Proc Natl Acad Sci USA* **79**: 1171–1174, 1982.
64. Chaudhri G, Hunt NH, Clark IA and Ceredig R, Antioxidants inhibit proliferation and cell surface expression of receptors for interleukin-2 and transferrin in T lymphocytes stimulated with phorbol myristate acetate and ionomycin. *Cell Immunol* **115**: 204–213, 1988.
65. Dröge W, Mihm S, Bockstette M and Roth S, Effect of reactive oxygen intermediates and antioxidants on proliferation and function of T lymphocytes. *Methods Enzymol* **234**: 135–151, 1994.
66. Jeitner TM, Kneale CL, Christopherson RI and Hunt NH, Thiol-bearing compounds selectively inhibit protein kinase C-dependent oxidative events and proliferation in human T cells. *Biochim Biophys Acta* **1223**: 15–22, 1994.
67. Dröge W, Pottmeyer-Gerber C, Schmidt H and Nick S, Glutathione augments the activation of cytotoxic T lymphocytes *in vivo*. *Immunobiology* **172**: 151–156, 1986.
68. Yamamauchi A and Bloom ET, Requirement of thiol compounds as reducing agents for IL-2-mediated induction of LAK activity and proliferation of human NK cells. *J Immunol* **151**: 5535–5544, 1993.
69. Yim C-Y, Hibbs JB Jr, McGregor JR, Galinsky RE and Samlowski WE, Use of *N*-acetyl cysteine to increase intracellular glutathione during the induction of antitumor responses by IL-2. *J Immunol* **152**: 5796–5805, 1994.