Linköping University medical Dissertations No. 1479

Birth Characteristics' Impact on Future Reproduction and Morbidity Among Twins and Singletons

Marie Bladh

Faculty of Medical and Health Sciences

Department of Obstetrics and Gynecology
Department of Clinical and Experimental Medicine
SE-581 85 Linköping, Sweden

Linköping 2015

© Marie Bladh, 2015

ISBN: 978-91-7685-961-2

ISSN: 0345-0082

Published articles have been reprinted with permission from the publishers: Twin
Reserach and Human Genetics

Printed in Sweden by LiU-tryck, Linköping 2015

*We must welcome the future,
remembering that soon it will be
the past; and we must respect
the past, remembering that it
was once all that was humanly
possible.*

George Santayana

ABSTRACT

Globally, in both developed and developing countries, the twinning rates have increased since the early 70's. A large proportion of twins are born preterm and/or small-for-gestational-age (SGA) and/or with a low birth weight. Several studies have been performed on the long-term effect of these non-optimal birth characteristics on future reproductive performance and morbidity. Yet, most studies exclude twins or higher order pregnancies and thus the findings are based on singleton pregnancies only.

The aim of the present thesis was therefore to investigate the impact of non-optimal birth characteristics in terms of preterm birth, small-for-gestational age, and low birth weight, on the reproductive pattern and morbidity among twins and singletons. Furthermore, the present thesis attempted to establish whether twins and singletons were affected in the same manner.

The studies included in this thesis are prospective population-based register studies, including all men and women, alive and living in Sweden at age 13, who were born between 1973 and 1983 (1,000,037 singletons and 16,561 twins) for the first three studies with follow-up till the end of 2006 and 2009. The last study included all men and women, alive and living in Sweden at age 13, who were born between 1973 and 1993 (2,051,479 singletons and 39,726 twins) with follow-up till the end of 2012.

In general, twins were found less likely to reproduce between 13 and 33 years of age compared with singletons. Stratifying data by different birth characteristics, it was found that twins had a lower likelihood of reproducing on several different birth characteristics (appropriate-for-gestational-age, normal birth weight, low birth weight, term birth, preterm birth). However, twins born very preterm had an increased likelihood of reproducing compared with singletons born very preterm.

Not taking birth characteristics into account, twinning was associated with a higher degree of hospitalization. However, accounting for the diverging birth characteristics this difference diminished and for some diagnoses the relationship was reversed such that twins were actually less likely to be hospitalized compared with singletons.

In terms of the heritability of non-optimal birth characteristics singleton mothers born preterm were more predisposed to give birth to a child that was preterm while singleton mothers born SGA more often gave birth to a child either born preterm or SGA. Among twins this heritability was not as evident. The only difference observed was among twin mothers born SGA who were more likely to give birth to a child born SGA.

In the extended cohort comprising those born between 1973 and 1993, male and female twins were found to be less likely to become parents compared with singletons. No difference was found among women in terms of having a second child, while male twins were more likely to have a second child compared with male singletons. It was also found that the likelihood of becoming a first-time parent and second-time parent was positively associated with the number of siblings.

Keywords: Twin, singleton, preterm birth, SGA, reproduction, hospitalization, siblings

CONTENTS

ABSTRACT	3
LIST OF PUBLICATIONS	7
LIST OF TABLES AND FIGURES	9
ABBREVIATIONS	11
EXPLANATIONS OF KEY TERMS	13
INTRODUCTION	17
BACKGROUND	19
Birth-characteristics in relation to outcomes in later stages of life	21
<i>Twinning</i>	21
<i>Mortality</i>	21
<i>Long-term morbidity</i>	22
<i>Socio-economic and lifestyle factors' effect on future pregnancy outcome</i>	24
<i>Childbearing and pregnancy outcome</i>	24
<i>Heritability and Epigenetics</i>	25
AIMS OF THE PRESENT THESIS	27
MATERIAL AND METHODS	29
Retrieval of data	29
International Classification of Diseases	30
Data sources	31
<i>The Swedish Medical Birth Register</i>	31
<i>The National Patient Register</i>	31
<i>The Cause of Death Register</i>	32
<i>The Total Population Register</i>	32
<i>The Multi-Generation Register</i>	32
<i>The Education Register and the 1970 Population and Housing Census</i>	32
The study population studies I-III	33
The study population study IV	36
Design of the studies	38
<i>Study I</i>	38
<i>Study II</i>	39
<i>Study III</i>	40
<i>Study IV</i>	41
Statistical methods	42
<i>Logistic Regression</i>	42
<i>Cox's proportional hazards model</i>	44
<i>Intra-pair correlation due to twinning</i>	45
<i>Analysis with respect to familial factors</i>	45
Exposure, outcomes and covariates	45
<i>Primary exposures</i>	45
<i>Primary outcomes</i>	47
<i>Covariates/confounders</i>	48
ETHICAL CONSIDERATIONS	51
RESULTS	53
Study I. Likelihood of giving birth among twins and singletons with respect to birth characteristics	53
Study II Hospitalization in adolescence and young adulthood among twins and singletons	54

Study III Intergenerational effect of preterm birth and small-for-gestational-age55

Study IV Reproductive pattern among twins and singletons in relation to number of siblings.....58

DISCUSSION 61

Summary of principal findings..... 61

Methodological considerations..... 62

Principal findings in relation to other research..... 64

Reproductive pattern 65

Hospitalization according to the main chapters in ICD-9 66

Intergenerational effect of non-optimal birth characteristics 66

Reproductive pattern in relation to number of siblings..... 67

GENERAL CONCLUSIONS..... 69

FUTURE PERSPECTIVES..... 71

POPULÄRVETENSKAPLIG SAMMANFATTNING..... 73

ACKNOWLEDGEMENTS..... 75

REFERENCES..... 77

LIST OF PUBLICATIONS

- Study I Bladh M, Josefsson A, Carstensen J, Finnström O, Sydsjö G.
Reproductive patterns among twins – A Swedish register
study of men and women born 1973-1983
BMC Pregnancy Childbirth, 2013. doi: 10.1186/1471-2393-13-
6)
- Study II Bladh M, Carstensen J, Josefsson A, Finnström O, Sydsjö G.
Hospitalization in adolescence and young adulthood among
twins and singletons: a Swedish cohort study of subjects born
between 1973 and 1983
Twin Research and Human Genetics. 2013 Jun;16(3):707-15.
doi: 10.1017/thg.2013.27. Epub 2013 May 10. PMID:
23659898
- Study III Bladh M, Josefsson A, Carstensen J, Finnström O, Sydsjö G.
Intergenerational cohort study of non-optimal birth
characteristics in twins and singletons
Accepted for publication in Twin Research and Human
Genetics
- Study IV Bladh M, Josefsson A, Carstensen J, Finnström O, Sydsjö G.
Reproductive pattern among twins and singletons in relation
to number of siblings – a Swedish cohort study of individuals
born between 1973 and 1993
Submitted

LIST OF TABLES AND FIGURES

List of tables and figures in the order in which they appear in the text.

Figure 1. *Different types of twinning.*

Figure 2. *Areas with long-term effect of preterm birth.*

Figure 3. *Consequences of non-optimal birth characteristics and future reproductive performance in young ages.*

Table 1. *Conversion of ICD-9 to ICD-10.*

Figure 4. *Flowchart of the exclusion of study participants in studies I-III.*

Figure 5. *Flowchart of the exclusion of study participants in study IV.*

Figure 6. *Design of study I.*

Figure 7. *Design of study II.*

Figure 8. *Design of study III.*

Figure 9. *Design of study IV.*

Figure 10. *Calculation of odds ratio.*

Figure 11. *Alternatives on when to enter and exit the Cox proportional hazards model.*

Table 2. *Study population and its distribution of twinning and sex, study I-III.*

Table 3. *Hazard ratio of becoming a parent among females and males in sub-groups defined by birth characteristics, comparing twins vs. singletons.*

Table 4. *Likelihood of being hospitalized stratified by optimality at birth.*

Table 5. *Likelihood of giving birth to a child born preterm or SGA, stratified by twinning status.*

Table 6. *Hazard ratio for becoming a parent in relation to number of siblings, stratified by twinning and sex.*

Table 7. *Hazard ratio of giving birth by number of siblings for twins and singletons.*

ABBREVIATIONS

ADHD	Attention Deficit Hyperactivity Disorder
AGA	Appropriate-for-Gestational-Age
BMI	Body Mass Index
CI	Confidence Interval
FOAD	Fetal Origins of Adult Disease
HR	Hazard Ratio
ICD	International Classification of Diseases
IVF	In-Vitro Fertilization
LBW	Low Birth Weight
LGA	Large-for-Gestational-Age
MBR	Medical Birth Register
NPR	National Patient Register
OR	Odds Ratio
PPV	Positive Predictive Value
PT	Preterm Birth
SD	Standard Deviation
SGA	Small-for-Gestational-Age
TPR	Total Population Register
VLBW	Very Low Birth Weight
VPT	Very Preterm Birth
WHO	World Health Organization

EXPLANATIONS OF KEY TERMS

Twinning

Twinning can be separated into dizygotic and monozygotic, fraternal or identical, twins. This is further complicated by whether the twins share the placenta or not, or as is only possible among monozygotic twins, also share the amniotic sac. See Figure 1 for detailed description of different types of twinning.

<http://twinpossible.com/twin-moms-to-be-want-to-know-are-my-twins-identical-or-fraternal>

Figure 1. *Different types of twinning.*

Birth characteristics

In this thesis birth characteristics refer to information regarding birth weight, gestational age at birth, and size for gestational age.

Non-optimal birth characteristics

Includes the following birth characteristics: small-for-gestational age (SGA), large-for-gestational-age (LGA), low birth weight, or preterm birth.

Optimal birth characteristics

Includes birth characteristics born at term, and appropriate-for-gestational-age.

Hospitalization

In study II hospitalization was studied, as a proxy for morbidity. A registered in- or outpatient visit in The Swedish National Patient Register served as the base for this variable. Two different variants of hospitalization were used a) any hospitalization vs. no hospitalization registered and b) 0-6 hospitalizations vs. more than 6 hospitalizations.

Preterm birth

Children born before gestational week 37.

Very preterm birth

Children born before gestational week 32.

Intra-uterine growth restriction/retardation

The fetus has not been able to achieve its potential size.

Body Mass Index (BMI)

BMI was defined according to the “standard” definition, i.e. as weight (in kilograms) divided by the squared height (in meters). The BMI was further divided into 4 categories:

- 1) Underweight <18.5
- 2) Normal weight 18.5-24.99
- 3) Overweight 25.00 – 29.99
- 4) Obese ≥ 30

Nicotine use

Indicator variable of whether the mother either smoked or used “snuff” 3 months prior to pregnancy or during pregnancy.

Socio-economic characteristics

Factors included were: maternal and paternal level of education, maternal and paternal country of origin, mother's civil status and mother's age when giving birth, parity, BMI, and nicotine use.

Large-for-gestational-age (LGA)

Born with a high birth weight given sex and the gestational age at birth. Defined as a birth weight above 2SDs of the mean birth weight according to the Swedish standard.

Small-for-gestational-age (SGA)

Born with a low birth weight given sex and the gestational age at birth. Defined as a birth weight below 2SDs of the mean birth weight according to the Swedish standard.

Appropriate-for-gestational-age (AGA)

Born with an appropriate birth weight given sex and the gestational age at birth, i.e. those not born SGA or LGA.

Low birth weight

Defined as a birth weight below 2,500 grams.

Very low birth weight

Defined as a birth weight below 1,500 grams.

Sibling

Children born to the same mother.

INTRODUCTION

In the past decades the number of twins has increased globally. In Sweden alone it rose from 8.2 births/1000 women in 1973 to 16.3 births/1000 women in 2003. From 2004 onward the twin birth rate has stabilized around 14 births/1000 women. During the same time period the proportion of preterm births has remained around 6 % in Sweden, though the absolute number has increased. According to the World Health Organization (WHO) it is estimated that approximately 13 million children are born preterm and approximately 20 million children are born with a low birth weight each year around the world. The increasing number of children born preterm and/or with low birth weight in combination with the improved medical care of these children have led to an increasing number of surviving children with non-optimal birth characteristics.

Several studies have established a relationship between non-optimal birth characteristics and long-term sequelae such as mental disorders, neurological disorders (including cerebral palsy), reduced reproduction, and metabolic syndrome, the latter being a known risk factor for type 2 diabetes, high blood pressure and cardiovascular disease. However, the majority of these studies have only included singleton births in their study populations. As twins are often born preterm and with a low birth weight it is reasonable to consider them a sub-group of all children born preterm, SGA and/or with a low birth weight.

In this thesis the aim was to investigate whether twins and singletons, born with non-optimal birth characteristics, exhibited different patterns with respect to reproductive pattern and general morbidity. For this purpose we had access to the unique Swedish population registers that provided an excellent opportunity to investigate these outcomes. Using the personal identification number assigned to each person residing in Sweden it was possible to link data on birth, hospitalizations and socio-demographics.

BACKGROUND

In 2005 it was estimated that approximately 9.6% of all births around the world were preterm births. Europe had the lowest rate of preterm births, 6.2% and Africa had the highest rate of preterm births 11.9% closely followed by North America with a 10.6% preterm birth rate¹. Another study estimated the worldwide incidence of preterm birth to be 11.1% in 2010².

It is well known that both short- and long-term survival among humans is highly dependent on the gestational week in which they were born, but also on their birth weight and birth weight in relation to gestational age³⁻⁵. In addition, restricted fetal growth is related to an elevated level of morbidity in general but also due to specific diagnoses. More specifically, children who are born preterm have an increased risk for hospitalizations due to physical conditions (e.g. type 2 diabetes, cardiovascular diseases, metabolic syndrome), mental conditions/diseases (e.g. psychiatric morbidity and drug abuse), and neurological disorders in adolescence and early adulthood⁶⁻⁸ (Figure 2).

Figure 2. Areas with long-term effect of preterm birth.

Prior research has shown that restricted fetal growth in general is related to increased likelihood of early childbearing (between the age of 13 and 27 years) in girls, whereas more prominent growth restriction is related to reduced likelihood⁹ (Figure 3).

Figure 3. *Consequences of non-optimal birth characteristics and future reproductive performance in young ages.*

It is possible that the short- and long-term effects of non-optimal birth characteristics are different for twins compared with singletons due to the difference in the etiology of preterm birth. Generally, the cause of preterm birth among twins is different than the cause of preterm birth among singletons. It has been quantified that, overall, somewhere between 30-40% of all preterm births are caused by maternal infections, while mechanical reasons such as increased uterine pressure and overdistention of the uterus, i.e. lack of space, are major causes of preterm birth among twins¹⁰⁻¹¹. In many cases twins are born preterm and with a low birth weight¹². Despite this most epidemiological studies on morbidity and reproductive patterns exclude twins and higher order pregnancies from their studies¹³⁻¹⁶. The reality today is that very few studies have focused on the potential effect twinning may have on reproductive pattern and general future health outcomes. Thus, there is a lack of knowledge on the risk for long-term morbidity associated with twin pregnancy and delivery when comparing twins with singletons.

Birth-characteristics in relation to outcomes in later stages of life

Twinning

Around the world the incidence of twin birth has increased since the mid 70's¹⁷⁻¹⁸. In Sweden alone the rate rose to 16.3 births/1000 women in 2003. However, since then the twin birth rate has stabilized around 14 births/1000 women¹⁹. Important factors explaining some of the increase in twin birth rates are IVF-treatment and hormone stimulation as is the increasing maternal age when giving birth^{18,20}.

The elevated risk associated with twin pregnancies is mainly due to the high number of preterm births and the increased proportion of children born growth restricted. Another important factor affecting the risk in twin pregnancies is delivery complications, although with the improved medical care of newborns born with non-optimal birth characteristics over the years this risk has declined some²¹⁻²².

Mortality

Both short- and long-term health and survival among children are highly dependent on the gestational week in which they were born, but also on size and weight in relation to gestational age³⁻⁵. Preterm birth and complications associated with preterm birth, such as respiratory distress, seizures, and infections are substantial contributors to neonatal and infant mortality. Close to 30% of all neonatal deaths can be attributed to preterm birth while 60-80% of all neonatal deaths are due to the infants being born with a low birth weight²³⁻²⁵. In a longer perspective, studies have reported slightly discrepant results. A study by Swamy et al reported that preterm birth was associated with a higher risk of mortality reaching into late childhood among both boys and girls¹³. This was not confirmed in a later study by Crump et al; they did however find that preterm birth was associated with an increased likelihood for mortality in early adulthood²⁶.

Long-term morbidity

In the past decades there has been an improvement in the medical treatment of children that are born preterm or with restricted fetal growth. This improvement has resulted not only in a much higher survival rate of these children, but also in a higher number of surviving children with both mental and physical chronic conditions. Restricted fetal growth and prematurity are related to an increased risk of morbidity in general but also due to several specific diagnoses. In particular, children born preterm have an increased risk for hospitalizations due to neurological sequelae, physical (congenital malformations, type 2 diabetes, cardiovascular diseases, metabolic syndrome), and mental conditions/diseases (psychiatric morbidity and drug abuse) in adolescence and early adulthood⁶⁻⁸. In a review by Räikkönen et al several long-term effects due to prematurity and SGA were listed on a number of areas including cognitive functioning, temperament and personality, mental disorders and increased sensitivity to stress²⁷ (Figure 2).

Physical morbidity

A large number of studies have proposed a connection between restricted fetal growth and an increased risk for major diseases later in life, e.g. the metabolic syndrome, which includes diabetes, high blood pressure, obesity, but also increases the risk for cardiovascular disease²⁸⁻³⁰. Different theories aimed at explaining the mechanisms behind these associations have been developed and perhaps the most debated of these is 'the fetal origins of adult disease hypothesis' (FOAD). The main concept of this theory is that "...an unfavorable development, or insults during fetal life, might induce lifetime effects on the subsequent development of body systems and hence give rise to major disease processes..."^{28,31}. This theory has received some criticism over the years both regarding its theory (too widely and vaguely formulated) and the methodology used i.e. missing important confounders, over-adjustment resulting in the "reversal paradox". This criticism has resulted in the

theory being slightly revised and in the development of an analytical scheme recommended when analyzing the FOAD related hypothesis³²⁻³⁸.

Previous research has also established that non-optimal birth characteristics, in addition to the well-established risks associated with pregnancy complications, are related to diseases of the nervous system, such as cerebral palsy as well as visual and hearing impairments³⁹⁻⁴⁰. The mechanisms by which these complications occur include intrauterine infections, central nervous system malformations, and reduced placental function⁴⁰. In addition, preterm birth and restricted fetal growth have been discovered to be related to respiratory disease and infectious diseases, and some evidence also exist of a relationship with malignancy⁴¹⁻⁴³.

Psychiatric morbidity and personality

Preterm birth and restricted fetal growth have recently been subject for research concerning mental illness. Low birth weight has been related to both psychiatric symptoms and psychiatric disorders, such as eating disorders, schizophrenia, and suicidal behavior⁴⁴⁻⁴⁶. Some evidence exist that high psychiatric morbidity increases between adolescence and young adulthood among children born SGA or with a low birth weight⁴⁷. An increased risk for attention deficit and hyperactivity disorder (ADHD) has also been noted among children born SGA and with a low birth weight⁴⁸. Similarly, an increased likelihood for autism spectrum disorders among children born preterm has been observed⁴⁹. This was partly validated in a within-twin study where it was concluded that low birth weight is associated with all forms of ADHD symptoms⁵⁰. The personality of those born with restricted fetal growth has also gained some attention. Some studies report that men and women born with very low birth weights display less negative emotions and are more cautious in late adolescence and young adulthood compared with controls^{41,51}, while others report an increased risk of delinquent, aggressive and externalizing problems⁵². Others on the other hand report a higher risk of internalizing symptoms, attention problems, and relational problems^{41,53}. In addition, preterm birth and low birth weight have

been studied with respect to future cognitive outcomes. Several studies provide evidence that future cognitive functioning is related to both prematurity⁵⁴ and low birth weight^{53,56}. It has also been suggested that these effects can extend all the way in to old age by causing "...a more rapid cognitive decline..." as the person becomes old²⁷.

Socio-economic and lifestyle factors' effect on future pregnancy outcome

Several socio-economic factors have been identified to influence the outcome of future pregnancies. Increasing maternal age when giving birth is a major factor influencing not only the likelihood of twinning (and higher order pregnancies) but is also a known predictor of children being born preterm as well as children born SGA. Other established factors related to having a child born preterm and/or a child born SGA include maternal educational level, unemployment, lack of social support and the mother's marital status⁵⁶⁻⁵⁸. Moreover, the mother's BMI and her smoking habits during pregnancy have been identified as risk factors for preterm birth and having a child with low birth weight⁵⁹⁻⁶⁰ as do the familial socio-economic position, during both childhood and adulthood, income, and childhood health and environment⁵⁹⁻⁶⁰.

Childbearing and pregnancy outcome

Hack et al⁴¹ have proposed that differences in personality and/or behavior may be involved in explaining the reduced childbearing among women born with restricted fetal growth. One of these potential factors influencing the future likelihood of becoming a parent is the onset of romantic relationships. Studies have indicated that children born preterm were less likely to be in a romantic relationship in early adulthood^{63,64}, or ever having been married or in a registered relationships⁶⁵, while children with a low/very low birth weight had a decreased likelihood of having experienced sexual relationships⁶⁶. This may in turn be caused by the reduced likelihood of very low birth weight children to leave the parental home⁶⁶.

It is also well known that restricted fetal growth, including low birth weight and SGA, may have an impact on organ structure and functioning, including the reproductive organs,^{8,28-29}, and therefore may be connected to future fertility and reproductive pattern. Studies have indicated that women born SGA may have a reduced ovarian and uterus size but also a reduced ovulation rate⁶⁷. Meanwhile, intra-uterine growth restricted men are at an increased risk for hypospadias, cryptorchidism^{x68} and men born preterm have an increased risk for testicular cancer⁶⁸⁻⁶⁹. Previous research on Swedish population data have found that preterm birth and low/very low birth weight were associated with a decreased likelihood of becoming a parent while being SGA did not decrease the likelihood of becoming a parent⁷⁰. The latter finding was in disagreement with a previous study on partly the same data (smaller cohort with a shorter follow-up time) where it appeared as if women born SGA had an increased likelihood of becoming a mother⁹.

Heritability and Epigenetics

The heritability of non-optimal birth characteristics has been widely studied previously, though focusing on singletons only. In a previous study, on partly the same data as this thesis, on the heritability of preterm birth and intrauterine growth restriction among singletons, it was found that women born SGA were more likely to themselves give birth to a child born SGA or preterm⁷¹. A more recent study added knowledge on the heritability by showing that women born in a spontaneous preterm delivery were more likely to repeat this in their own pregnancies⁷². Studies have also indicated that mothers giving birth to preterm twins are, in subsequent deliveries, more likely to give birth to preterm singletons⁷³⁻⁷⁴.

It has further been shown that genetic and environmental factors are equally important factors affecting birth characteristics of the newborn in terms of birth weight, prematurity and size for gestational age⁷⁵⁻⁷⁶. Several studies have also focused on explaining the mechanisms that have been hypothesized and shown to

cause the alteration of how certain genes are expressed and thereby affecting the risk for preterm birth⁷⁷⁻⁸³.

AIMS OF THE PRESENT THESIS

As most studies on reproductive pattern and its outcomes, morbidity and heritability of non-optimal birth characteristics are limited to the inclusion of only singleton births it is of importance to elucidate whether twins and singletons differ on these outcomes.

Hence, the principal aim of this research was to investigate the impact of non-optimal birth characteristics, i.e. preterm birth and SGA, on somatic health and reproductive pattern among adolescent and young adult men and women born as either twins or singletons. To this end, four separate sets of questions were formulated, investigated, and answered in the course of research leading up to the present thesis:

Firstly, if twins and singletons exhibit different reproductive patterns and if potential differences can be explained by their non-optimal birth characteristics? (Study I)

Secondly, having survived till the age of 13, is the future morbidity among twins and singletons different and is the morbidity affected differently due to non-optimal birth characteristics? (Study II)

Thirdly, does an intergenerational effect of non-optimal birth characteristics, as measured by preterm birth and SGA exist among twins and singletons and are twins and singletons affected differently? (Study III)

Fourthly and lastly, how does family size affect future reproductive pattern among twins and singletons? Do twins and singletons differ in the likelihood of becoming 1st and 2nd time parents? (Study IV)

MATERIAL AND METHODS

Retrieval of data

In Sweden it is possible to obtain unique data on a personal level due to the long tradition of registrations but also due to the unique personal identification number assigned to every person residing in Sweden, enabling linking information from different registers. Prior to retrieving any data several steps have to be taken. First, to be able to request data from any of the population-based registers an ethical approval has to be obtained from the Human Research Ethics Committee. After approval from the Human Research Ethics Committee a request to each agency responsible for the registers from which data are to be retrieved. One should bear in mind that an ethical approval does not imply that the agencies will be able to deliver the requested data, the legal expertise at each agency has first to approve of the data request making sure that delivery of the data are in line with current ethical principles and Swedish legislation. However, once data have been delivered, usually in the form of several data files, the data have to be set up into a manageable database. In setting up a database, comprising three generations, where data have been retrieved from several population-based registers, the data files have to be carefully merged. In addition, upon receiving the data a careful investigation must be undertaken to ensure that delivered data are indeed the desired data. Delivered data have further to be scrutinized meticulously for inconsistencies. This includes, e.g. crosschecking the different variables that could be used to identify twinning, birth weight and gestational age as well as birth weight with respect to gestational age. This also entails making sure that predefined variables in the data define what is to be investigated. If not, re-coding of these variables has to be performed. An example of this is that the Swedish Board of Health and Welfare changed the definition of parity. Previously it was defined as number of deliveries, where the delivery of multiples counted as one. This was later changed in to number of children delivered (i.e. if a primiparous woman delivered a pair of twins the first child had parity=1 and second child had parity=2).

International Classification of Diseases

The first version of the International Classification of Diseases (ICD) was published in 1949. Over the years the ICD has been through a couple of revisions and for the purpose of this thesis the three latest versions have been used. ICD-10 was released in 1994 and the use of this release in Sweden started in 1996, replacing the previous version, ICD-9, which had been in use since 1987 when ICD-8 was replaced (in use since 1969). In order to obtain definitions of diseases that were comparable over the years the classifications of ICD-10 had to be converted to the classifications in ICD-9. ICD-9 was chosen as baseline since the individuals in study I-III were born in 1973 and 1983 but also since the aim was to study morbidity in late childhood and early adolescence. The conversion of ICD-10 to ICD-9 is presented in Table 1.

Diagnosis codes for each of the different ICD version used in this thesis		
ICD-9 Text	ICD-9	ICD-10
Infections and parasitic diseases (1)	001-139	A00-B99
Neoplasm (2)	140-239	C00-D48
Endocrine, nutritional and metabolic diseases and immunity disorders (3)	240-279	E00-E90 D80-D99
Blood diseases (4)	280-289	D50-D77
Mental disorders (5)	290-319	F00-F99
Diseases of the nervous system and sense organs (6)	320-389	G00-H95
Diseases of the circulatory system (7)	390-459	I00-I99
Diseases of the respiratory system (8)	460-519	J00-J99
Diseases of the digestive system (9)	520-579	K00-K93
Diseases of the genitourinary system (10)	580-589	N00-N99
Diseases of the skin and subcutaneous tissue (12)	680-686	L00-L99
Diseases of the musculoskeletal system and connective tissue (13)	710-739	M00-M99
Congenital anomalies (14)	740-759	Q00-Q99
Symptoms, signs and ill-defined conditions (16)	780-789	R00-R99
External causes of morbidity and mortality (17,18)	800-999	S00-T98
	E807-E999	V01-X59

Table 1. Conversion of ICD-9 to ICD-10.

Data sources

The Swedish Medical Birth Register

The Swedish Medical Birth Register (MBR) was introduced in 1973. From the start information regarding prenatal, obstetrical, delivery and neonatal factors was based on “Medical Birth Reports” (Medicinska Födelsemeddelande, MFM). However, some inconsistencies in reporting were present and after an evaluation in 1976 a revised procedure for data collection was implemented in 1982 to minimize reporting errors. At the same time, it was decided to include information pertaining to diseases during pregnancy. Since then the information collected has been through additional changes in 1990, 1994 and 1998. Today MBR contains information on prenatal care (age, maternal height and weight, nicotine use, previous pregnancies and diagnoses prior and during pregnancy), delivery (mode of delivery, fetal presentation, pain relief used during labor and delivery) and neonatal care (birth weight, body length, gestational age and newborns’ diagnoses) as well as maternal chronic diseases⁸⁴. The Swedish Medical Birth Register has a high coverage rate with only 0.5-3.0% missing records⁸⁴⁻⁸⁵.

The National Patient Register

The National Patient Register, previously known as Hospital Discharge Register, was established in 1964. From the start it mainly included information on psychiatric care. In 1984 it was decided that all inpatient care should be reported to the register and this was implemented in 1987. The register was further revised in 2001, when all outpatient visits (including outpatient surgery and psychiatric care from both private and public caregivers) were included in the mandatory reporting to the register. As of today the register contains information on e.g. sex, age, date of admission and discharge, main and secondary diagnoses⁸⁶⁻⁸⁷. The register has a high total coverage rate of about 99%, and in a recent validation of the register it was concluded that for the most part the diagnoses in the register are of a high quality with an the overall positive predictive value (PPV) in the range 85-95%.

The Cause of Death Register

The Cause of Death Register, in its current form, was established in 1961 and contains information regarding the cause of death for all persons residing in Sweden. This means that cases where the death occurred abroad are also included in the register. The register does not include infants who died at birth. Starting from 2012 all deaths occurring in Sweden are registered in the Cause of Death register⁸⁸⁻⁸⁹, implying that those, at time of death, who currently does not reside in Sweden but who dies while in Sweden are also included. The coverage rate is close to 100% but approximately 2.5% of the entries are incomplete.

The Total Population Register

The Total Population Register (TPR) was established in 1968 and contains information regarding a person's citizenship, country of birth, marital status, migration (both immigration and emigration) and deaths and births⁹⁰⁻⁹¹. The register is of good quality, though some over-coverage exists due to missing reports of deaths and emigrations (i.e. people are registered though they should not be).

The Multi-Generation Register

The Multi-Generation Register, which is based on the TPR, includes all persons who were born in Sweden from 1932 and onward, but also include persons who, at some point, have been registered in Sweden since 1961. The register is used to identify the biological parents or adoptive parents of these "indexpersons"⁹²⁻⁹³.

The Education Register and the 1970 Population and Housing Census

Prior to 1985 when the Swedish Register of Education was established, data on level of education were retrieved from the Population and Housing Census that took place in 1970. Today the register is updated regularly and contains information on highest level of education, year of completion and some basic demographic data such as age and sex. The register is limited to individuals between 16 and 74 years of age⁹⁴⁻⁹⁶.

The study population studies I-III

The study population in studies I-III was defined as all men and women born in Sweden between 1973 and 1983 who were alive and living in Sweden at the age of 13.

The starting population, consisting of all men and women born between 1973 and 1983, endpoints included, included a total of 1,070,380 individuals. However, individuals who were deceased before their 13th birthday, individuals who did not reside in Sweden at the age of 13, individuals with missing values on birth weight and/or gestational length and individuals who had an unlikely birth weight with respect to the gestational age (Figure 4) were removed from the study population. The limits for removal due to extremely high birth weight with respect to gestational age were defined as:

- > = 2000 grams for gestational weeks, < =28,
- > = 2500 grams for gestational weeks 29 and 30,
- > = 3000 grams for gestational weeks 31 and 32,
- > = 3500 grams for gestational weeks 33 and 34,
- birth weights for other gestational weeks were within limit.

Limits for removal due to extremely low birth-weights with respect to gestational age were defined as:

- <= 400 grams in gestational week 29,
- <= 800 grams in gestational week 30,
- <=1000 grams in gestational week 31,
- <= 1150 grams in gestational week 32,
- <= 1250 grams in gestational week 33,
- <= 1450 grams in gestational week 34,
- <= 1600 grams in gestational week 35,
- <= 1700 grams in gestational week 36,
- <= 1800 grams in gestational week 37,
- <= 1950 grams in gestational week 39,

<= 2000 grams in gestational weeks 40, 41 and 42,

<= 2500 grams in gestational weeks 43 and 44.

After removal of individuals due to these events, the final study population consisted of 1,016,908 individual of which 16,561 were identified as twins.

Figure 4. Flowchart of the exclusion of study participants in studies I-III.

The study population study IV

The study population in study IV was defined as all men and women born in Sweden between 1973 and 1993 who were alive and living in Sweden at the age of 13.

The starting population, consisting of all men and women born between 1973 and 1993, endpoints included, encompassed a total of 2,148,960 individuals. However, after removal of individuals who had deceased before their 13th birthday, individuals who did not reside in Sweden at the age of 13, individuals with missing values on birth weight and/or gestational length and finally removal of individuals who had an unlikely birth weight with respect to the gestational age (Figure 5), the final study population consisted of 2,091,205 individual of which 39,726 were identified as twins.

The limits for removal due to an extremely high birth weight or extremely low birth-weights were defined as for studies I-III.

Figure 5. Flowchart of the exclusion of study participants in study IV.

Design of the studies

Study I

In this study the primary focus was to investigate the effect of twinning on future reproductive performance and to investigate if the reproductive performance among both twins and singletons were affected by non-optimal birth characteristics in the same way. For this purpose all twins and singletons born between 1973 and 1983 in Sweden and who were alive and residing in Sweden at age 13 were included and followed until the end of 2006, for more detailed description see Figure 4. Data on each study subject’s own birth as well as the birth of their first offspring, and the study object’s parental socio-demographic factors were collected from Swedish population based registers, Figure 6. Stratified analysis on gender where hazard ratios of the likelihood of becoming a parent and corresponding 95% CI were calculated using Cox proportional hazards model. Two sets of models were considered, in the first model estimates were adjusted for socio-demographic factors and in the second model estimates were further adjusted for the men’s and women’s own birth characteristics.

Figure 6. Design of study I.

Study II

In this national population-based register study, all individuals born between 1973 and 1983 who were alive and living in Sweden at 13 years of age constituted the sample. The purpose of this study was to examine the morbidity, where hospitalization was used as a proxy for morbidity, among both twins and singletons and to evaluate if the outcome differed depending on twinning status and to what extent non-optimal birth characteristics affected the morbidity. Data on each study subject’s own birth, parental socio-demographic factors as well as data on hospitalizations were collected from Swedish population based registers, Figure 7. Odds ratios of the likelihood of being hospitalized and corresponding 95% CI were calculated using logistic regression. Several different outcomes were evaluated, general hospitalization due to any cause but also due to each the groups of diagnosis present in each of the main chapters in ICD-9. Moreover, due to an intra-class correlation among twins all confidence intervals were adjusted accordingly⁹⁷.

Figure 7. Design of study II.

Study III

This is a population-based register study of mother-firstborn offspring pairs recorded in the Swedish Medical Birth Register. The study included 268,867 mothers born in 1973-1983 and their firstborns born in 1986-2009. The main focus of this study was to investigate the intergenerational effect of preterm birth and SGA among both twins and singletons and try to elucidate if any differences in the intergenerational effect between twins and singletons were present. Socio-demographic data on grandmothers as well as mothers, birth characteristics of the mothers and their offspring were collected from validated Swedish registers, Figure 8. Multiple logistic regression models were performed to estimate the likelihood of giving birth to a child born preterm or SGA. In these models adjustment were made for both grandmothers’ and mothers’ socio-economic background but also for the mothers’ own birth characteristics.

Figure 8. Design of study III.

Study IV

This is a register-based study where the study population consisted of all men and women born in Sweden between 1973 and 1993 who were alive and living in Sweden at 13 years of age. Data on the study objects’ own births as well as their offspring, parental socio-demographic factors were collected from Swedish population based registers, Figure 9. Hazard ratios for the likelihood of becoming a parent were estimated using Cox’s proportion hazard models. All models were adjusted for socio-demographic factors as well as birth characteristics and a categorized indicator on the number of siblings, approximated by the number of biological brothers and sisters with the same mother.

Figure 9. Design of study IV.

Statistical methods

Logistic Regression

Binary logistic regression can, if simplified, be separated into two model types - "single logistic regression" and "multiple logistic regression". The first model type is limited to one nominal (e.g. yes/no) dependent variable and one independent variable that can be either categorical or continuous and is used when one wants to predict the probability of the outcome. The second model type pertains to models with two or more independent variables. When performing logistic regression it is assumed that the observations are independent and that the relationship between the natural log of odds ratio and the independent variable is linear. The logistic regression model predicts the odds ratio of the studied event happening and is relevant when studying the cumulative incidence of an event (i.e. proportion of new events occurring within a pre-specified time period). A short description of how the odds ratio is calculated is provided in Figure 10.

In studies II and study III this was the chosen method of analysis in order to test each studies hypothesis. In study II multiple logistic regression was used to calculate the odds ratio for hospitalization due to the main chapters in ICD-9 and total hospitalization, comparing twins with singletons. Each main chapter was modeled separately. We also considered models where data were stratified by twinning status but also according to optimality at birth (non-optimal vs. optimal birth characteristics).

In study III multiple logistic regression was used to estimate the odds ratio of giving birth to a child that was either born preterm or SGA. The analyses were stratified upon whether the mother was born preterm or SGA.

		EVENT	
		YES	NO
EXPOSURE	YES	A	B
	NO	C	D

OR = $\frac{A/C}{B/D}$ = proportion of events among those exposed compared with those not exposed

B/D = proportion of non-events among those exposed compared with those not exposed

Figure 10. Calculation of odds ratio.

Cox's proportional hazards model

In short, Cox proportional hazards model is a time-to-event model strategy in which the relative risk for an event happening at a specific time is estimated (hazard ratio). As in logistic regression models it is possible to have more than one independent variable in the model. Moreover, when analyzing data through Cox proportional hazards models it is possible to allow censoring of the study subjects. This means that study subject can exit from the model prior to the studied event happening, or enter the study at different time-points and estimates of the hazard will take this into account as displayed in Figure 11.

Figure 11. Alternatives on when to enter and exit the Cox proportional hazards model.

In Cox proportional hazards model, it is assumed that the hazard ratio (e.g. between twins and singletons) remains constant over time. If this is substantially violated it is possible to model data with time dependency or to use separate models in different time periods of follow-up.

In study I and study IV, Cox proportional hazards model was used to estimate the likelihood of becoming a parent. The time variable was defined as age, and

individuals exited from the models when they became parents, emigrated, died or reached the end of follow-up, depending on which of these events was the first event to occur.

Intra-pair correlation due to twinning

Due to a correlation of the incidence of hospitalization and 1st and 2nd twin, the confidence intervals for the OR's in study II were adjusted using cluster design effect formulas⁹⁷.

Analysis with respect to familial factors

In analyzing the effect birth characteristics have on future reproduction, pregnancy outcome and morbidity it is important to also acknowledge familial factors affecting the outcome. Examples of such factors include family size, social position (employment, income, education, social support network, place of residence), dietary intake, and the general environment into which the child is born. Due to the nature of the registers we were not able to obtain data on all of these factors. In study IV, we used number of maternal full-siblings as a proxy for family size and in all studies information on maternal and paternal level of education were included in all studies as were parental region of origin. Also, in studying the pregnancy outcome of the index persons' own pregnancies BMI and maternal nicotine use were included.

Exposure, outcomes and covariates

Primary exposures

Gestational age at birth

Gestational age at birth was separated into four categories: < 32 completed gestational weeks (very preterm), 32-36 completed gestational weeks (moderately preterm), 37-42 completed gestational weeks and lastly >42 completed gestational weeks (post-term), this definition of gestational age was used in all studies. The persons in this study were born between 1973 and 1983 in studies I-III and 1973 and 1993 in study IV. During this time period the gestational length has mainly been

measured by the last menstrual period but also with the assistance of ultra sonogram. These two ways to determine the gestational age may yield slightly different estimates since estimates using the menstrual period usually are 2-3 days higher than the estimates from an ultra sonogram^{84,98}.

Birth weight

Birth weight were split in to three categories: very low birth weight was defined as a birth weight < 1500 grams, low birth weight between 1500 grams and 2499 grams, and normal birth weight ≥ 2500 grams (used in all studies).

Small for gestational age

For the purpose of this thesis small for gestational age was defined as a birth weight < -2 SD of the mean weight for the gestational length⁹⁹ according to the Swedish standard and currently used by the National Board of Health and Welfare.

Non-optimal birth characteristics

Non-optimal birth characteristics, used in study II, was defined as being either SGA or born preterm while optimal birth characteristics were defined as being born at term and appropriate for gestational age (AGA).

Twinning

Twinning is a key exposure in this thesis and was used in all studies. Twinning was dichotomized into two levels; singletons and twins. All higher order pregnancies were excluded from the studies due to a very limited number (n=220 in studies I-III and n=346 in study IV).

Primary outcomes

Likelihood of giving birth

In studies I and IV the outcomes studied were the likelihood of giving birth, estimated using Cox proportional hazards model. In study I the individuals were followed till the end of 2006 thus among those that we could follow through all years the endpoint was anywhere between 23 and 33 years of age. In study IV the endpoints were 29 and 39 years of age. This implies that most of the participants only had begun reproducing; mean age in Sweden when women had their first child was 28.5 years in 2013¹⁰⁰. Only a limited number of participants were approaching the end of their reproductive years.

Likelihood of being hospitalized

In study II we wanted to investigate the morbidity in adolescence and young adulthood. As a proxy for morbidity we analyzed the likelihood of being hospitalized according to the main chapters in ICD-9 and total (all-cause) hospitalization. Total number of hospitalizations was dichotomized into two levels, <6 and ≥6 hospitalizations.

Likelihood of giving birth to a child born preterm or SGA

The primary aim in study III was to study the intergenerational effect of preterm birth and SGA, subsequently the outcomes studied were the likelihood of giving birth to a child born preterm or SGA.

Covariates/confounders

Several different covariates were considered in this thesis and can be divided into two main groups. The first group consists of socio-demographic factors related to the childhood of the study population and included parental level of education divided into “elementary school” (9 to 10 years of schooling), “high school” (11-13 years of schooling) and “graduate/postgraduate education” (≥ 14 years of schooling) and since there was a great number of missing values on this variable a separate category labeled “unknown” was included. The parents’ origin was categorized into “both parents from Nordic countries” and “one or both parents from non-Nordic countries”. Finally, factors only related to the mothers of the study population were included; mother’s marital status when giving birth was divided into “married”, “unmarried” and “divorced or widowed” while mother’s age when giving birth were divided into four categories “13-19 years of age”, “20-26 years of age”, “27-33 years of age” and “34 years of age or older”. Mother’s parity was divided into “primiparous” and “multiparous”.

The second group of variables relates to the study population when they themselves became parents and included, age when giving birth, level of education, and marital status. All of these variables were coded as for the mothers of the study objects except age when giving birth which was divided into “ ≤ 19 years of age”, “20-24 years of age” and “ ≥ 25 years of age”. In addition, for the study population we had access to information on the women’s BMI in early pregnancy and nicotine use prior and during pregnancy. Women’s BMI was divided into four categories “ <18.5 ”, “18.5-24.9”, “25.0-29.9” and “ ≥ 30.0 ” according to WHO classification of BMI¹⁰¹ Nicotine use was defined as using either cigarettes or snuff, both before and during pregnancy, and was categorized into “yes” and “no”.

In study IV we also used a family size variable. As a proxy for family size we used the number of siblings with the same mother, and this was further categorized into, “0-2” and “ ≥ 3 ” number of siblings.

ETHICAL CONSIDERATIONS

Population-based registers are widely used in medical research today. In Sweden Statistics Sweden and the National Board of Health and Welfare frequently supply researchers with data, either from just one agency or, which is often the case, merged information on individuals from both agencies.

For the agencies to consider releasing data to the researcher an ethical approval has to be obtained from the Human Research Ethics Committee. However, an approval does not guarantee that the researcher will obtain any data. It is up to each agency to establish that current ethical principles and Swedish legislation will not be violated prior to delivering any data to the researcher, thus it is possible to be denied access to data. According to current legislation in Sweden the Regional Ethical Review Committee has the right to make a judgment based on balancing the individual's right to have his/her integrity inviolate, against the societal benefits from a specific research project. This means that the individuals in the registers have no possibility to either approve or disprove of the usage of their information.

The registers of today provide the researchers with invaluable research opportunities, and if consents were to be collected from each and every person in the registers much research would not be performed, and hence important knowledge would be missed. However, is it ethically viable to include persons in a research project, based on population-based registers, without their consent? In most cases it may be assumed that the public benefits outweigh the integrity problem. In most cases this does not cause a problem since data are anonymized and the results are reported on a group level, not on an individual level. However, in some cases when the focus is on rare outcomes the risk for identifying individuals increases. It is the researcher's responsibility to assure that the personal integrity is not violated, and one way of accomplishing this is to publish data on group levels only.

Another ethical consideration when working with large population-based studies is the very large sample size. Having a large sample size will often yield a statistical significance even when the differences are very small which may be without clinical importance. This requires the researcher to be aware of the implication of extremely large sample sizes and interpret the findings with caution. A statistical significance does not always equal clinical significance.

The studies included in this thesis were approved by the Regional Ethical Review Board in Linköping 03-556, 07-M66 08 – 08-M 233-8.

RESULTS

Study I. Likelihood of giving birth among twins and singletons with respect to birth characteristics

A total of 1,016,688 men and women were included in this study of which 16,561 (1.6%) were born as twins, Table 1. Among both twins and singletons the proportion of males and females was equal (50.1% were men and 49.9% were women among twins and 51.4% were men and 48.6% were women among singletons). During the follow-up, which ended in 2006, 40% of the twin females and 24% of the twin males had become parents. Corresponding percentages for singletons were 44% among females and 26% among males.

	Men	Women	Total
Singleton	513,801 (98.4%)	486,326 (98.3%)	1,000,037 (98.4%)
Twin	8,313 (1.6%)	8,248 (1.7%)	16,561 (1.6%)
Total	522,114 (51.4%)	494,574 (48.6%)	1,016,688 (100.0%)

Table 1. Study population and its distribution of twinning and sex.

The primary finding in this study was that twins in general had a reduced likelihood of reproducing compared with singletons, HR=0.87 (95% CI=0.84-0.90) among females and HR=0.89 (95% CI=0.85-0.94) among males. Moreover comparisons on the likelihood of reproducing among twins vs. singletons were further elaborated using sub-group analyses where the sub-groups were defined by birth characteristics. It was found that for a number of birth characteristics twins had a reduced likelihood of reproducing compared with singletons. Among females, twins born AGA, had a normal to low birth weight, born at term or moderately preterm had a lower likelihood of reproducing compared with singletons. Among males, twins born AGA, with a lower birth weight, born moderately preterm had a decreased likelihood of reproducing, Table 2.

	Females HR (95% CI)	Males HR (95% CI)
Size for gestational age		
Appropriate for gestational age	0.89 (0.85-0.93)	0.93 (0.88-0.99)
Small for gestational age	0.96 (0.88-1.03)	1.09 (0.97-1.23)
Large for gestational age	1.22 (0.74-2.02)	0.78 (0.36-1.69)
Birth weight		
Normal birth weight (≥2500 g)	0.90 (0.86-0.94)	0.92 (0.87-0.98)
Low birth weight (1500-2499 g)	0.92 (0.86-0.98)	1.04 (0.95-1.15)
Very low birth weight (<1500 g)	1.10 (0.83-1.46)	0.79 (0.51-1.24)
Gestational age		
Term (37-42 weeks)	0.92 (0.88-0.96)	0.97 (0.91-1.03)
Preterm (32-36 weeks)	0.86 (0.79-0.92)	0.90 (0.81-0.99)
Very preterm (<32 weeks)	1.29 (1.02-1.63)	1.33 (1.00-1.77)
Post-term (>42 weeks)	0.70 (0.44-1.12)	1.03 (0.57-1.88)
Optimality at birth		
Optimal	0.90 (0.86-0.93)	0.92 (0.87-0.97)
Non-optimal	0.91 (0.86-0.96)	0.99 (0.92-1.07)

Table 2. Hazard ratio of becoming a parent among females and males in sub-groups defined by birth characteristics, comparing twins vs. singletons.

Study II Hospitalization in adolescence and young adulthood among twins and singletons

The study population in this study was the same as for study I, see Table I for distribution of twinning and gender.

The main aim of this study was to study the morbidity, where hospitalization was used as proxy variable, among twins and singletons. For this purpose we studied the main chapters of the Swedish version of ICD-9 in order to capture general morbidity. Twinning was, in general, associated with a slightly increased likelihood of being hospitalized, (Total hospitalization, OR=1.04 95% CI=1.00-1.07; Hospitalized ≥ 6 times, OR=1.17 95% CI=1.09-1.25) compared with singletons. They were also at an

increased risk for hospitalization due to Infections and parasitic diseases, Diseases of the nervous system and sense organs, Diseases of the digestive system, Congenital anomalies and External causes of morbidity and mortality (see Study II for details). When stratifying data by optimality at birth, many of the relationships were reversed indicating that twins seem to fare somewhat better when accounting for birth characteristics (Table 3).

Hospitalization diagnosis according to ICD-9	Optimal twin vs. optimal singletons	Non-optimal twin vs. non-optimal singleton
Endocrine, nutritional and metabolic diseases and immune disorders	1.03 (0.86-1.23)	0.82 (0.69-0.97)
Blood diseases	0.99 (0.80-1.23)	0.70 (0.55-0.89)
Mental disorders	0.83 (0.74-0.94)	0.88 (0.79-0.98)
Diseases of the respiratory system	0.87 (0.82-0.93)	0.87 (0.82-0.93)
Diseases of the genitourinary system	0.92 (0.83-1.02)	0.82 (0.74-0.91)
Diseases of the musculoskeletal system and connective tissue	0.97 (0.87-1.09)	0.88 (0.79-0.99)
Symptoms, signs and ill defined conditions	0.87 (0.81-0.93)	0.81 (0.76-0.87)
External causes of morbidity and mortality	1.07 (1.02-1.13)	1.02 (0.97-1.07)
Total (any hospitalization)	0.96 (0.92-1.01)	0.90 (0.86-0.95)

Table 3. Likelihood of being hospitalized stratified by optimality at birth.

However, also adjusting for birth characteristics non-optimal twins had a reduced likelihood of being hospitalized due to congenital anomalies (OR=0.86 95% CI=0.77-0.96) but also exhibited a lower likelihood of being hospitalized more than 5 times (OR=0.86 95% CI=0.78-0.94) compared with non-optimal singletons.

Study III Intergenerational effect of preterm birth and small-for-gestational-age

In study III 268,867 women born between 1973 and 1983 and their firstborn offspring, born between 1986 and 2009, were included. Among these women 4,073

(1.5%) were born as twins while 264,794 (98.5%) were born as singletons. Analyzing the intergenerational effect of SGA it was found that women born SGA were more likely to give birth to a SGA child among both twins and singletons while an increased likelihood of giving birth to a preterm child was only present among mothers born as singletons. If the mother herself was born preterm singleton she was more likely to give birth to preterm child. However, no statistical differences were present in the difference between twins and singletons even though the parameter estimates, for most estimates, were lower among twins compared with singletons. As this could be caused by a low power of the study, due to a relatively low number of twins included, linear regression models were also considered. In these models, a statistically significant interaction term between twinning and mothers' size for gestational age was present, indicating that singleton mothers born SGA had a higher likelihood of having a child with low birth weight compared with twin mothers born SGA, Table 4.

Logistic regression	Mother Singleton OR (95% CI)	Mother Twin OR (95% CI)	Twin vs. Singleton
Child preterm			
Mother preterm	1.40 (1.28-1.52)	1.11 (0.80-1.55)	0.191
Mother SGA	1.35 (1.25-1.46)	1.13 (0.80-1.60)	0.380
Child SGA			
Mother preterm	1.11 (0.97-1.28)	1.39 (0.90-2.14)	0.265
Mother SGA	2.96 (2.71-3.24)	2.15 (1.40-3.31)	0.232
Linear regression	β (95% CI)	β (95% CI)	Twin vs. Singleton
Child birth weight			
Mother preterm	-53.88 (-65.72 - -42.04)	-58.56 (-96.77 - -20.34)	0.296
Mother SGA	-255.79 (-237.10 - -214.47)	-116.41 (-157.18 - -75.65)	<0.001
Child gestational age			
Mother preterm	-0.28 (-0.32 - -0.24)	-0.22 (-0.35 - -0.09)	0.068
Mother SGA	-0.20 (-0.24 - -0.16)	-0.06 (-0.19 - -0.08)	0.061

Table 4. *Likelihood of giving birth to a child born preterm or SGA, stratified by twinning status.*

Study IV Reproductive pattern among twins and singletons in relation to number of siblings

The study population for study IV was an extended cohort compared with studies I-III and with a longer follow-up. The study population for this study was all men and women born between 1973 and 1993 and they were followed till the end of 2012. In total 2,091,205 men and women were included of which 39,726 were born as twins. In Table 5 the distribution of twinning and sex is presented.

	Men	Women	Total
Singleton	1,053,819 (98.1%)	997,660 (98.1%)	2,051,479 (98.1%)
Twin	20,014 (1.9%)	19,712 (1.9%)	39,726 (1.9%)
Total	1,073,833 (100.0%)	1,017,372 (100.0%)	2,091,205 (100.0%)

Table 5. Study population and its distribution of twinning and sex.

In this study it was found that twins had a lower likelihood of reproducing (men, HR=0.96 95% CI=0.93-0.99; women, HR=0.90 95% CI=0.88-0.93) compared with singletons. No difference was detected in the likelihood of having a second child among women, however among men the likelihood of having a second child was larger among twins compared with singletons (HR=1.07, 95% CI=1.02-1.11). Including the dichotomous variable on the number of siblings, 0-2 siblings and >2 siblings the likelihood of reproducing was higher, among both male and female twins and singletons, if the person had more than two siblings. This effect remained when analyzing the likelihood of having a second child.

	Male		Female	
	Twins HR (95% CI)	Singletons HR (95% CI)	Twins HR (95% CI)	Singletons HR (95% CI)
Number of siblings	HR for becoming a first-time parent			
0-2	Reference	Reference	Reference	Reference
>2	1.17 (1.08-1.27)	1.16 (1.15-1.18)	1.18 (1.10-1.26)	1.22 (1.21-1.24)
Number of siblings	HR for becoming parent to a second child			
0-2	Reference	Reference	Reference	Reference
>2	1.13 (1.01-1.26)	1.13 (1.12-1.15)	1.16 (1.06-1.27)	1.19 (1.18-1.21)

Table 6. Hazard ratio for becoming a parent in relation to number of siblings, stratified by twinning and sex.

DISCUSSION

Summary of principal findings

- It was found that twins, both males and females, in general have a lower likelihood of reproducing in ages 13-33 compared with singletons. The diverging birth characteristics among twins and singletons fail to fully explain this difference.
- Overall, twins had a higher likelihood of being hospitalized in adolescence and early adulthood compared with singletons. However, even though twin pregnancies increases the risk for the newborn to be born with a low birth weight, preterm or SGA it appears as twins actually had a lower risk for being hospitalized due to several diagnoses when taking their diverging birth characteristic into account.
- An intergenerational effect of preterm birth and SGA births was present among both twins and singletons. Mothers born as singletons and preterm or SGA had an increased likelihood of giving birth to a child that was either preterm or SGA. The intergenerational effect was most pronounced among women born a singleton SGA who had a 3 times higher hazard rate of giving birth to a child SGA compared with singleton women not born SGA, which also was the only risk present among twins.
- Twin males and females have a lower likelihood of becoming a first-time parent compared with singletons. These differences vanished or were reversed when studying the chance of becoming a parent again. The likelihood of male twins becoming a second-time parent was higher compared with singletons while there was no difference between female twins and singletons.

- The number of siblings was found to be an important factor when studying future reproductive performance among both twins and singletons. Having three or more siblings was associated with an increased likelihood of becoming parents among male and female twins and singletons alike.

Methodological considerations

As with all research, the designs of these studies have their advantages and disadvantages. A major strength of all studies in this thesis is that data are prospectively collected from large population-based registers yielding a large study population. This may also impose a problem, as even very small differences will become statistically significant.

In using population-based register data it is important to be aware of the quality of the data and the possible effects it could have on the research. Even though all registers used have been validated and are of good quality some flaws still exist that may affect the results despite our efforts to minimize these risks. The first consideration pertains to the birth characteristics of the study population. In the first three studies the study population comprised men and women born between 1973 and 1983, and in the last study the cohort was extended to include men and women born between 1973 and 1993. It is possible that intrauterine growth curves provided by Marsal et al in 1996⁹⁹ to calculate SGA and LGA are inappropriate for those born prior to 1996 as the mean birth weight of children has increased over time. Other possible problems involve the estimation of gestational age. From 1973 to the mid 80's, when ultrasonogram was introduced in the estimation of gestational age, the last menstrual period was the method used. These two methods yield slightly different estimates on gestational age, as estimates using the last menstrual period are usually two to three days higher^{84,98}. Moreover, validation of the birth weights registered in MBR has shown that some measurement errors exist at both endpoints, i.e. among those with extremely low or extremely high birth weights. The majority of

these errors are simply due to errors when transferring data from paper to computer. In an attempt to alleviate this problem we excluded individuals with an extremely high or extremely low birth weight with respect to the gestational length.

The second consideration pertains to the reproductive performance of the study population. As of today it is not possible to obtain any information on whether the women had undergone an induced abortion or had experienced a spontaneous abortion. This could cause an underestimation of the general reproductive rate as our estimates are based on live born infants. However, there is no obvious reason to expect that either twins or singletons are more prone to have had induced abortions. Thus, this is not expected to cause any bias regarding the comparisons of reproductive pattern of twins and singletons. There was also a lack of information on whether the men and women had tried to conceive. Moreover, in the studies of reproductive pattern the follow-up time was limited with follow-up ending in 2009 in study I and 2012 in study IV. Hence, the oldest persons were 33 years old at the end of follow-up in study I, and 36 years old in study IV. In 2013 women were, on average, 28.5 years old when they gave birth to their first child and usually men are on average somewhat older than women when starting a family. Thus, many of the participants have just started their reproduction and have many reproductive years left.

The third consideration is related to twinning. Nationally and internationally the twinning rates are at a much higher today compared with some 20 years ago¹⁷⁻¹⁸. A possible explanation to this is the increasing maternal age at childbirth, but the introduction of IVF-treatment of couples having problems conceiving “naturally” could also be a contributing factor¹⁸. As all individuals in study I were born prior to the introduction of IVF-treatment this cannot affect the twinning rate, nor should hormonal treatment have a major effect on the twinning rate. It is likely, however, that some of their offspring is the result of a successful IVF-treatment. Moreover, we

did not have information on whether the twins were monozygotic or dizygotic. This was not considered to be a major problem as the research focused on reproductive pattern, hospitalization, and the intergenerational effect of non-optimality at birth indifferent of zygosity.

The fourth consideration is focused on the estimation of morbidity. For this work the presence of hospitalization according to any of the main chapters in ICD-9 was used. Since primary care is not part of the NPR we only managed to study the most severe cases were studied, i.e. those requiring treatment as an inpatient or outpatient at a hospital, of each main chapter.

The last consideration pertains to population-based register studies in general. Selecting a large cohort of individuals, potentially, a large number of statistically significant differences will be found. This means that even small differences, which are negligible or not have clinical relevance, will become statistically significant. It is thus very important to also consider the estimates (and confidence limits) of the associations when interpreting the findings.

Principal findings in relation to other research

In comparing the findings in this thesis with previous research presently available it is important to be aware of how the researches have defined their variables as e.g. SGA. Depending on study design and different standards the definition of SGA varies from study to study. In our studies we have used $<2SD$ of the mean birth weight depending on sex and gestational age. In one study analyzing the risk for thrombocytopenia among SGA infants SGA was defined as below the 10th percentile¹⁰²⁻¹⁰³ while in a study by Diaz et al in 2015 SGA was defined as $<2SD$ for gestational age¹⁰⁴ and in yet another study $<2.5SD$ for gestational age was used¹⁰⁵. Even minor differences in the definition of SGA birth are likely to affect the results and thus the comparability across studies.

In addition to the different definitions of SGA differences in the definition of study population are likely to be present – all depending on the availability of register data. In comparing studies and their findings it is important to take these differences into account, as is the design of the study. One should ask oneself several questions such as: Which study design was used? How was the data collected? How were the controls selected? Any matching, and if so which variables were used? How is migration and deaths accounted for in the follow-up? All of these factors will inevitably affect the comparability across studies.

Reproductive pattern

Several studies have provided evidence that birth characteristics is related to the future likelihood of giving birth^{9,70-71}, although the findings have not been conclusive. The studies in the present thesis found that women born SGA, or with a low/very low birth weight or preterm/very preterm had a decreased likelihood of giving birth but also very preterm males had a decreased likelihood of becoming a parent. These findings are in agreement with previous research on Swedish register data^{9,70-71}. Performed studies were also able to verify the findings of women born SGA being less likely in having a child, which is the conclusion drawn from a previous study on partly the same data but with a smaller cohort and where the follow-up time was shorter⁹.

In the present work it was further found that twins had a lower likelihood of reproducing compared with singletons. This could be due to a variety of causes. Twins often form relationships at a later stage in life¹⁰⁶⁻¹⁰⁷ and more often have a university degree¹⁰⁸. The first explanation may very well cause twins postponement in entering parenthood, while the latter explanation could be either way – twins more often pursue an academic education due to not being able to have children *or* the postponement is caused by their educational focus. Also, though the models have been adjusted for several socio-demographic factors, there is always the

possibility of additional important confounding factors not adjusted for that influence the estimates¹⁰⁹.

Hospitalization according to the main chapters in ICD-9

Twinning per se has been found to be associated with morbidity and mortality in several studies, especially during the neonatal period. In the present work it was found that twins had an increased risk for being hospitalized due to several diagnoses, defined by the main chapters in ICD-9, including congenital malformations, which is in accordance with a previous study by Cooke et al¹¹⁰. It was also found that being born a twin increased the risk for hospitalization due to mental disorders; this is also in alignment with previous research¹¹⁰⁻¹¹¹.

However, stratifying data by optimality at birth and adjusting for socio-demographic factors and birth characteristics, it was found that in the non-optimal group twins appeared to fare somewhat better compared with singletons. A possible explanation to this is the etiology of preterm birth among twins and singletons. Twins are more likely to be delivered preterm due to mechanical reasons as lack of space, while among singletons the major cause for preterm birth are maternal complications during pregnancy such as infections^{10-11,18}.

Intergenerational effect of non-optimal birth characteristics

As several previous studies⁷¹⁻⁷⁴, the studies included in this thesis were able to establish an intergenerational effect of non-optimal birth characteristics. A slightly different intergenerational effect of preterm birth among mothers who themselves were born preterm was detected. Mothers who were born as singletons and born preterm had an increased risk for delivering a preterm child as well as a SGA child while there was no effect of preterm birth among mothers born as twins. In addition, we established a relationship among mothers born as twins and singletons with respect to SGA, mothers born as singletons and SGA had an increased risk to deliver

a child either preterm or SGA, twin mothers born SGA also had an increased risk to deliver a child who was SGA.

These findings are in accordance with several previous studies on singletons in which being born preterm or having a sibling born preterm, increases the risk for having a preterm child⁷². These findings are also in accordance with a previous study by Selling et al⁷¹, which found an intergenerational effect of mothers born SGA having a greater risk for delivering a SGA child. However, unlike this study we were able to establish that singleton women born preterm were more likely to deliver their own children both preterm or SGA.

Reproductive pattern in relation to number of siblings

Having 3 or more siblings was found to increase the chances of becoming a first-time parent among both twins and singletons as well as becoming parent a second time. This is in part in agreement with a previous study by Murphy and Knudsen who found that one of the most important intergenerational factors with respect to future reproductive pattern was the total number of siblings¹¹². Nevertheless, the number of siblings fail to fully explain the difference in reproductive pattern among twins and singletons after adjustment for socio-demographic and birth characteristics.

We also found an indication of male twins being more likely to have a second child compared with singletons, while this difference was not present among females. There are several potential explanations to this phenomenon. First, men are usually older when becoming parents for the first time, and evidence has been presented that male twins have a reduced likelihood of reproducing as well. Thus, it is plausible to assume that male twins are older than female twins when they become parents and therefore have a higher reproductive rate once the firstborn has arrived. This finding is in alignment with the results from a Norwegian study where male reproduction increased with the number of siblings, this effect was not present among females¹¹³. Another study has also claimed that future reproduction and at

what age one decides to start a family is highly dependent on family values and may be of greater importance than “traditional” socio-demographic factors¹¹⁴.

GENERAL CONCLUSIONS

Up till now, most epidemiological studies on reproductive pattern and morbidity exclude twins (and higher order pregnancies), without explaining why. This thesis provides an addition to the current knowledge, mainly derived from intra-twin studies, on twin morbidity and reproductive pattern.

- Twins aged 13-33 years of age had a lower likelihood of reproducing compared with singletons after adjustment for both socio-demographic factors and birth characteristics (Study I).
- Having survived till the age of 13, and thereby surviving the most vulnerable time among newborn twins, twins still had an increased risk for being hospitalized in adolescence and young adulthood compared with singletons. Adjusting for both socio-demographic factors and optimality at birth this effect was diminished and even reversed, as twins with non-optimal birth characteristics seemed to fare better than non-optimal singletons (Study II).
- The intergenerational effect of preterm birth and SGA was present among both twins and singletons, though the effect seems to be greater among singletons compared with twins (Study III).
- Both male and female twins had a lower likelihood of becoming a first time parent. This decreased likelihood vanished in studying the likelihood of becoming a second-time parent and among males twins exhibited an increased likelihood of becoming a second-time parent compared with singletons. In terms of familial factors, the number of siblings affects future reproductive pattern among both twins and singletons (Study IV).

FUTURE PERSPECTIVES

This thesis provides indications that twins have a lower likelihood of reproducing compared with singletons. Since the studies had a limited follow-up, with respect to reproductive age, it is of importance to continue these studies but with a follow-up until the end of their reproductive years. This would further elucidate the importance of twinning on future reproductive performance as well as provide a measure of “total” fertility.

As the study population in this thesis mainly were born prior to IVF-treatment it is also important to study if there are any differences on optimality at birth and subsequent fertility between twins conceived by IVF and “natural” twins.

In study IV we found that the number of siblings a person has affects the future likelihood of becoming a parent. In the future it is of importance to compare if the reproductive pattern among twins and their singleton siblings are different. This could provide further knowledge on the importance of the epigenetic effect (due to the intrauterine environment) on reproductive performance and outcomes.

In study II twins showed an increased likelihood of being hospitalized due to external causes of morbidity and mortality, after adjusting for both socio-demographic background factors as well as birth characteristics. In the future it would be of interest to investigate how the personality of the child is affected by different birth characteristics and how this in turn affects the likelihood of being hospitalized due to e.g. external causes of morbidity and mortality.

In the future it would be interesting to perform a clinical study including both twins and singletons where hormonal data, uterus size, ovulation rate, sperm quality etc. are collected in addition to asking the participants questions regarding chronic conditions, medications, infertility treatment, induced and spontaneous abortions.

Adding population-based register data on parental level of education, country of birth, employment, their own birth characteristics, mother's pregnancy induced conditions, pregnancy outcome etc. would allow us to gain valuable knowledge on the reproductive pattern and morbidity among both twins and singletons.

POPULÄRVETENSKAPLIG SAMMANFATTNING

Från 70-talet fram till mitten av 2000-talet ökade andelen tvillingfödslar både i Sverige men även i ett flertal andra länder runt om i världen. Tvillingar är ofta födda för tidigt, dvs. före vecka 37, har ofta en låg födelsevikt eller är "lätt-för-tiden" vilket innebär att barnet väger lite i relation till vilken graviditetsvecka det föddes i - dessa olika utfall kallar vi icke-optimala. Ett flertal vetenskapliga studier har påvisat ett samband mellan dessa icke-optimala födelsekaraktistika och framtida reproduktionsmönster men även med sjuklighet. Trots detta inkluderar de flesta studier inte tvillingar då man undersöker reproduktiv förmåga eller sjuklighet, och således vet man inte hur tvillingars påverkan av dessa födelsekaraktistika förhåller sig i jämförelse med enlingar.

Syftet med denna studie var således att undersöka hur både tvillingars och enlingars reproduktion och sjuklighet påverkas av att vara född antingen för tidigt, med låg födelsevikt eller liten-för-tiden. För att besvara dessa frågor inhämtades anonymiserade persondata från Statistiska Centralbyrån och Socialstyrelsen.

I denna avhandling visade studierna att mellan åldrarna 13 och 33 år har tvillingar, generellt sett, en lägre sannolikhet att få barn även när man tar hänsyn till sociodemografiska faktorer såsom föräldrarnas utbildningsnivå, moderns ålder då personen själv föddes, om modern var förstföderska eller omföderska och föräldrars födelseland.

Dessutom har tvillingar överlag en högre risk för att ha kontakt med specialiserad sjukvård (öppenvård eller slutenvård) under tonåren och som unga vuxna. Men, när hänsyn tas till de olika födelsekaraktistika som tvillingar och enlingar har så försvinner många av skillnaderna och tvillingar med icke-optimala födelsekaraktistika tycks till och med ha ett lägre antal vårdbesök vid sjukhus jämfört med "icke-optimala" enlingar för ett flertal diagnoser.

Dessutom fanns ärftlighet av icke-optimala födelsekaraktistika hos både tvillingar och enlingar, men mest uttalat hos enlingar då mödrar födda lätt-för-tiden hade en

ökad risk för att få ett barn som var antingen född före graviditetsvecka 37 eller som var lätt-för tiden. Mödrar födda lätt-för-tiden hade en ökad risk för att själva få ett barn som var lätt-för-tiden. Hos tvillingar förelåg en generationseffekt endast hos de som själva var födda lätt-för-tiden som hade en ökad risk för att själva få barn som var lätt-för-tiden.

I sista studien, som var en utökad studie av första delarbetet, fanns återigen att tvillingar har en längre sannolikhet att få ett första barn jämfört med enlingar. Detta gäller både män och kvinnor. Däremot upptäcktes att bland de som fått barn, fanns det ingen skillnad i sannolikheten att få ett andra barn bland kvinnor, noterbart är att bland männen så hade tvillingar en ökad sannolikhet att få ett andra barn jämfört med enlingar. Dessa resultat håller även då man tar hänsyn till hur många syskon man har med gemensam mamma. Dessutom, ju fler syskon man har desto större är sannolikheten att få både ett och två barn för både män och kvinnor liksom tvillingar och enlingar.

ACKNOWLEDGEMENTS

There are a great number of people who deserve to be shown appreciation and being properly thanked for their support in the making of this thesis.

Professor Gunilla Sydsjö, my supervisor, thank you for introducing me to the magical world of research on pregnancy and subsequent outcomes and for letting me become a part of your research team! You have been a constant source of encouragement and support, always giving me amazingly quick and valuable feedback.

Associate professor Ann Josefsson, my co-supervisor, thank you for all valuable input and explanations regarding pregnancy and childbirth and the etiology of preterm birth. Your enthusiasm has been inspiring and your input on my research has been very valuable and appreciated.

Professor John Carstensen, my co-supervisor and mentor in statistics! Thank you for your patience and letting me ask you questions. But I'd also like to thank you for your brilliant ideas on how I can improve my statistical analyses and for your careful proofreading of manuscripts and all tables I have produced!

Professor Orvar Finnström, my co-supervisor, I would like to thank you for your careful review of all manuscripts and tables, your expertise on the neonatal care of newborns and for sharing some of your knowledge with me.

Ninni Borendal-Wodlin, thank you for your support during this process.

Jan Brynhildsen, Mats Hammar, Anders Nordgaard, Ann-Britt Wihrén, Erik Hildebrand, Caroline Lilliecreutz and Anna-Clara Spetz Holm – for reading my manuscripts and “kappa” and providing me with valuable input.

Sara Agnafors and Louise Möller, thank you for all interesting discussions, laughs and for being there when I needed someone to talk to and for giving me valuable input on my work!

Josefine Vikström, thank you for your company, discussions and support – my room is empty without you.

Kristina Kernell, Åsa Kastbom, Linda Jonsson and Maria Sarberg for many discussions on research but also everyday life discussions!

Chatarina Malm, Ingela Olsson and Maria Rundberg, thank you for all administrative help and guidance you have provided me with.

All employees at the Division of Obstetrics and Gynecology – thank you for small talk in the coffee room and hallways.

Maud, Sandra, Eva – thank you for joining me in countless number of walks and discussions on everyday matters!

Jimmy and Karin Forsberg, my brother and sister-in-law, thank you for providing me with bed and food when work and classes made it impossible to go home - your B&B is a solid 5 star!

Pia Bladh and Erik Rönnqvist, my sister-in-law and brother-in-law, thank you for all your support.

Bernt Forsberg and Birgitta Forsberg, my father and mother, your upbringing shaped me into the stubborn and curious person I am today. Without you I would not be where I am today.

Ingegerd Bladh and Paul Marcusson, my mother in-law and father in-law, your support, especially in caring for my children, during this process have been invaluable.

Above all I would like to thank my husband Ronnie and my children, Ewelina and Alexander, for their never-ending support in every conceivable manner! I hope that I in the future can compensate you for all weekends, nights and vacations where my research have made me physically but not mentally present.

REFERENCES

1. Beck S, Wojdyla D, Say L, Pilar Betran A, Merialdi M, Harris Requejo J, Rubens C, Menon R, FA Van P. The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity. *Bulletin of the World Health Organization* 2010;88:31–38
2. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller A, Narwal R, Adler A, Garcia CV, Rohde S, Say L, Lawn JE. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet* 2012; 379: 2162–72)
3. Andersen, A.M. and M. Osler, Birth dimensions, parental mortality, and mortality in early adult age: a cohort study of Danish men born in 1953. *International journal of epidemiology* 2004;33:92-9.
4. Ashworth, A., Effects of intrauterine growth retardation on mortality and morbidity in infants and young children. *European journal of clinical nutrition* 1998. 52 Suppl 1: S34-41; discussion S41-2.
5. Clausson, B., S. Cnattingius, and O. Axelsson, Preterm and term births of small for gestational age infants: a population-based study of risk factors among nulliparous women. *British journal of obstetrics and gynaecology* 1998;105:1011-7.
6. Selling K Ekholm, Carstensen J, Finnström O, Josefsson A, Sydsjö G. Hospitalizations in Adolescence and Early Adulthood Among Swedish Men and Women Born Preterm or Small for Gestational Age. *Epidemiology* 2008;19:63-70.
7. Monfils Gustafsson W, Josefsson A, Ekholm Selling K, Sydsjö G. Preterm birth or foetal growth impairment and psychiatric hospitalization in adolescence and early adulthood in a Swedish population-based birth cohort. *Acta psychiatrica et neurologica Scandinavica* 2009;119:54-61
8. Longo S, Bollani L, Decembrino L, Di Comite A, Angelini M, Stronati M. Short-term and long-term sequelae in intrauterine growth retardation. *The journal of maternal - Fetal and neonatal medicine* 2013;26:222-225.
9. Ekholm K, Carstensen J, Finnström O, Sydsjö G. The Probability of Giving Birth among Women Who Were Born Preterm or with Impaired Fetal Growth: A Swedish Population-based Registry Study. *American journal of epidemiology* 2005;161:725-33.

10. Goldenberg RL, Culhane JF, Iams JD, Romero R: Epidemiology and causes of preterm birth. *Lancet* 2008;371:75-84.
11. Romero R., Mazor M, Munoz H, Gomez R, Galasso M, Sherer DM. The preterm labor syndrome. *Annals of the New York Academy of Sciences* 2008;734:414-29.
12. Warner BB, Kiely JL, Donovan EF: Multiple births and outcome. *Clinics in perinatology* 2000;27:347-61.
13. Swamy GK, Östbye T, Skjaerven R: Association of preterm birth with long-term survival, reproduction and next-generation preterm birth. *Journal of the American Medical Association* 2008;229:1429-36.
14. Nohr EA, Vaeth M, Baker JL, Sorensen TI, Olsen J, Rasmussen KM: Pregnancy outcomes related to gestational weight gain in women defined by their body mass index, parity, height and smoking status. *The American journal of clinical nutrition* 2009;90:1288-94.
15. Rasmussen S, Kiserud T, Albrechtsen S: Foetal size and body proportion at 17-19 weeks of gestation and neonatal size, proportion and outcome. *Early human development* 2006;82:683-90.
16. Beard JR, Lincoln D, Donoghue D, Taylor D, Summerhayes R, Dunn TM, Earnest A, Morgan G: Socio-economic and maternal determinants of small-for-gestational age births: patterns of increasing disparity. *Acta obstetrica et gynecologica Scandinavica* 2009;88:575-83.
17. Imaizumi Y: Trends of twinning rates in ten countries, 1972-1996. *Acta geneticae medicae et gemellologiae* 1997;46:209-18.
18. Tandberg A, Bjørge T, Bør Dahl PE, Skjaerven R: Increasing twinning rates in Norway, 1967-2004: the influence of maternal age and assisted reproductive technology (ART). *Acta obstetrica et gynecologica Scandinavica* 2007;86:833-39.
19. National Board of Health and Welfare, Centre for Epidemiology. Statistics – Health and Medical Care, Pregnancies, Deliveries and Newborn Infants, The Swedish Medical Birth Register 1973–2007, Assisted Reproduction, treatment 1991–2006. 2009. (Article No.2009-125-5)
http://www.socialstyrelsen.se/pressrum/Documents/20091255_rev3.pdf
20. Fellman J, Erikson AW: Variations in the maternal age effect on twinning rates: the Nordic experience. *Twin research and human genetics* 2005;8:515-23.

21. Montgomery KS, Cubera S, Belcher C, Patrick D, Funderburk H, Melton C, Fastenau M, Childbirth education for multiple pregnancy part1: prenatal considerations. *The Journal of perinatal education* 2005;14:26-35
22. Montgomery KS, Cubera S, Belcher C, Patrick D, Funderburk H, Melton C, Fastenau M, Childbirth education for multiple pregnancy part1: intrapartum and postpartum considerations. *The Journal of perinatal education* 2005;14:33-38
23. Doyle LW, Saigal S. Preterm Birth 3 An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet* 2008;371: 261- 269
24. Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, Bassani DG, Jha P, Campbell H, Fischer Walker C, Cibulskis R, Eisele T, Liu L, Mathers C, for the Child Health Epidemiology Reference Group of WHO and UNICEF. Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet* 2010;375:1969–87
25. Lawn JE, Cousens S, Zupan J, for the Lancet Neonatal Survival Steering Team. Neonatal Survival 1 4 million neonatal deaths: When? Where? Why? *Lancet* 2005;365:891–900)
26. Crump C, Sundquist K, Sundquist J, Winkleby MA. Gestational age at birth and mortality in young adulthood. *Journal of the American Medical Association* 2011;306:1233-1240.
27. Räikkönen K, Pesonen AK, Roseboom TJ, Eriksson JG. Early determinants of mental health. *Best practice & research clinical endocrinology & metabolism* 2012;26:599-611.
28. Barker DJP. Mothers, babies and health in later life. 2 ed., Edinburgh, Scotland: Churchill Livingstone, 1998.
29. Hodgson ED. Perinatal Programming: Early Life Determinants of Adult Health and Disease. Taylor & Francis, 2006.
30. Svensson J, Carstensen B, Mortensen HB, Borch-Johnsen K; Danish Study Group of Childhood Diabetes Early childhood risk factors associated with type 1 diabetes--is gender important? *European journal of epidemiology* 2005;20:429-34.
31. Lucas A, Fewtrell MS, Cole TJ. Fetal origins of adult disease – the hypothesis revisited. *British medical journal* 1999;319:245-249

32. Skogen JC, Øverland S. The fetal origins of adult disease: a narrative review of the epidemiological literature. *Journal of the Royal Society of Medicine short reports* 2012;59. doi:10.1258/shorts.2012.012048
33. Susser M, Levin B. Ordeals for the fetal programming hypothesis. *British medical journal* 1999;218:885-886
34. Paneth N, Susser M. Early origin of coronary heart disease (the 'Barker hypothesis'). *British medical journal* 1995;310:411-412
35. James WPT, Nelson M, Ralph A, Leather S. Socioeconomic determinants of health: The contribution of nutrition to inequalities in health. *British medical journal* 1997;314:1545
36. Tu YK, West R, Ellison TH, Gilthorpe MS. Why evidence for the Fetal origins of adult disease might be a statistical artifact: The 'reversal paradox' for the relationship between weight and blood pressure in later life. *American Journal of Epidemiology* 2005;161:27-32
37. Tu YK, Gunnel D, Gilthorpe MS. Simpson's paradox, Lord's paradox, and suppression effects are the same phenomenon – the reversal paradox. *Emerging Themes in Epidemiology* 2008;5:2
38. Tu YK, West R, Ellison GT, Gilthorpe MS. Why evidence for the fetal origins of adult disease might be a statistical artifact: the "reversal paradox" for the relation between birth weight and blood pressure in later life. *American journal of epidemiology* 2005;161:27-32.
39. Ericson A. and Källén B. Very low birthweight boys at the age of 19. *Archives of disease in childhood. Fetal and neonatal edition* 1998;78:F171-4.
40. Yanney M. and Marlow N. Paediatric consequences of fetal growth restriction. *Seminars in fetal & neonatal medicine* 2004;9:411-8.
41. Hack M. Young adult outcomes of very-low-birth-weight children. *Seminars in fetal & neonatal medicine* 2006;11:127-37.
42. Mellekjaer L, Olsen ML, Sørensen HT, Thulstrup AM, Olsen J, Olsen JH. Birth weight and risk of early-onset breast cancer. *Cancer Causes Control* 2003;14:61-4.
43. Yuan W, Basso O, Sorensen HT, Olsen J. Indicators of fetal growth and infectious disease in childhood--a birth cohort with hospitalization as outcome. *European journal of epidemiology* 2001;17:829-3

44. Cnattingius S, Hultman CM, Dahl M, Sparén P. Very preterm birth, birth trauma, and the risk of anorexia nervosa among girls. *Archives of general psychiatry* 1999;56:634-8.
45. Mittendorfer-Rutz E, Rasmussen F, Wasserman D. Restricted fetal growth and adverse maternal psychosocial and socioeconomic conditions as risk factors for suicidal behaviour of offspring: a cohort study. *Lancet* 2004;364:1135-40.
46. Nilsson E, Ståhlberg G, Lichtenstein P, Cnattingius S, Olausson PO, Hultman CM. Fetal growth restriction and schizophrenia: a Swedish twin study. *Twin research and human genetics* 2005;8:402-8.
47. Lund LK, Vik T, Skranes J, Lydersen S, Brubakk A-M, Indredavik MS. Low birth weight and psychiatric morbidity: stability and change between adolescence and young adulthood. *Early Human Development* 2012;88:623-629.
48. Heinonen K, Räikkönen K, Pesonen A-K, Andersson S, Kajantie E, Eriksson JG, Wolke D, Lano A. Behavioural symptoms of attention deficit/hyperactivity disorder in preterm and term children born small and appropriate for gestational age; A longitudinal study. *BMC Pediatrics* 2010;10:91
49. Johnson S, Hollis C, Kochhar P, Hennessy E, Wolke D, Marlow N. Autism Spectrum Disorders in Extremely Preterm Children. *The Journal of pediatrics* 2010;156:525-31
50. Pettersson E, Sjölander A, Almqvist C, Anckarsäter H, D'Onofrio BM, Lichtenstein P, Larsson H. Birth weight as an independent predictor of ADHD symptoms; a within-twin pair analysis. *Journal of child psychology and psychiatry* 2015;56:453-459
51. Pesonen A-K, Räikkönen K, Heinonen K, Andersson S, Hovi P, Järvenpää A-L, Eriksson JG, Kajantie E. Personality of young adults born prematurely: the Helsinki study of very low birth weight adults. *Journal of child psychology and psychiatry* 2008;49:609-617
52. Gäddlin P-O, Finnström O, Samuelsson S, Wadsby M, Wang C, Leijon I. Academic achievement, behavioural outcomes and MRI findings at 15 years of age in very low birthweight children. *Acta Paediatrica* 2008;97:1426-1432
53. Thompson C, Syddall H, Rodin I, Osmond C, Barker DJ. Birth weight and the risk of depressive disorder in late life. *The British journal of psychiatry* 2001;179:450-455.

54. Stålnacke J, Lundequist A, Böhm B, Forssberg H, Smedler A-C. Individual cognitive patterns and developmental trajectories after preterm birth. *Child Neuropsychology* 2014, <http://dx.doi.org/10.1080/09297049.2014.958071>
55. Løhaugen GCC, Gramstad A, Evensen KAI, Marinussen M, Lindqvist S, Indredavik M, Vik T, Brubakk A-M. Cognitive profile in young adults born preterm at very low birthweight. *Developmental medicine & child neurology* 2010;1133-1136
56. Ancel PY, Saurel-Cubizolles MJ, Di Renzo GC, Papiernik E, Bréart G. Very and moderate preterm births: are the risk factors different? *British Journal of Obstetrics and Gynaecology* 1999;106:1162-1170
57. Clausson B, Cnattingius S, Axelsson O. Preterm and term births of small for gestational age infants: a population-based study of risk factors among nulliparous women. *British Journal of Obstetrics and Gynaecology* 1998; 105:1011-1017
58. Snelgrove JW, Murphy KE. Preterm birth and social inequality: assessing the effects of material and psychosocial disadvantage in a UK birth cohort. *ACTA Obstetrica et Gynecologica* 2015;94:766-775
59. Räisänen S, Gissler M, Saari J, Kramer M, Heinonen S. Contribution of Risk Factor to Extremely, Very and Moderately Preterm Births – Register-Based Analysis of 1,290,742 Singleton Births. *PLoS One* 2013;8:e60660.
60. Räisänen S, Gissler M, Sankilampi U, Saari J, Kramer MR, Heinonen S. Contribution of socioeconomic status to the risk of small for gestational age infants – a population-based study of 1,290,165 singleton live births in Finland. *International Journal for Equity in Health* 2013;12:s8,
61. Astone NM, Misra D, Lynch C. The effect of maternal socio-economic status throughout the lifespan on infant birthweight. *Paediatric and Perinatal Epidemiology* 2007;21:310-318
62. Correia S, Barros H. Small-for-gestational age Portuguese babies: The effect of childhood social environment, growth and adult socioeconomic conditions. *Preventive Medicine* 2014;70:102-107.
63. Moster D, Lie RT, Markestad T. Long-term medical and social consequences of preterm birth. *The New England journal of medicine* 2008;359:262-273
64. Männistö T, Väärasmäki M, Sipola-Leppänen M, Tikanmäki M, Matinoli H-M, Pesonen A-K, Räikkönen k, Järvelin M-R, Hovi P, Kajantie E. Independent living and romantic relations among young adults born preterm. *Pediatrics* 2015;135:290-297

65. D'Onofrio BM, Class QA, Rickert ME, Larsson H, Långström N, Lichtenstein P. Journal of the American Medical Association psychiatry 2013;70:1-9
66. Kajantie E, Hovi P, Räikkönen K, Pesonen A-K, Heinonen K, Järvenpää A-L, Eriksson JG, Strang-Karlsson S, Andersson S. Young adults with very low birth weight: Leaving the parental home and sexual relationships – Helsinki study of very low birth weight adults. Pediatrics 2008;122:e62-e72
67. Ibáñez I, Potau N, Ferrer A, Rodriguez-Hierro F, Marcos MV, de Zegher F. Reduced ovulation rate in adolescent girls born small for gestational age. Journal of clinical endocrinology and metabolism 2002;87:3391-3393
68. Main KM, Jensen RB, Askund C, Hoi-Hansen CE, Skakkebaek NE. Low birth weight and male reproductive function. Hormone research 2006;31:116-122
69. Crump C, Sundquist k, Winkleby MA, Sieh W, Sundquist J. Gestational age at birth and risk of testicular cancer. International journal of cancer 2011;131:446-451
70. deKeyser N, Josefsson A, Bladh M, Carstensen J, Finnström O, Sydsjö G. Premature and low birthweight are associated with a lower rate of reproduction in adulthood: a Swedish population-based registry study. Human reproduction 2012;27:1170-1178
71. Selling KE, Carstensen J, Finnström O, Sydsjö G. Intergenerational effects of preterm birth and reduced intrauterine growth: a population-based study of Swedish mother-offspring pairs. British journal of obstetrics and gynaecology 2006;113:430-40.
72. Bhattacharya S, Raja EA, Mirazo ER, Campbell DM, Lee AJ, Norman JE, Bhattacharya S. Inherited predisposition to spontaneous preterm delivery. Obstetrics and gynecology 2010;115:1125-33.
73. Schaaf JM, Hof MHP, Mol BWJ, Abu-Hanna A, Ravelli ACJ. Recurrence risk of preterm birth in subsequent singleton pregnancy after preterm twin delivery. American journal of obstetrics & gynecology 2012;207:279.e1-7
74. Schaaf JM, Hof MHP, Mol BWJ, Abu-Hanna A, Ravelli ACJ. Recurrence risk of preterm birth in subsequent twin pregnancy after preterm singleton delivery. British journal of obstetrics and gynaecology 2012;119:1624-9
75. De Stavola BL, Leon DA, Koupil I. Intergenerational correlations in size at birth and the contribution of environmental factors. American journal of epidemiology 2011;174:52-62

76. Jaquet D, Swaninathan S, Alexander GR, Czwrnichow P, Collin D, Salihu HM, Kirby RS Lévy-Marchal C. Significant paternal contribution to the risk for small for gestational age. *British journal of obstetrics and gynaecology* 2005;112:153-159
77. Wu G, Imhoff-Kunsch B, Webb Girard A. Biological mechanisms for nutritional regulation of maternal health and fetal development. *Paediatric and perinatal epidemiology* 2012;123:2929-2937
78. Galjard S, Devlieger R, Van Assche FA. Fetal growth and developmental programming. *Journal of perinatal medicine* 2013;41:101-105
79. Gordon L, Joo J, Powell JE. Neonatal DNA methylation profile in human twins is specified by complex interplay between intrauterine environmental and genetic factors, subject to tissue-specific influence. *Genome research* 2012;22:1395-1406
80. Souren NYP, Lutsik P, Gasparoni G, Tierling S, Gries J, Riemenschneider M, Fryns JP, Derom C, Zeegers MP, Waljter J. Adult monozygotic twins discordant for intra-uterine growth have indistinguishable genome-wide DNA methylation profiles. *Genome biology* 2013;14:R44
81. Ollikainen M, Smith Kr, Joo EJ, Andronikos R, Novakovic B, Abdul Aziz NK, Morley R, Saffery R, Craig JM. DNA methylation analysis of multiple tissues from newborn twins reveals both genetic and intrauterine components to variation in the human neonatal epigenome. *Human molecular genetics* 2010;19:4176-4188
82. Banister CE, Koestler DC, Maccani MA, Padbury JF, Houseman EA, Marsit CJ. Infant growth restriction is associated with distinct patterns of DNA methylation in human placentas. *Epigenetics* 2011;6:920-927
83. Menon R, Conneely KN, Smith AK. DNA methylation: An epigenetic risk factor in preterm birth. *Reproductive sciences* 2012;19:6-13
84. Centre for Epidemiology, National Board of Health and Welfare. The Swedish Medical Birth Register; A summary of content and quality () National Board of Health and Welfare, 2003, Stockholm, Sweden
85. Sveriges kommuner och landsting. Kvalitet i sjukvårdsdata, indikatorbeskrivningar och vårdkonsumtion. Bilagor till Öppna jämförelser av hälso- och sjukvårdens kvalitet och effektivitet 2008.
(<http://skl.se/download/18.33ccf562145ac94e998f15ac/1401187917981/skl-OJvard08-bilaga1-3.pdf>) 2008, Sveriges kommuner och landsting: Solna, Sweden

86. National Board of Health and Welfare. In English – the National Patient Register. (<http://www.socialstyrelsen.se/register/halsodataregister/patientregistret/inenglish>)
87. Ludvigsson JF, Andersson E, Ekbom A, Feychting M, Kim JL, Reuterwall C, Heurgren M, Otterblad Olausson P. External review and validation of the Swedish national inpatient register. BMC Public Health 2011;11:450
88. Centre for Epidemiology, National Board of Health and Welfare. Dödsorsaksstatistik Historik, produktionsmetoder och tillförlitlighet. Komplement till rapporten Dödsorsaker 2008. (<http://www.socialstyrelsen.se/Lists/Artikelkatalog/Attachments/18019/2010-4-33.pdf>) National Board of Health and Welfare, 2008, Stockholm, Sweden
89. Centre for Epidemiology, National Board of Health and Welfare. Dödsorsaksregistret (<http://www.socialstyrelsen.se/register/dodsorsaksregistret>)
90. Statistics Sweden. Registret över totalbefolkningen (RTB) (http://www.scb.se/Pages/List_257499.aspx)
91. Population and welfare department, Statistics Sweden. The total population register. The population as at 31 December 2011. (http://ec.europa.eu/eurostat/cache/metadata/Annexes/demo_gind_esms_se_a_n1.docx) Statistics Sweden, 2012, Örebro, Sweden.
92. Statistics Sweden. Multi-generation register 2002; a description of contents and quality (Serial no. 2003:5 (http://www.scb.se/statistik/publikationer/BE9999_2011A01_BR_BE96BR1102.pdf)). 2003, Statistics Sweden: Örebro, Sweden
93. <http://www.jpi-dataproject.eu/Home/Database/360?topicId=10>
94. Statistics Sweden. Educational Attainment of the Population 2002. (Publication no. UF0506). (<http://www.scb.se>). 2003, Statistics Sweden: Örebro, Sweden
95. Statistics Sweden. Population and Housing Census 1970 (SOS). Part 12. National Central Bureau of Statistics: Stockholm, Sweden
96. Statistics Sweden. The Swedish Register of Education. (http://www.scb.se/statistik/UF/UF0506/Produktbeskrivning_short_English_UF0506_20040101r.doc) 2004, Statistics Sweden, Örebro, Sweden
97. Kirkwood BR, Sterne JAC. Medical Statistics. 2nd ed. Oxford: Blackwell Science, 2003.

98. Morin I, Morin L, Zhang X, Platt RW, Blondel B, Bréart G, Usher R, Kramer MS. Determinants and consequences of discrepancies in menstrual and ultrasonographic gestational age estimates. *British journal of obstetrics and gynaecology* 2005;112:145-142
99. Marsal K, Persson PH, Larsen T, Lilja H, Selbing A, Sultan B. Intrauterine growth curves based on ultrasonically estimated foetal weights. *Acta Paediatrica* 1996;85:843-848.
100. Centre for Epidemiology, National Board of Health and Welfare. Graviditeter, förlossningar och nyfödda barn.
(<http://www.socialstyrelsen.se/Lists/Artikelkatalog/Attachments/19627/2014-12-19.pdf>). 2014, National Board of Health and Welfare, Stockholm, Sweden
101. World Health Organization. BMI classification
(http://apps.who.int/bmi/index.jsp?introPage=intro_3.html). 2015, World Health Organization
102. Christensen RD, Baer VL, Henry E, Snow GL, Butler A, Martha C. Sola-Visne MC. Thrombocytopenia in Small-for-Gestational-Age Infants. *Pediatrics* 2015;136:e361-e370
103. Varella MH, Moss WJ. Early growth patterns are associated with intelligence quotient scores in children born small-for-gestational age. *Early Human Development* 2015;91:491-497
104. Díaz M, Bassols J, Sebastiani G, López-Bermejo A, Ibáñez L, F de Zegher F. Circulating GLP-1 in infants born small-for-gestational-age: breast-feeding versus formula-feeding. *International Journal of Obesity* 2015;1–3
105. Ibáñez L, Potau N, Enriquez G, Marcos MV, de Zegher F. Hypergonadotrophinaemia with reduced uterine and ovarian size in women born small-for-gestational-age. *Human Reproduction* 2003;18:1565-1569
106. Petersen I, Martinussen T, McGue M, Bingley P, Christensen K. Lower Marriage and divorce rates among twins compared to singletons in Danish birth cohorts 1940-1964. *Twin research and human genetics* 2011;2:150-157
107. Middeldorp CM, Cath DC, Vink JM, Boomsma DI. Twin and genetic effects on life events. *Twin research and human genetics* 2005;3:224-231
108. Hjern A, Ekéus C, Rasmussen F, Lindblad F. Educational achievement and vocational career in twins – a Swedish national cohort study. *ACTA Paediatrica* 2012;101:591-596.

109. Blakely T, Hunt D, Woodward A. Confounding by socio-economic position remains after adjusting for neighbourhood deprivation: an example using smoking and mortality. *Journal of epidemiology and community health* 2004;58:1030-1031
110. Cooke RWI. Does neonatal and infant neurodevelopmental morbidity of multiples and singletons differ? *Seminars in fetal & neonatal medicine* 2010;15:362-366
111. Scher AI, Petterson B, Blari E, Ellenberg JH, Grether JK, Haan E, Reddiough DS, Yeargin-Allsop DS, Nelson KB. The risk of mortality or cerebral palsy in twins: a collaborative population-based study. *Pediatric Research* 2002;52:671-681
112. Murphy M, Knudsen LB. The intergenerational transmission of fertility in contemporary Denmark: The effects of number of siblings (full and half), birth order, and whether male or female. *Population Studies* 2010;56:235-248
113. Statistics Norway. The effect of childhood family size on fertility in adulthood. New evidence from IV estimation (Serial no 2015:802). 2015, Statistics Norway, Oslo, Norway
114. Kolk M. Understanding transmission of fertility across multiple generations – Socialization or socioeconomics. *Research in social stratification and mobility* 2014;35:89-103

Papers

The articles associated with this thesis have been removed for copyright reasons. For more details about these see:

<http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-121279>