

One- and two-year follow-up of a randomized trial of neck-specific exercise with or without a behavioural approach compared with prescription of physical activity in chronic whiplash disorder

Maria Landén Ludvigsson, Gunnel Peterson, Asa Dederling and Anneli Peolsson

Linköping University Post Print



N.B.: When citing this work, cite the original article.

Original Publication:

Maria Landén Ludvigsson, Gunnel Peterson, Asa Dederling and Anneli Peolsson, One- and two-year follow-up of a randomized trial of neck-specific exercise with or without a behavioural approach compared with prescription of physical activity in chronic whiplash disorder, 2016, Journal of Rehabilitation Medicine, (48), 1, 56-64.

<http://dx.doi.org/10.2340/16501977-2041>

Copyright: Foundation for Rehabilitation Information

<http://www.medicaljournals.se/jrm/>

Postprint available at: Linköping University Electronic Press

<http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-126145>

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

One- and two-year follow-up of a randomized trial of neck-specific exercise with or without a behavioural approach or prescription of physical activity in chronic whiplash.

Maria Landén Ludvigsson, MSc, PhD-student^{1,2}, Gunnel Peterson MSc, PhD-student^{1,3}, Åsa Dederig, PhD, Associate Professor⁴, Anneli Peolsson, PhD, Associate Professor¹

¹ Department of Medical and Health Sciences, Division of Physiotherapy, Linköping University, Linköping, Sweden, ² Rehab Väst, County Council of Östergötland, Motala, Sweden, ³ Centre for Clinical Research Sörmland, Uppsala University, Uppsala, Sweden and county council of Sörmland, Katrineholm, Sweden. ⁴ Division of Physiotherapy, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, and Department of Physical Therapy, Karolinska University Hospital, Stockholm, Sweden

Financial disclosure and conflicts of interest The study was supported by funding from the Swedish government through the REHSAM foundation, the Swedish Research Council, the regional Center for Clinical Research and the County Council of Östergötland, Centre for Clinical Research Sörmland at Uppsala University, the Medical Research Council of Southeast Sweden, and the Uppsala-Örebro Regional Research Council, Sweden. The authors declare no conflict of interest.

Running head: Long-term follow-up of exercise in chronic whiplash

Abstract

Objective We aimed to explore whether neck-specific exercise, with or without a behavioural approach, has benefits after 1 and 2 years over prescribed physical activity regarding pain, self-rated functioning/disability, and self-efficacy in chronic whiplash management. **Design** This was a follow-up of a randomized, assessor blinded, clinical trial. **Patients** Participants were 216 volunteers with chronic whiplash associated disorders, grade 2 or grade 3. **Methods** Participants were randomized to one of three exercise interventions: neck-specific exercise with or without a behavioural approach, or physical activity prescription. Self-rated pain (Visual Analogue Scale), disability (Neck Disability Index/Patient Specific Functional Scale), and self-efficacy (Self-Efficacy Scale) were evaluated after 1 and 2 years. **Results** Both neck-specific exercise groups maintained more improvement regarding disability/functioning than the prescribed physical activity group at both time points ($p \leq 0.02$). At 1 year, 61% of those in the neck-specific group reported at least 50% pain reduction, compared with 26% of those in the physical activity prescription group ($p < 0.001$), but at 2 years the difference was not significant. **Conclusion** After 1 to 2 years, participants with chronic whiplash who were randomized to neck-specific exercise, with or without a behavioural approach, remained more improved than participants who were prescribed general physical activity.

Key words: whiplash, chronic, exercise, randomized, follow-up study, spine, behavior therapy

Journal of Rehabilitation Medicine

Guarantor: Maria Landén Ludvigsson, Department of Medical and Health Sciences, Division of Physiotherapy, Linköping University, SE-581 83 Linköping, Sweden. E-mail: Maria.Landen.Ludvigsson@liu.se

Introduction

At 1 year post-injury, 50% of people with whiplash-associated disorders (WAD) still report neck pain (1). Despite the significant impact of WAD, there is still no clear evidence regarding which treatment is most effective (2-4). The persistence of symptoms in individuals with WAD has been attributed to both physical and psychosocial factors (5). It is therefore reasonable to assume that a behavioural approach may be of benefit in the management of chronic WAD, as in chronic back pain (6). However, impairments and altered patterns of muscle activation (7-11) and muscle deformation (12) in the cervical spine are also features of chronic WAD, suggesting that treatments aimed at improving muscle function might also be of importance. Although exercise is considered a safe treatment for neck pain, with temporary and benign side effects, its efficacy in the context of chronic WAD remains unclear (2, 3). Short-term pain reduction may be achieved (13), but there is no evidence regarding what the optimal exercise approach would be in WAD (13). Although physical activity prescription (self-directed general physical activity outside the health care system, PPA) is often recommended for patients with chronic pain (14), the relative effectiveness of different exercise regimens in relieving chronic WAD remains unknown (13). Regarding chronic WAD grades 1-2, Michaleff et al. found that a comprehensive exercise programme including neck-specific exercise was not more effective for pain reduction than advice alone (15). Another randomized controlled trial previously reported by our group, found that patients with chronic WAD grades 2-3 experienced greater reductions in neck pain and neck-related disability after neck-specific exercise, with or without the addition of a behavioural approach, compared with PPA (16). However, those results were only analyzed up to 6 months post-inclusion.

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

The aim of this study was to explore whether, after 1 and 2 years, neck-specific exercise with or without a behavioural approach has long-term benefits over PPA regarding pain, self-rated functioning/disability, and self-efficacy in the management of grade 2 or grade 3 chronic WAD.

Methods

Design

This is a 1- and 2-year follow-up of a multi-centre, prospective, randomized clinical trial with assessor and group allocation blinding (Clinical Trials.gov, NCT01528579 (17)). Short-term results have been presented previously (16). The study, conducted in accordance with the Helsinki declaration, was approved by the Regional Ethics Committee of Linköping University, Sweden.

Participants and settings

Two hundred and sixteen individuals with chronic WAD were recruited between February 2011 and May 2012, including 142 (65%) women and 74 (35%) men with a mean age of 40 (range 18-63, SD 11.4) years (Table I). Inclusion criteria were age 18-63 years, and a grade 2 or 3 whiplash injury according to the Quebec Task Force (18) in the preceding 6-36 months that was nominated as the cause of current symptoms. Additional inclusion criteria were a Neck Disability Index score (NDI) (19) of at least 10/50 points, and/or an average pain rating on the Visual Analogue Scale (VAS) of >20/100 mm (0 = no pain, 100 = worst imaginable pain) (20) for the preceding week. Exclusion criteria were signs of traumatic brain injury, previous neck trauma with unresolved symptoms, more dominant pain elsewhere, neck pain causing more than 1 month of absence from work in the year preceding the whiplash injury, myelopathy, spinal infection or tumor, previous neck surgery, conditions that were potentially

detrimental to completing the study interventions (e.g., severe psychiatric disorders or known drug abuse) or insufficient knowledge of the Swedish language. Experienced physiotherapists conducted the interventions in a primary care setting. The physiotherapists were provided with standardized oral information about their interventions. Those in charge of physiotherapist-led interventions participated in a 1-day workshop of standardized practical and theoretical training held by the project leaders.

One hundred seventy patients (79%) completed the 1-year follow-up, and 123 patients (57%) completed the 2-year follow-up (Fig. 1). With imputed values (see statistics), data was available from 193 participants (89%) at 1 year and from 184 (85%) participants at 2 years. There was no baseline difference ($p>0.27$ for all) between those who completed the questionnaires and drop-outs at either time point regarding allocation, gender, WAD grade, pain or pain bothersomeness, NDI (1 year only), or age (2 years only). Drop-outs at 1 year were somewhat younger [age 37 years (SD11)] than those who completed the study [age 41 years (SD 11), $p=0.04$]. Drop-outs at 2 years reported more baseline disability than those who completed the study [NDI 17.9 (SD 6.9) vs. NDI 15.7 (SD 6.5), $p<0.04$]. At 2 years, drop-outs and completers did not differ regarding preceding improvement from baseline to 1 year (NDI and neck pain VAS, $p>0.94$).

Table 1. Baseline characteristics of participants in the three randomized groups

Variable	NSE (n=76)	NSEB (n=71)	PPA (n=69)	P-value
Gender, female, n (%)	57 (75)	47 (66)	38 (55)	0.04
Age, mean (range) SD	38 (18-62)11.3	40 (19-63)11.6	43 (19-63)10.7	0.03
Months since injury, mean(range)SD	19 (6-36)8.7	20 (6-36) 8.9	20 (6-36) 10.3	0.69
WAD grade 2/3	49/27 (64/36)	33/38 (46/54)	41/28 (58/42)	0.08
Smoker, n (%)	17 (22)	8 (11)	12 (18)	0.22
Educational level				0.44
Elementary school, n (%)	4 (5)	6 (9)	6 (9)	
High school, n (%)	38 (50)	40 (57)	34 (51)	
University, n (%)	31 (41)	21 (30)	24 (36)	
Other, n (%)	3(4)	3(4)	3 (4)	
Employed n (%)	61 (80)	57 (80)	52 (75)	0.71
General Health, EQ-5D-score, median (IQR)	0.72 (0.69-0.76)	0.73 (0.23-0.80)	0.73 (0.66-0.80)	0.62
Physical activity level, IPAQ				0.64
Low, n (%)	40 (56)	36 (56)	39 (58)	
Medium, n (%)	12 (17)	8 (13)	15 (22)	
High, n (%)	19 (27)	20 (21)	13 (19)	
Unsettled insurance claim, n (%)	16 (21)	17 (24)	11 (16)	0.39
Baseline outcome measurements:				
NDI, mean (SD) / median (IQR)	16(6)/15(12-21)	17(7)/17(11-21)	17(7)/17(13-22)	0.47
VAS P, mean (SD) / median (IQR)	40(24)/38(20-62)	45(24)/50(19-65)	42(25)/42 (20-60)	0.51
VAS B, mean (SD) / median (IQR)	49(22)/49(32-65)	50(23)/52(31-70)	48(22)/49(30-67)	0.87
SES, mean (SD) / median (IQR)	150(34)/154(130-174)	153(35)/159(127-184)	147(41)/157(124-182)	0.52
PSFS, mean (SD) / median (IQR)	4.4(1.7)/ 4.3(3.3-6.0)	4.5(2.1)/4.3(3.0-6.0)	4.6(1.8)/5.0(3.3-5.7)	0.80
Use of analgesic drugs, yes (%)	40 (53)	44 (62)	45 (67)	0.23

NSE = Neck-specific exercise group, NSEB = Neck-specific exercise group with a behavioral approach, PPA = Prescription of physical activity group, EQ-5D = Euroqol-5D health questionnaire, IPAQ = International Physical Activity Questionnaire, NDI= Neck Disability Index, VAS P = Pain Visual Analogue Scale, VAS B = Pain Bothersomeness Visual Analogue Scale, SES = Self Efficacy Scale, PSFS = Pain Specific Functional Scale

Procedure

Potential participants were identified from the health care registries of six Swedish counties, including primary health care centers and hospital outpatient services. Participants were screened for eligibility through the following four-step process (Fig. 1): 1) an initial screening

letter that contained study information, basic inclusion/exclusion criteria (Fig.1), NDI (19) and Pain VAS (P-VAS) measures 2) a telephone interview with one of the project leaders; 3) a review of medical records, if needed due to any uncertainties; and 4) a physical examination by an experienced physiotherapist (mean 18 years of experience) to confirm findings consistent with either WAD grade 2 (neck pain and clinical findings) or WAD grade 3 (addition of neurological signs) (18).

All participants received verbal and written information about the study. Informed consent and baseline measurements were collected before allocation. Allocation from a computer-generated randomization list was made by an independent researcher, who also put the individual results in sealed opaque envelopes for further distribution to the treating physiotherapists. Collection and entering of data was made by staff blinded to allocation.




Figure 1

Figure 1. Patient disposition throughout the study.

*Whiplash injury in the preceding 6 to 36 months, reported as the onset of current symptoms, excluding unconsciousness/loss of memory in connection to the whiplash injury; previous neck trauma with unresolved symptoms; previous neck surgery; ongoing malignant disease; severe psychiatric disorders; drug abuse; difficulty understanding the Swedish language. NDI=Neck Disability Index; VAS=Visual Analogue Scale; WAD=whiplash-associated disorders.

Interventions

First, all patients were examined by their treating physiotherapist. All three interventions were undertaken during a 12-week period, and participants were encouraged to continue exercising on their own after the interventions. The interventions, as previously described in more detail (16), were: physiotherapist-led neck-specific exercise (NSE), NSE with the addition of a behavioural approach (NSEB), or prescribed physical activity (PPA). No serious harms were reported.

Neck-specific exercise, NSE

Participants in the NSE group undertook supervised, neck-specific exercise twice weekly. They initially also practiced daily at home, focusing on activity of the deep cervical muscles. Next, gym exercise within participants' symptom tolerance was gradually introduced, with progressive head resistance training that focused on low load endurance in flexion, extension, rotation, and lateral flexion. Exercise-related pain provocation was not accepted in this group. A detailed description of the exercises can be found at the Academic Archive On-line (21). Participants also received a written individualized exercise program that contained exercises from the interventions and general physical activity towards the end of the 12-week intervention.

Neck-specific exercise with a behavioural approach, NSEB

The protocol of exercises in the NSEB group was the same as that undertaken by the NSE group, with the addition of a behavioural approach. The behavioral approach was aimed to be basic and manageable by experienced physiotherapists in primary care with some previous knowledge of behavioral approaches. In accordance with the concept of graded exercise, patients were encouraged not to focus on temporary increases in neck pain. They also received physiotherapist-guided behavioural interventions, including education and

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

introduction to activities aimed at pain management and problem-solving. Timeframes and specific components of the interventions have been described previously (16).

Prescribed physical activity, PPA

Participants in the PPA group initially received a short motivational interview by a physiotherapist. Based on this interview and a physical examination, they were prescribed individualized general physical activity to be performed independently. The purpose of this prescription was to increase overall physical activity, either with individualized home exercise or activities performed in public gyms or elsewhere, outside the health care system. Neck-specific exercises that included any form of head resistance were not prescribed in this group. A single follow-up visit or phone call was encouraged. Participants in this group were also encouraged to continue exercise post-interventions.

Outcomes

All measurements were recorded at baseline and after 1 and 2 years. The individually chosen activities were recorded verbally in the Patient-Specific Functional Scale (PSFS) by a blinded investigator in connection to other clinical tests, to ensure that they were individually standardized with regard to time and/or repetitions.

Disability and functioning

The primary outcome measurement was the NDI (19), which consists of 10 items that grade neck disability from 0 (no activity limitations) to 5 (major activity limitations), with a total maximum score of 50 points (19). A higher score represents a greater level of disability. The NDI is considered a reliable and valid measurement of disability in patients with neck pain disorders (22).

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

The PSFS was used to measure functioning. For the PSFS, each patient nominated three individual activities related to work, leisure, and physical activity/exercise that they were unable to do or experienced difficulty performing because of their neck condition. These activities were ranked according to functional level on a scale from 0 (unable to do) to 10 (functional level equal to preinjury status), and the average of the scores was calculated. The PSFS has been shown to have excellent reliability (23) and responsiveness in the chronic WAD population (24).

Pain

Current neck pain intensity was measured using a Pain VAS (P-VAS) anchored by 0 = no pain, and 100 = worst imaginable pain (20). Bothersomeness of neck pain (B-VAS) was recorded for the preceding 24 hours (anchored by 0 = not bothersome at all, 100 = extremely bothersome). Pain bothersomeness is reportedly more responsive than pain intensity in individuals with WAD (24). Patients also recorded whether they used analgesics to manage their neck pain (yes/no).

Self-efficacy

The participants' confidence in their ability to perform activities despite pain was evaluated using the Self-Efficacy Scale (SES) (25, 26). The SES is a reliable instrument in WAD populations (5, 27); it consists of 20 different physical and psychosocial activity items that are scored from 0 = not confident at all, to 10 = very confident, thus generating a total score from 0-200.

Adherence

Participants were asked to estimate their adherence to their post-intervention prescribed exercise on a 4-point scale: full, fair, some, or no adherence. Participants who reported some to full adherence were classified as adherent in the analysis.

Clinical relevance

The Initiative on Methods, Measurement and Pain Assessment in Clinical Trials (IMMPACT) concluded that it is impossible to provide specific guidelines for determining whether or not a group difference is clinically meaningful (28). Therefore, we also calculated the proportion of patients that achieved clinically important improvement where such cut-offs have been established. The Minimal Clinically Important Difference (MCID) of the NDI score is suggested to be 3.5-5/50 points (19), with a reduction of 5 set as the cut-off score in this study. This exceeds the measurement error in this study sample, as previously reported [Minimum Detectable Change (MDC) of 3.3/50 points (16). In the PSFS, the MCID is reported as 2.3 and the MDC is reported as 2.1 (29). An increase of 2.3 was set as the cut-off score in this study. A reduction in pain intensity of $\geq 50\%$ is suggested to indicate substantial improvement or treatment success according to IMMPACT recommendations (30). This level is also used in this study to define clinically important improvement. To our knowledge, the MCID has not been established in the SES.

Statistics

The required sample size for the original randomized controlled trial was determined on the basis of the expected difference between the three groups (alpha 5%, power 80%) for main outcome, NDI (3.5/50, SD 7), and allowing for a 10% dropout rate, rendering a sample size

of 216. If only one item of data was missing from the NDI scores or two items were missing from the SES, the missing data were substituted by the average item score of the questionnaire for that participant. If more data items were missing, that particular score was omitted from the analysis. Missing scores were considered MAR (missing at random), such that the closest match is considered a reliable and efficacious imputation method in repeated measures data. Closest match replaces a participant's missing time point with a value obtained from another participant who has similar scores on the same measure assessed at other time points (31). Only participants with complete measurements from three time points served as possible donors, and participants with missing data served as recipients. The score obtained by the donor at that time point was imputed to the missing time point for the recipient. Closest match was defined as all donors with less than 15% absolute differences between the recipient's scores and donor's scores at all other time points for which data was present, and with the same trend over time (better, worse, or unchanged). Priority was given to absolute matches, and each outcome was analyzed and imputed separately. When more than one possible donor was identified, the computer (SPSS version 22) selected the closest match donor at random from the possible donors. Because baseline adherence data cannot be measured, these values were not imputed. The presented results are from the imputed data set. Use of imputed versus non-imputed values in the analysis did not cause a significant difference in the results.

The primary analysis was conducted using SPSS version 22 on an intent-to-treat basis, including all patients completing each measurement and imputed data. Descriptive statistics were calculated, and between-group comparisons were evaluated with the Kruskal-Wallis test for non-parametric data, with the Mann-Whitney U test for post-hoc, or with one-way ANOVA for normally distributed parametric data. In binary outcomes, chi-square tests were

used. Within-group differences at baseline, 6, 12, and 24 months were calculated using a Friedman's ANOVA with post-hoc Wilcoxon's signed rank test. Correlations were calculated using Spearman's rho. To determine the proportion of responders in each group, we performed sub-analyses of participants who had reached the predefined cut-off values as described previously. Statistical significance was set at $p < 0.05$ (with Bonferroni post-hoc correction at 0.017). To aid comparability to previous and other publications in this field, all outcomes are presented with both mean and median values, and data is also presented for separate WAD grades.

Results

Disability and functioning

The between-group analysis indicated that both physiotherapist-led neck-specific exercise groups reported greater reductions of neck disability and improved functioning ($p < 0.01$) than the PPA group at the 1-year follow-up. At 2 years, the NSEB group reported significant reduction in neck disability compared with the PPA group ($p \leq 0.02$); however the NSE group reported significant improvement in functioning compared with the PPA group ($p = 0.02$) (Table II). There was however no difference between the two physiotherapist-led groups. A greater proportion of individuals in the two physiotherapist-led neck-specific groups than in the PPA group experienced clinically relevant improvement in disability/functioning at 1 year ($p < 0.01$) and 2 years (disability $p = 0.01$); however, the difference regarding functioning was not statistically significant at 2 years ($p = 0.32$) (Fig. 2).

The within-group results indicated that both neck-specific exercise groups (but not the PPA group) improved over time (disability $p =$ NSE 0.01, NSEB < 0.001 , PPA 0.13, functioning $p =$

NSE/NSEB <0.001, PPA 0.22). The improvements were gained in the first 6 months (16) and were maintained over time (table III).

Table 2. Change score from baseline to 1 and 2 years (median(IQR)/mean(SD), with between group significance level

1 year							
Outcome	n	NSE		n	NSEB		p
		median(IQR)/mean(SD)			median(IQR)/mean(SD)		
NDI	65	-3.0 (-6.5– 3.5)/-2.8 (7.0)		68	-4.0 (-8.0– 0)/-3.9 (6.5)		<0.001
grade 2	44	-2.6 (-6.0–2.5)/-2.3 (7.0)		31	-3.0(-7.0– 0)/-3.4 (6.5)		
grade 3	21	-4.0 (-9.0– 1.0)/-3.8 (7.1)		37	-4.0 (-10.0– 0.5)/-4.4 (6.5)		
PSFS	62	2.3 (0.3– 4.4)/2.3 (2.4)		66	2.0 (0– 4.3)/2.1 (2.7)		<0.01
grade 2	40	2.5(0.3–3.9)/2.2(2.4)		30	1.8 (0–4.4)/2.1 (2.6)		
grade 3	22	2.0 (1.1– 4.5)/2.4 (2.4)		36	2.0 (0.2– 4.2)/2.0 (2.9)		
VAS P	66	-11 (-30– 4)/-13 (23)		68	-9 (-24– 3)/-12 (25)		0.15
grade 2	44	-9 (-29– 4)/-13 (22)		31	-6 (-21– 3)/9 (23)		
grade 3	22	-17 (-33– 8)/-13 (26)		37	-14 (-35– 7)/-15 (26)		
VAS B	66	-16 (-35– 6)/-15 (27)		68	-12 (-30– 1)/-13 (26)		0.29
grade 2	44	-20 (-39– 7)/-15 (29)		31	-12 (-30– 0)/-12 (27)		
grade 3	22	-7 (-32– 6)/-14 (24)		37	-12 (-32– 2)/-14 (27)		
SES	64	5 (-10–21)/6 (26)		68	3 (-11– 26)/8 (31)		0.41
grade 2	42	6 (-10– 23)/6 (28)		31	0 (-8– 15)/4 (21)		
grade 3	22	4 (-10– 15)/5 (24)		37	8 (-14– 36)/11 (38)		
2 years							
Outcome	n	NSE		n	NSEB		p
		median(IQR)/mean(SD)			median(IQR)/mean(SD)		
NDI	61	-2.0 (-5.5– 3.5)/-1.8 (6.1)		66	-3.0 (-9– -1)/-3.7 (6.4)		0.001²
grade 2	41	-1.0(-5.0– 4.0)/-0.8 (6.1)		31	-3.0 (-10.0– 1.0)/-3.8 (7.1)		
grade 3	20	-2.5 (-6.8– 1.0)/-3.2 (6.0)		35	-3.0 (-9.0 1.0)/-3.5 (5.9)		
PSFS	62	1.3 (-0.2– 3.8)/1.7 (2.4)		63	1.2 (0–3.3)/1.5 (2.7)		0.02¹
grade 2	40	1.3 (-0.5–3.9)/1.7 (2.7)		29	0.8 (-0.1– 3.3)/1.4 (2.8)		
grade 3	22	1.6 (-0.1– 3.3)/1.7 (2.0)		34	1.8 (-0.1– 3.4)/1.5 (2.5)		
VAS P	62	-11 (-33– 0)/-14 (25)		54	-13 (-33– 6)/-14 (22)		0.67

NSE = neck-specific exercise group, NSEB = neck-specific exercise group with a behavioral approach, PPA = prescription of physical activity group, NDI = Neck Disability Index, PSFS = Pain Specific Functional Scale, VAS = Visual Analogue Scale, SES = Self-efficacy Scale

Pain

There were no significant between-group differences ($p \geq 0.15$) regarding reduction in neck pain or pain bothersomeness at the 1-year or 2-year follow-up (Table II). Although significantly fewer participants in the two neck-specific groups reported taking analgesics for neck pain (NSE 47%, NSEB 50%, PPA 69%, $p=0.04$) at 1 year, the difference was non-significant at 2 years (NSE 44%, NSEB 53%, PPA 61%, $p=0.31$). However greater

proportions of individuals in the two physiotherapist-led neck-specific groups reported clinically relevant improvements of pain/pain bothersomeness after 1 year, as indicated by 50% pain reduction ($p \leq 0.02$); but the difference was non-significant at 2 years ($p \geq 0.11$; Fig. 2).

The within-group results showed that current pain and pain bothersomeness improved significantly over time in both neck-specific groups ($p = \text{NSE}, \text{NSEB} \leq 0.001$). Pain bothersomeness improved in the PPA group ($p = 0.02$), although current pain did not ($p = 0.07$). The improvements were gained in the first 6 months (16) and maintained over time without significant changes thereafter for either outcome (table III).

Self-Efficacy

There were no significant between-group differences regarding self-efficacy ($p < 0.17$) at the 1-year or 2-year follow-up. The within-group results demonstrated improvement for the NSE group only over time ($p = \text{NSE } 0.02, \text{NSEB } 0.07, \text{PPA } 0.86$), and there were no changes in any of the groups after 6 months ($p > 0.12$) (table III).

Adherence

At the 1-year follow-up, there was no between-group difference regarding self-reported adherence to post-intervention prescribed exercise ($p = 0.23$). Seventy-nine percent of the patients in the NSE and PPA group and 69% of patients in the NSEB group reported some to full adherence. At 2 years there was a significant difference ($p = 0.02$): the PPA group reported the highest percentage of some to full adherence to post-intervention exercise (NSE 69%, NSEB 60%, PPA 74%).

Table 3. Within-group results over time. Values are median (IQR)/ mean (SD) scores

	n	Baseline	6 months	12 months	24 months	P-value
NSE						
NDI	61	15 (12-21)/16 (6)	12 (7-19)/13 (8)	11 (8-19)/13 (8)	13 (8-20)/13 (8)	0.01
PSFS	62	4.3 (3.3-6.0)/4.4 (1.7)	7.0 (5.0-8.0)/6.3 (2.4)	7.0 (5.3-9)/6.7 (2.3)	6.8 (4.5-8.1)/6.3 (2.5)	<0.001
VAS Pain	62	38 (20-62)/40 (24)	18 (7-46)/27 (24)	15 (3-44)/25 (26)	17 (5-48)/ 25 (24)	<0.001
VAS Bothersomeness	59	49 (32-65)/49 (22)	27 (10-52)/32 (25)	24 (6-56)/32 (27)	23 (7-57)/33 (27)	0.001
SES	62	154 (130-174)/150 (34)	173 (138 -192)/165 (31)	173 (143-188)/160 (36)	161 (143-189)/ 158 (34)	0.02
NSEB						
NDI	66	17 (11-21)/17 (7)	13 (9-19)/13 (8)	12(7-17)/13 (8)	13(7-18)/13 (8)	<0.001
PSFS	63	4.3 (3.0-6.0)/4.5 (2.1)	6.8 (5.3-8.4)/6.7 (2.1)	7.0 (4.9 -8.7)/6.6 (2.5)	6.3 (4.2-7.7)/6.1 (2.5)	<0.001
VAS Pain	54	50 (19-65)/45 (24)	31 (7-49)/31 (23)	30 (7-57)/33 (27)	30 (7-51)/31 (25)	<0.01
VAS Bothersomeness	65	52 (31-70)/50 (23)	33 (17-52)/34 (23)	36 (10-61)/37 (27)	33 (9-59)/36 (27)	<0.001
SES	62	159 (127-184)/153 (35)	170 (134-189)/ 160 (33)	172 (149-196)/167 (31)	172 (143-192)/162 (37)	0.07
PPA						
NDI	55	17 (13-22)/17 (7)	16 (11-22)/16 (9)	17(8-22)/16 (8)	17(11-22)/17 (8)	0.13
PSFS	55	5.0 (3.3-5.7)/4.6 (1.8)	6.0 (4.0-7.3)/5.5 (2.5)	6.0 (4.0-7.3)/5.6 (2.2)	5.3 (3.7-7.0)/5.3 (2.2)	0.22
VAS Pain	52	42 (20-60)/42 (25)	23 (8-56)/32 (27)	27 (8-57)/32 (26)	27 (8-57)/31 (26)	0.07
VAS Bothersomeness	52	49 (30-67)/48 (22)	36 (18-61)/39 (27)	37 (14-64)/39 (25)	32 (16-57)/ 36 (26)	0.02
SES	56	157 (124-182)/147 (41)	159 (127-187)/149 (46)	156 (121-185)/147 (44)	153 (118-190)/ 147 (44)	0.86

NSE = neck-specific exercise group, NSEB = neck-specific exercise group with a behavioral approach, PPA = prescription of physical activity group, NDI = Neck Disability Index, PSFS = Pain Specific Functional Scale, VAS = Visual Analogue Scale, SES = Self-efficacy Scale.




Figure 2. Percentage of responders to treatment in the three intervention groups at the 1-year and 2-year follow-ups.

The proportion of responders with some to full adherence to post-intervention exercise that experienced a clinical important difference in the Neck Disability Index (NDI, cut-off change 5/50), Patient Specific Functional Scale (PSFS, cut-off change 2.3), Visual Analogue Scale-Bothersomeness (VAS-B, 50% reduction), Current Pain Visual Analogue Scale (VAS-P, 50% reduction). NSE=neck-specific exercise, NSEB=neck-specific exercise with a behavioural approach, PPA= Prescription of Physical Activity. Significant between-group differences: *P=0.02, **P<0.01.

Discussion

The results of this study indicate that a substantial number of individuals with chronic WAD (mean duration 20 months) can obtain long-lasting symptom reduction, particularly following neck-specific exercise. The two neck-specific exercise groups reported greater improvements regarding disability/functioning than the PPA group, which reported no improvement over time. Pain bothersomeness was reduced in all three groups over time, but current pain was only reduced in the NSE/NSEB groups, however there was no difference among mean group results. However, regarding clinically important improvement, defined as the proportion of participants reaching the MCID of the NDI, PSFS, or VAS, greater proportions of participants in the two neck-specific groups displayed clinically important improvements regarding pain, disability, and functioning at 1 year; the NSE group reported the greatest proportion (up to 61%). At 2 years, the between-group difference was only significant regarding disability, although the two neck-specific groups exhibited clear trends toward greater proportions of participants with clinically important improvement in all outcomes (40-59%), compared with the PPA group (12-33%). The lack of significance may be because of the lower number of participants in this subanalysis after excluding non-adherent patients. Neck-specific exercise did not produce any significant differences regarding any outcomes when applied alone versus in combination with a behavioural approach.

Our findings are in contrast with those of Michaleff et al. (15), who observed similar improvements for both treatment arms, whereas in our study the improvements seen in the PPA group were generally smaller or non-existing. Stewart et al. observed only short-term differences between groups (32). Possible reasons may be that Michaleff et al. included lower WAD grades (1-2) and did not include patients with WAD grade 3. WAD grade 3 has been associated with treatment success in neck-specific exercise (33). Disability improvements

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

also tended to be greater among patients with WAD grade 3 in the neck-specific groups, while there was a trend toward deterioration in the PPA group. The neck-specific exercise regimes may also have differed somewhat, as Stewart et al. (32) did not specify whether or not the endurance training was neck-specific. Although Michaleff et al. (15) reported progressive training in neck flexion/extension, it is unclear whether this included resistance. In our study, endurance training included head-resisted neck exercise, and apart from flexion and extension, rotation and lateral flexion were also part of the programme.

Consistent with our findings, the PSFS is reportedly more sensitive to change than the NDI (24). This difference in sensitivity might be because the chosen activities are based on what is most important to each individual. Therefore, the results must be interpreted as changes of functioning regarding meaningful individual activities for each participant. Even so, the PSFS is suggested to be an appropriate measure for statistical comparisons between groups in clinical research (34), and can also complement the NDI.

The proportion of participants that reported clinical improvement even after 2 years was high among those who continued exercising to some degree, especially in the two neck-specific groups. Whether this was because improvement led to greater motivation to continue exercising, or continuous exercise postulated improvement is uncertain. However, whether or not all participants were included in the analysis (regardless of adherence), the results were the same, except for a tendency toward lower improvements and a significant between-group difference favouring the two neck-specific groups regarding P-VAS in the complete set analysis.

Even though the PPA group reported better adherence to their prescribed exercise activity than the NSEB group at 2 years, the NSEB group still reported less disability. One reason

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

might be that the behavioural approach included the discussion of strategies to handle relapse/periods of worsening. This may also be a reason why the proportion of responders tended to increase over time. It may also indicate that choice of exercise intervention is important even after 2 years, even when continuous adherence is not optimal. This is in accordance with findings by Ylinen et al., who reported that improvements achieved through long-term training in chronic neck pain were maintained at a 3-year follow-up despite faltering adherence (35).

There were no improvements or between-group differences in self-efficacy at the 1- or 2-year follow-ups. There was a large variance that could not be explained by level of adherence. Self-efficacy has previously been reported to increase with the completion of an exercise program (36), however, there was no correlation between either baseline SES score or change in SES score and adherence (Spearman's ρ -0.10, -0.11, $p > 0.23$). One reason might be the relatively high baseline level of self-efficacy in this study, indicating that self-efficacy was not a major issue. This might be because participants were recruited mainly from primary care, not pain clinics. Patients in pain clinics reportedly have higher levels of functional impairment and psychosocial difficulties, but may thus not be as representative of individuals in general who suffer from chronic pain (37).

Opinions differ regarding whether rating scales and larger scores from questionnaires should be analyzed with non-parametric or parametric statistics. We also checked our results using parametric statistics, and observed no significant differences from the non-parametric results presented, in accordance with the policy of this journal.

Study limitations

Forty-three percent of the participants were lost to follow-up at 2 years, possibly because the clinical tests, in which they also participated for up to 1 year, ended, which may have reduced their motivation to continue answering questionnaires. This scenario introduces possible bias into the results. However, analysis indicated that the only difference between drop-outs and completers was that drop-outs at 1 year were somewhat younger than completers, and drop-outs at 2 years reported more baseline disability than completers. As reported previously, age was not associated with outcome; however, in this study sample a higher NDI score at baseline was associated with greater improvement regarding disability, but not pain (33). Furthermore, the use of imputed versus non-imputed data in the analyses did not significantly affect the results.

The closest match imputation method used in this study did not substitute values in cases in which only baseline data was available. This approach resulted in some missing values, even after imputation. However, closest match is reported to perform well even when missing data is substantial, and is recommended over common methods such as last value carried forward, which are reported to perform poorly (31). There was a small, but significant baseline difference between randomization groups regarding gender and age; however, neither factor was associated with outcomes in this study sample (33).

Although these results are very promising, and the study sample was representative regarding age, gender, and level of pain compared with those who declined study participation (16), they need to be repeated in another study population. It is also important for future studies to identify predictors of who will benefit from what specific treatments. As seen in this study, there were participants that exhibited clinically important improvement in all groups

Long-term follow-up of exercise in chronic whiplash, Landén Ludvigsson et al.

(including the PPA group, in which the mean disability rather tended to worsen slightly). In conclusion, after 1 to 2 years, participants with chronic WAD grade 2 or 3 who were randomized to neck-specific exercise (with or without a behavioural approach) remained more improved than participants who were prescribed general physical activity. However due to the loss to follow-up, the results need to be interpreted with some caution.

Acknowledgements

We wish to thank all participants in this study, including WAD participants, physiotherapists, and staff involved at any stage of the study.

Accepted manuscript JRM

References

1. Carroll LJ, Holm LW, Hogg-Johnson S, Cote P, Cassidy JD, Haldeman S, et al. Course and prognostic factors for neck pain in whiplash-associated disorders (WAD): results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)* 2008;33(4 Suppl):S83-92.
2. Verhagen AP, Scholten-Peeters GG, van Wijngaarden S, de Bie RA, Bierma-Zeinstra SM. Conservative treatments for whiplash. *Cochrane Database Syst Rev.* 2007:CD003338.
3. Hurwitz EL, Carragee EJ, van der Velde G, Carroll LJ, Nordin M, Guzman J, et al. Treatment of neck pain: noninvasive interventions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)* 2008;33(4 Suppl):S123-52.
4. Southerst D, Nordin MC, Cote P, Shearer HM, Varatharajan S, Yu H, et al. Is exercise effective for the management of neck pain and associated disorders or whiplash-associated disorders? A systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMA) Collaboration. *Spine J* 2014; (in press).
5. Soderlund A, Denison E. Classification of patients with whiplash associated disorders (WAD): reliable and valid subgroups based on the Multidimensional Pain Inventory (MPI-S). *Eur J Pain* 2006;10:113-9.
6. Chou R, Huffman LH. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med* 2007;147:492-504.
7. Nederhand MJ, Hermens HJ, MJ IJ, Turk DC, Zilvold G. Cervical muscle dysfunction in chronic whiplash-associated disorder grade 2: the relevance of the trauma. *Spine (Phila Pa 1976)* 2002;27:1056-61.
8. Sterling M, Jull G, Vicenzino B, Kenardy J, Darnell R. Development of motor system dysfunction following whiplash injury. *Pain* 2003;103:65-73.
9. Jull G, Kristjansson E, Dall'Alba P. Impairment in the cervical flexors: a comparison of whiplash and insidious onset neck pain patients. *Man Ther* 2004;9:89-94.
10. Woodhouse A, Vasseljen O. Altered motor control patterns in whiplash and chronic neck pain. *BMC Musculoskelet Disord* 2008;9:90.
11. Juul-Kristensen B, Clausen B, Ris I, Jensen RV, Steffensen RF, Chreiteh SS, et al. Increased neck muscle activity and impaired balance among females with whiplash-related chronic neck pain: a cross-sectional study. *J Rehabil Med* 2013;45:376-84.
12. Peterson G, Dederling A, Andersson E, Nilsson D, Trygg J, Peolsson M, et al. Altered ventral neck muscle deformation for individuals with whiplash associated disorder compared to healthy controls - A case-control ultrasound study. *Man Ther* 2015;20:319-27..
13. Teasell RW, McClure JA, Walton D, Pretty J, Salter K, Meyer M, et al. A research synthesis of therapeutic interventions for whiplash-associated disorder (WAD): part 4 - noninvasive interventions for chronic WAD. *Pain Res Manag* 2010;15:313-22.
14. Professional Associations for Physical Activity, Swedish National Institute of Public Health Physical activity in the prevention and treatment of disease. Stockholm: Swedish National Institute of Public Health; 2010. Available from: <http://www.fyss.se/fyss-in-english/>.
15. Michaleff ZA, Maher CG, Lin CW, Rebeck T, Jull G, Latimer J, et al. Comprehensive physiotherapy exercise programme or advice for chronic whiplash (PROMISE): a pragmatic randomised controlled trial. *Lancet* 2014 ;12;384:133-41 .
16. Ludvigsson ML, Peterson G, O'Leary S, Dederling A, Peolsson A. The Effect of Neck-specific Exercise With, or Without a Behavioral Approach, on Pain, Disability, and Self-Efficacy in Chronic Whiplash-associated Disorders: A Randomized Clinical Trial. *Clin J Pain* 2015;31:294-303.

17. Peolsson A, Ludvigsson ML, Overmeer T, Dederling A, Bernfort L, Johansson G, et al. Effects of neck-specific exercise with or without a behavioural approach in addition to prescribed physical activity for individuals with chronic whiplash-associated disorders: a prospective randomised study. *Bmc Musculoskel Dis* 2013;14:311.
18. Spitzer WO, Skovron ML, Salmi LR, Cassidy JD, Duranceau J, Suissa S, et al. Scientific monograph of the Quebec Task Force on Whiplash-Associated Disorders: redefining "whiplash" and its management. *Spine (Phila Pa 1976)* 1995;20(8 Suppl):1S-73S.
19. Vernon H. The Neck Disability Index: state-of-the-art, 1991-2008. *J Manipulative Physiol Ther* 2008;31:491-502.
20. Carlsson AM. Assessment of chronic pain. I. Aspects of the reliability and validity of the visual analogue scale. *Pain* 1983;16:87-101.
21. Landén Ludvigsson M, Peolsson A, Peterson G, 2015 [cited 2015-02-09]. Available from: <http://dx.doi.org/10.3384/report.diva-113865>
22. MacDermid JC, Walton DM, Avery S, Blanchard A, Etruw E, McAlpine C, et al. Measurement properties of the neck disability index: a systematic review. *J Orthop Sports Phys Ther* 2009;39:400-17.
23. Westaway MD, Stratford PW, Binkley JM. The patient-specific functional scale: validation of its use in persons with neck dysfunction. *J Orthop Sports Phys Ther* 1998;27:331-8.
24. Stewart M, Maher CG, Refshauge KM, Bogduk N, Nicholas M. Responsiveness of pain and disability measures for chronic whiplash. *Spine (Phila Pa 1976)* 2007;32:580-5.
25. Altmaier EM, Russell DW, Kao CF, Lehmann TR, Weinstein JN. Role of self-efficacy in rehabilitation outcome among chronic low back pain patients. *Journal of Counseling Psychology* 1993;40:335-9.
26. Denison E, Asenlof P, Lindberg P. Self-efficacy, fear avoidance, and pain intensity as predictors of disability in subacute and chronic musculoskeletal pain patients in primary health care. *Pain* 2004;111:245-52.
27. Bunketorp L, Carlsson J, Kowalski J, Stener-Victorin E. Evaluating the reliability of multi-item scales: a non-parametric approach to the ordered categorical structure of data collected with the Swedish version of the Tampa Scale for Kinesiophobia and the Self-Efficacy Scale. *J Rehabil Med* 2005;37:330-4.
28. Dworkin RH, Turk DC, McDermott MP, Peirce-Sandner S, Burke LB, Cowan P, et al. Interpreting the clinical importance of group differences in chronic pain clinical trials: IMMPACT recommendations. *Pain* 2009;146:238-44.
29. Cleland JA, Fritz JM, Whitman JM, Palmer JA. The reliability and construct validity of the Neck Disability Index and patient specific functional scale in patients with cervical radiculopathy. *Spine (Phila Pa 1976)* 2006;31:598-602.
30. Dworkin RH, Turk DC, Wyrwich KW, Beaton D, Cleeland CS, Farrar JT, et al. Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain* 2008;9:105-21.
31. Elliott P, Hawthorne G. Imputing missing repeated measures data: how should we proceed? *Aust N Z J Psychiatry* 2005;39:575-82.
32. Stewart MJ, Maher CG, Refshauge KM, Herbert RD, Bogduk N, Nicholas M. Randomized controlled trial of exercise for chronic whiplash-associated disorders. *Pain* 2007;128:59-68.
33. Landén Ludvigsson M, Peterson G, Dederling Å, Falla D, Peolsson A. Factors associated with pain and disability reduction following exercise interventions in chronic whiplash. *Eur J Pain* 2015 (in press, doi: 10.1002/ejp.729).
34. Abbott JH, Schmitt JS. The Patient-Specific Functional Scale was valid for group-level change comparisons and between-group discrimination. *J Clin Epidemiol* 2014;67:681-8.

35. Ylinen J, Hakkinen A, Nykanen M, Kautiainen H, Takala EP. Neck muscle training in the treatment of chronic neck pain: a three-year follow-up study. *Eura Medicophys* 2007;43:161-9.
36. Jones F, Harris P, Waller H, Coggins A. Adherence to an exercise prescription scheme: the role of expectations, self-efficacy, stage of change and psychological well-being. *Br J Health Psychol* 2005;10:359-78.
37. Mailis-Gagnon A, Yegneswaran B, Lakha SF, Nicholson K, Steiman AJ, Ng D, et al. Pain characteristics and demographics of patients attending a university-affiliated pain clinic in Toronto, Ontario. *Pain Res Manag* 2007;12:93-9.

Accepted manuscript JRM

Accepted manuscript JRM